



絶滅危惧種メダカに関する文献集

久米幸毅・久保喜計・細谷和海

近畿大学農学部水産学科

Bibliography on an endangerd fish, Medaka, *Oryzias latipes*

(Temmink & Schlegel)

Kouki Kume, Yoshikazu Kubo and Kazumi Hosoya

Department of Fisheries, Faculty of Agriculture, Kinki University,
3327-204 Nakamachi, Nara, 631-8505, Japan

Synopsis

Bibliography on Medaka fish, *Oryzias latipes*, was compiled as a list. Medaka is known as a paddy fish, whose wild populations have recently reduced. This is soon followed by that they are categorized as an endangered species in the Japanese Red Data Book (2003). Their decline in nature is chiefly due to the decrease of paddy fields suitable for their propagation, and to the separation of migratory pathway between paddy fields and ditches. On Medaka, much information has been accumulated so far as about experimental fish and ornamented fish. However, it lacks in basic biology on the wild Medaka. This bibliography lists scientific papers, articles of newspaper and other miscellaneous printed materials with specific order both in English and Japanese. This aims to provide necessary information to the persons such as researchers and citizen volunteers who willingly take part in the protection activities of the wild Medaka.

はしがき

メダカは流れのゆるやかな河川、小川に生息する卵生の淡水魚で、東アジアの熱帯から温帯域にかけ分布する。日本では特に親しまれ、属名に“稻”(*Oryza sativa*)に由来する*Oryziast*が用いられていることからも、水田環境を象徴する魚とされている。メダカは水温と光条件が満たされれば1年中産卵するほど繁殖力が旺盛で、しかも温度耐性が幅広く、塩分耐性も強い。このように、本種はもともと多様な水質環境に適応することができ、飼育が容易で、かつ非常に安価で入手することができるため、実験動物や観賞魚として頻繁に用いられてきた。

戦後、農業の効率化を目的とした大規模な圃場整備事業によって我が国の水田環境は一変し、野生メダカは激減してしまった。野生メダカは水田を繁殖場所・生育場所として積極的に利用することが知られており、その生活環を水田ネットワークに合わせ個体群を維持してきたと考えられている。そのため、生息地としての水田そのものの減少、水田と農業用水路との間の大きな段差、パイプライン化とともになうネットワークの分断は大きな影響を及ぼしていると思われる。

これを受け、旧環境庁は1999年にメダカを「絶滅危惧Ⅱ類」に指定した。各都道府県も地方版レッドデータブックにおいて本種を絶滅危惧種に位置づけている。それを契機に野生メダカ復活の機運は全国的に高まっているが、ヒメダカの放流で代表されるように、地域特性や系統を無視した無計画な放流が公然と行われている。そのため、生物多様性本来の目標にかなう施策を立案するためにも、野生メダカに関する正確な情報を数多く収集し、保護の担い手に提供することが急務となっている。

山本時男博士は1975年に総説書“The Medaka”を刊行されているが、これを含め、メダカに関する情報はこれまで発生学、遺伝学、生理学、薬理学などヒメダカを用いた実験室のデータに基づくものにかたより、野生メダカの生態に関する情報は絶対的に不足しているのが現状である。

本資料は野生メダカの保全に関する情報をはじめ、文献をさまざまな分野に整理したものである。ただし、本資料は発生工学や遺伝に関する情報提供を目的としていない。それらの分野に関しては、

名古屋大学ホームページのメダカデータベースを参照していただきたい。目録作成では原典に当たることを心がけたが、入手できなかった文献については孫引きによって収録した。文献出典の表示方法は、日本魚類学会刊行の魚類学雑誌最新号(51巻1号)に従った。目録は研究分野別に示し、最後に全文献を文献番号順に並べた。また、研究分野が多岐にわたるものでは重複して配置した。資料の編集は水産生物学研究室の北川忠生博士、本学部博士後期課程の藤田朝彦氏、平成14年度農学研究科水産学専攻の岡谷佳澄氏(現日本メンテナス株式会社環境試験所)、平成13年度水産生物学専攻生の朝井俊亘氏(現近畿大学付属新宮高校教諭)、“近大メダカの学校”的学生諸氏、特に八田実紀・松永京子・横田彰子さん、および森本昌平君の協力に負うところが大きい。なお、編集作業の一部は、財団法人リバーフロント整備センターに設置された外来種対策・検討委員会のケーススタディのための予算によった。合わせて謝意を表したい。本資料は1950年以降の文献を中心に整理したもので、発生や遺伝に関する資料はいまだ不完全であることは承知しているが、野生メダカに関する情報が不足している今日、それらを整理し社会に提供することは重要であると思われる。本資料が野生メダカを保護するための一助となることを願ってやまない。

分類(Taxonomy)

- Chen Y. R., H. Uwa and X. L. Chu. 1989. Taxonomy and distribution of the genus *Oryzias* in Yunnan China Cyprinodontiforms Oryziidae. *Acta Zootaxonomica Sinica*, 14: 239-246. [ref. 30]
- Kim, I-S and E-H. Lee. 1992. New record of ricefish, *Oryzias latipes sinensis* (Pisces Oryziidae) from Korea. *Korean Journal of Systematic Zoology*, 8(2): 177-182. [ref. 185]
- Matsuda, M., T. Yamagishi, M. Sakaizumi and S-R. Joen. 1997. Mitochondrial DNA variation in the Korean wild population of medaka, *Oryzias latipes*. *Korean J. Limnol*, 30(2): 119-128. [ref. 235]

- Parenti, L. R. and B. Soeroto. 2004. *Adrianichthys roseni* and *Oryzias nebulosus*, two new ricefishes (Atherinomorpha: Beloniformes: Adrianichtyidae) from Lake Poso, Sulawesi, Indonesia. Ichthyol. Res., 51(1): 10-19. [ref. 289]
- Rosen, D. and L. Parenti. 1981. Relationship of *Oryzias*, and the groups of atherinomorph fishes. American Museum of Natural History, 2719: 1-25. [ref. 296]
- Takehana, Y., S. Uchiyama, M. Sakaizumi, M. Matsuda and S-R. Jeon. 2004. Geographic variation and diversity of the cytochrome b gene in wild populations of medaka (*Oryzias latipes*) from Korea and China. Zool. Sci., 21(4): 483-491. [ref. 344]
- Uwa, H., R. F. Wang and Y. R. Chen. 1988. Karyotype and geographical distribution of ricefishes from Yunnan southwestern China. Japanese J. Ichthyol., 35(3): 332-340. [ref. 380]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]
- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 和泉克雄. 1971. 熱帯メダカ族百科. 東京書店, 357pp. [ref. 480]
- Egami, N. 1959. Note on sexual difference in the shape of the body in the fish, *Oryzias latipes*. Annot. Zool. Japon., 32: 59-64. [ref. 40]
- Foscarini, R. 1989. A comparative study of the skin and gill structure in oviparous and viviparous fresh water fish larvae. J. Fish. Biol., 34(1): 31-40. [ref. 54]
- Fukamachi, S., H. Mitani, A. Shima and M. Sugimoto. 2004. Somatolactin selectively regulates proliferation and morphogenesis of neural-crest derived pigment cells in medaka. Proc. Natl. Acad. Sci. U.S.A., 101(29): 10661-10666. [ref. 57]
- Ghoneum, M. and N. Egami. 1982. Age related changes in morphology of the thymus of the fish, *Oryzias latipes*. Exp. Gerontol., 17: 33-40. [ref. 62]
- Hamaguchi, S. 1985. Changes in the morphology of the germinal dense bodies in primordial germ cells of the teleost, *Oryzias latipes*. Cell Tissue Res., 240(3): 669-673. [ref. 77]
- Hamaguchi, S. 1996. Bilaterally asymmetrical testes in fishes of the genus *Oryzias*. Zool. Sci., 13(5): 757-763. [ref. 78]
- Hayashida, Y., T. Kawamura, R. Horie and I. Yamashita. 2004. Retionic acid and its receptors are required for expression of aryl hydrocarbon receptor mRNA and embryonic development of blood vessel and bone in the medaka fish, *Oryzias latipes*. Zool. Sci., 21(5): 541-551. [ref. 91]
- Hendreson, V. 1983. Scale morphologic response to aromatic hydrocarbons. Bull. Environ. Contam. Toxicol., 31(3): 315-321. [ref. 96]
- Hinton, D. E., E. R. Walker, C. A. Pinkstaff and E. M. Zuchelkowski. 1984. Morphological survey of teleost organs important in carcinogenesis with attention to fixation. Natl. Cancer Inst. Monogr., 65: 291-320. [ref. 100]
- Hirose, K. 1972. The ultrastructure of the ovarian follicle of medaka, *Oryzias latipes*. Z. Zellforsch. Mikrosk. Anat., 123(3): 316-329. [ref. 108]

形 態 (Morphology)

- Ali, M. Y. and C. C. Lindsey. 1974. Heritable and temperature-induced meristic in the medaka, *Oryzias latipes*. Can. J. Zool., 52(8): 959-976. [ref. 2]
- Benton, M. J., S. A. Diamond and S. I. Guttman. 1994. A genetic and morphometric comparison of *helisoma trivolvis* and *gambusia holbrooki* form clean and contaminated habitats. Ecotoxicol. Environ. Saf., 29(1): 20-37. [ref. 16]
- Bunton, T. E. 1991. Ultrastructure of hepatic hemangiopericytoma in the medaka (*Oryzias latipes*). Exp. Mol. Pathol., 54(2): 87-98. [ref. 23]

- Hirshfield, M., G. A. Kaufman and R. J. Beyers. 1972. Relationships of weight, length and body composition in the medaka, *Oryzias latipes*. Am. Midl. Nat., 88: 239-245. [ref. 109]
- Inohaya, K. and A. Kudo. 2002. Analysis of the vertebral ligament formation in living transgenic medaka. Zool. Sci., 19(12): 1450. [ref. 131]
- Inohaya, K., A. Kudo and T. Suzuki. 2003. Analysis of the vertebral ligament formation in the teleost, *Oryzias latipes*. Zool. Sci., 20(12): 1559. [ref. 132]
- Ishikawa, Y. 1990. Development of muscle nerve in the teleost fish, medaka. Neurosci. Res., Suppl(13): S152-S156. [ref. 138]
- Ishikawa, Y. 1994. Innervation of lateral line system in the medaka, *Oryzias latipes*. The Fish Biology journal MEDAKA, 6: 17-24. [ref. 139]
- Ishikawa, Y., T. Yasuda, K. Maruyama, T. Kage, N. Yamamoto, M. Yoshimoto and H. Ito. 2004. Axonogenesis in the medaka embryonic brain. J. Neurochemistry, 43 (2-3): 481. [ref. 140]
- Iwamatsu, T., K. Onitake, K. Matsuyama, M. Satoh and S. Yukawa. 1997. Effect of micropylar morphology and size on rapid sperm entry into the eggs of the medaka. Zool. Sci., 14(4): 623-628. [ref. 152]
- Iwata, A. and H. Uwa. 1981. Maintenance and growth of the horny processes of the medaka, *Oryzias latipes*, in vitro. Dev. Growth Differ., 23(3): 245-248. [ref. 159]
- Jaffe, L. F. 1986. Calcium and morphogenetic fields. Ciba. Found. Symp., 122: 271-288. [ref. 165]
- Kaneko, T. and S. Hasegawa. 1999. Application of laser scanning microscopy to morphological observations on drinking in freshwater medaka larvae and those exposed to 80% seawater. Fish Sci., 65(3): 492-493. [ref. 170]
- Katoh, F. and T. Kaneko. 2002. Effects of environmental Ca '2+' levels on branchial chloride cell morphology in freshwater-adapted killifish *Fundulus heteroclitus*. Fish Sci., 68(2): 347-355. [ref. 173]
- Kaufman, G. A. and R. J. Beyers. 1972. Relationships of weight, length and body composition in the medaka, *Oryzias latipes*. Am. Midl. Nat., 88: 239-245. [ref. 174]
- Khoo, G., V. P. E. Phang and T. M. Lim. 1992. The confocal laser scanning microscope as a tool for studying xanthophores of the swordtail (*Xiphophorus helleri*). Zool. Sci., 9(3): 665-669. [ref. 182]
- Kikuchi, S. and A. Imaizumi. 1983. A note on morphological changes in the thymus after rejection in the fish, *Oryzias latipes*. Zool. Mag., 92(3): 428-430. [ref. 183]
- Lindsey, C. C. 1965. The effect of alternating temperature on vertebral count in the medaka (*Oryzias latipes*). Can. J. Zool., 43: 99-104. [ref. 221]
- Nakamura, H., K. Ozato, Y. Wakamatsu and T. Iwamatsu. 2003. Normal growth of the "See-Through" medaka. Zool. Sci., 20(5): 607-615. [ref. 249]
- Ogawa, Y. 1958. On the local variation in the number of anal fin-rays of Japanese killifish, *Oryzias latipes* in Tajima district, Hyogo Prefecture. Collecting and Breeding, 20: 48-51. [ref. 268]
- Parenti, L. R. and B. Soeroto. 2004. *Adrianichthys roseni* and *Oryzias nebulosus*, two new ricefishes (Atherinomorpha: Beloniformes: Adrianichtyidae) from Lake Poso, Sulawesi, Indonesia. Ichthyol. Res., 51(1): 10-19. [ref. 289]
- Rocchetta, G., M-L. Vanelli and C. Pancaldi. 1983. Size and shape in *Poecilia reticulata*. Experientia, 39(4): 409-410. [ref. 295]
- Rosenthal, G. G. and C. S. Evans. 1998. Female preference for swords in *Xiphophorus helleri* reflects a bias for large apparent size. Proc. Natl. Acad. Sci. U.S.A., 95(8): 4431-4436. [ref. 297]
- Suzuki, A. and N. Shibata. 2004. Developmental

- process of genital ducts in the medaka, *Oryzias latipes*. Zool. Sci., 21(4): 397-406. [ref. 337]
- Takeuchi, K. 1967. Large tooth formation in female medaka, *Oryzias latipes*, given methyl testosterone. J. Dent., 46(4): 750. [ref. 351]
- Takeuchi, K. 1967. Normal development of teeth of medaka, *Oryzias latipes*. Aichi Gakuin Daigaku Shigakkaishi, 5(1): 81-83. [ref. 352]
- Takeuchi, K. 1968. Inhibition of large distal tooth formation in male medaka, *Oryzias latipes*, by estradiol. Experientia, 24(10): 1061-1062. [ref. 354]
- Uwa, H. and L. R. Parenti. 1988. Morphometric and meristic variation in ricefishes, genus *Oryzias*: A comparison with cytogenetic data. Ichthyol. Res., 35(2): 159-166. [ref. 379]
- Yamada, G., Y. Ogino, S. Ohta, Y. Sato, H. Ogi, H. Kato, M. Kamikawa, R. Haraguchi and K. Suzuki. 2003. Molecular analysis of external genitalia formation: morphogenesis for a copulatory organ. Biochemical J., 75(8): 695. [ref. 397]
- Yano, S. and H. Kaizu. 1954. Malformation in the medaka. Collecting and Breeding, 16: 213. [ref. 411]
- Yasutake, J., K. Inohaya and A. Kudo. 2003. Twist functions in the Medaka vertebral column formation. Zool. Sci., 20(12): 1559. [ref. 413]
- Yoshizaki, K. 1981. A morphological study on the melanophores of swordtail *Xiphophorus helleri*. Nippon Suisan Gakkaishi, 47(6): 747-751. [ref. 417]
- 青木 清・大木浩子. 1985. メダカの視覚と定位. 生物の科学 遺伝, 39(8): 37-38. [ref. 424]
- 江上信雄. 1953. メダカの尻鰭軟条数の変異に関する研究 I. 日本各地産野生メダカの軟条数の変異. 魚類学雑誌, 3(33-3): 87-89. [ref. 434]
- 江上信雄. 1953. メダカの尻鰭軟条数の変異に関する研究 II. 軟条数についての交配実験. 魚類学雑誌, 3: 171-178. [ref. 435]
- 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp. [ref. 436]
- 江上信雄・山上健次郎・嶋 昭紘. 1990. メダカの生物学. 東京大学出版会, 315pp. [ref. 437]
- 江上信雄・吉野道仁. 1954. メダカの尻鰭軟条数の変異に関する研究 III. 野生メダカ軟条数の地理的変異 (資料の追加). 魚類学雑誌, 7: 83-88. [ref. 438]
- 古川壇美・野呂晋一・長尾和右・井尻憲一・江口星雄. 1999. メダカ稚魚の成長に伴う視覚の発達および微小重力下での挙動. 宇宙生物科学, 13(3): 164-165. [ref. 446]
- 平岡幸夫. 1980. 水質汚濁の生物指標としてのメダカのせきつい骨異常 I 某農業かんがい用水路における経時的観察所見. 広島大学医学雑誌, 28(5): 611-618. [ref. 460]
- 岩切 稔・高尾立実・渡辺 清・中畠公宏. 1986. メダカうろこ色素胞の神経支配 II 光学顕微鏡と走査電子顕微鏡の対比による銀染色標本の観察. 福岡教育大学紀要 第3分冊 数学・理科・技術科編, 35(1985): 97-104. [ref. 472]
- 岩切 稔・坂中一恵・竹重宏子. 1987. メダカの形態的体色変化. 福岡教育大学紀要 第3分冊 数学・理科・技術科編, 36(1986): 63-72. [ref. 473]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]
- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp. [ref. 479]
- 松原喜代松. 1995. 魚類の形態と検索. 石崎書店, pp.367-370. [ref. 509]
- 水野利恵・井尻憲一. 1999. メダカ突然変異体haの発生・成長過程における耳石の観察. 宇宙生物科学, 13(3): 166-167. [ref. 515]
- 長尾和右・井尻憲一・西脇優子. 1999. 3Dクリノスタットによる疑似微小重力下におけるメダカはい発生 網膜および軟骨形成. 宇宙

- 生物科学, 13(3): 162-163. [ref. 520]
- 長屋雅人・工藤 明. 2001. 小型硬骨魚類(ゼブラフィッシュ/メダカ)を用いた再生研究. 遺伝子医学, 5(4): 653-658. [ref. 521]
- 瀬能 宏. 1993. カダヤシ科・メダカ科. 中坊 徹次(編) 日本産魚類検索 - 全種の同定. 東海大学出版会, pp.414-415. [ref. 555]
- 杉本雅純. 1998. 魚の色と模様 色素胞の数や形態の変化による体色変化 メダカの背地適応を例として. 海洋と生物, 20(6): 488-494, I. [ref. 567]
- 宇和ひろし. 1985. メダカ属の種と系統. 生物の科学 遺伝, 39(8): 6-11. [ref. 604]
- 籐本美孝・上野輝彌. 1984. メダカ *Oryzias latipes* の骨学的研究. 北九州市立自然史博物館誌, 5: 143-161. [ref. 613]

遺伝的分化 (Genetic diversification)

- Egami, N. 1954. Geographical variation in the male characters of the fish *Oryzias latipes*. Annot. Zool. Japon., 27: 7-12. [ref. 37]
- Kondo, R., S. Kaneko, H. Sun, S. I. Chigusa and M. Sakaizumi. 2002. Diversification of olfactory receptor genes in the Japanese medaka fish, *Oryzias latipes*. Gene, 282(1-2): 113-120. [ref. 206]
- Matsuda, M., H. Yonekawa, S. Hamaguchi and M. Sakaizumi. 1997. Geographic variation and diversity in the mitochondrial DNA of the medaka, *Oryzias latipes*, as determined by restriction endonuclease analysis. Zool. Sci., 14: 517-526. [ref. 233]
- Matsuda, M., T. Yamagishi, M. Sakaizumi and S-R. Jeon. 1997. Mitochondrial DNA variation in the Korean wild population of medaka, *Oryzias latipes*. Korean J. Limnol, 30(2): 119-128. [ref. 235]
- Sakaizumi, M., K. Moriwaki and N. Egami. 1983. Allozymic variation and regional differentiation in wild population of the fish *Oryzias latipes*. Copeia, 1983: 311-318. [ref. 305]
- Takehana, Y., S. Uchiyama, M. Sakaizumi, M. Matsuda and S-R. Jeon. 2004. Geographic

- variation and diversity of the cytochrome b gene in wild populations of medaka (*Oryzias latipes*) from Korea and China. Zool. Sci., 21(4): 483-491. [ref. 344]
- Takenaka, Y. and M. Sakaizumi. 2001. Geographic variation in the cytochrome b gene of the Japanese medaka, *Oryzias latipes*. Zool. Sci., 18(Supplement): 29. [ref. 347]
- Uwa, H. and L. R. Parentil. 1988. Morphometric and meristic variation in ricefishes, genus *Oryzias*: A comparison with cytogenetic data. Ichthyol. Res., 35(2): 159-166. [ref. 379]
- Uwa, H., R. F. Wang, Y. R. Chen. 1988. Karyotype and geographical distribution of ricefishes from Yunnan southwestern China. Japanese J. Ichthyol., 35(3): 332-340. [ref. 380]
- 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp. [ref. 436]
- 江上信雄・山上健次郎・嶋 昭絢. 1990. メダカの生物学. 東京大学出版会, 315pp. [ref. 437]
- 偏 寛. 1999. 生物多様性の分子機構へのアプローチ 脊索動物ゲノムの倍数化と多様性の分子機構 トランスポゾン関与の可能性. 蛋白質 核酸 酶素, 44(3): 234-244. [ref. 462]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]
- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp. [ref. 479]
- 太田克明・吉田元一・小野珠乙・山岸 哲. 1994. 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成5年度研究成果報告集2, pp.270-273. [ref. 537]
- 小澤祥司. 1999. 生きもの豊かな水田を再び. 水情報, 19(4): 8-11. [ref. 542]
- 小澤祥司. 2000. メダカが消える日 - 自然の再

- 生をめざして-. 岩波書店, 224 pp.
[ref. 543]
- 酒泉 満. 1987. メダカの分子生物地理学. 水野信彦・後藤 晃(編)日本の淡水魚類 - その分布, 変異, 種分化をめぐって-. 東海大学出版会, pp.81-90. [ref. 545]
- 酒泉 満. 1997. 淡水魚地方個体群の遺伝的特性と系統保存. 長田芳和・細谷和海(編)日本の希少淡水魚の現状と系統保存, よみがえれ日本産淡水魚. 緑書房, pp.218-227. [ref. 546]
- 酒泉 満. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してIV ゲノム・進化・種分化 メダカの系統と種内構造. 蛋白質 核酸 酵素, 45(17): 2909-2917. [ref. 547]
- 酒泉 満. 2002. メダカの遺伝的地域差と地域個体群の保全. 日本水環境学会セミナー, 43: 90-100. [ref. 550]
- 酒泉 満・久保孝一. 1981. カラー自然シリーズ35, メダカ. 偕成社, 38pp.[ref. 551]
- 竹花佑介・酒泉 満. 2002. 日本産希少淡水魚の保護 メダカの遺伝的多様性の危機. 生物の科学 遺伝, 56(6): 66-71. [ref. 576]
- 椿 宜高. 2000. 現代社会と遺伝学 生物多様性の危機と遺伝学. 生物の科学 遺伝, 54(10): 58-64. [ref. 590]
- 椿 宜高・高村健二・永田尚志・富山清升・太田克明・吉田元一・山岸 哲・河田雅圭・土田浩治. 1996. 野生生物の種の減少 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成7年度終了研究成果報告書4, 271-287. [ref. 591]
- 椿 宜高・高村健二・富山清升・太田克明・吉田元一・山岸 哲・粕谷栄一・河田雅圭・伊沢雅子. 1995. 野生生物の種の減少 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成6年度研究成果報告集2, pp.280-284. [ref. 592]
- 遺伝学一般 (General Genetics)
- Araki, K., H. Okamoto, I. Nakayama, H. Nagoya and A. C. Graveson. 2001. Analysis of haploid development based on expression patterns of developmental genes in the medaka *Oryzias latipes*. Dev. Growth Differ., 43(5): 591-599. [ref. 9]
- Benton, M. J., S. A. Diamond and S. I. Guttman. 1994. A genetic and morphometric comparison of *helisoma trivolvis* and *gambusia holbrooki* form clean and contaminated habitats. Ecotoxicol. Environ. Saf., 29(1): 20-37. [ref. 16]
- Dodd, J. M. 1960. Genetic and environmental aspects of sex determination in cold blooded vertebrates. Mem. Soc. Endocrinol., 7: 17-44. [ref. 36]
- Fukamachi, S., A. Shimada and A. Shima. 2002. Positional cloning of the responsible gene for medaka orangered variants. Zool. Sci., 19(12): 1420-1421. [ref. 55]
- Fukamachi, S., A. Shimada, K. Naruse and A. Shima. 2001. Genomic analysis of γ -ray-induced germ-cell mutations at the b locus recovered from the medaka specific-locus test. Mutat. Res., 458(1-2): 19-29. [ref. 56]
- Funayama, T., H. Mitani and A. Shima. 1993. Ultraviolet-induced DNA damage and its photorepair in tail fin cells of the medaka, *Oryzias latipes*. Photochemistry and Photobiology, 58(3): 380-385. [ref. 59]
- Furutani, M. and H. Kondoh. 2002. A genome-wide screen for mutations affecting embryonic pattern formation in the medaka, *Oryzias latipes*. Zool. Sci., 19(12): 1422. [ref. 60]
- Furutani, M. and J. Wittbrodt. 2004. Medaka and zebrafish, an evolutionary twin study. Mech. Dev., 121(7-8): 629-637. [ref. 61]
- Grabher, C., T. Henrich, A. Arenz, J. Wittbrodt, T. Sasado and M. Frutani. 2003. Transposon-mediated enhancer trapping in medaka. Gene, 322: 57-66. [ref. 66]
- Harumi, T. and T. Watanabe. 2003. Expression

- of membrane-bound and soluble guanylyl cyclase mRNAs in embryonic and adult retina of the medaka fish *Oryzias latipes*. Zool. Sci., 20(2): 133-140. [ref. 83]
- Harumi, T., T. Watanabe, T. Yamamoto, Y. Tanabe and N. Suzuki. 2002. Expression of particulate and soluble guanylyl cyclase mRNA in the medaka retina. Zool. Sci., 19(12): 1436. [ref. 84]
- Higuchi, T., S. Kitagawa, H. Hori and A. Kanamori. 2002. Cloning and expression of a meiosis-specific gene, DMC1, in medaka. Zool. Sci., 19(12): 1438. [ref. 99]
- Howard, R. D., J. A. Dewoody and W. M. Muir. 2004. Transgenic male mating advantage provides opportunity for Trojan gene effect in a fish. Proc. Natl. Acad. Sci. U.S.A., 101(9): 2934-2938. [ref. 119]
- Iida, A., H. Inagaki, M. Suzuki, Y. Wakamatsu, H. Hori and A. Koga. 2004. The tyrosinase gene of the i(b) albino mutant of the medaka fish carries a transposable element insertion in the promoter region. Pigment Cell Res., 17(2): 158-164. [ref. 125]
- Inagaki, H., A. Koga, Y. Bessho and H. Hori. 1998. The tyrosinase gene from medaka fish; transgenic expression rescues albino mutation. Pigment Cell Res., 11(5): 283-290. [ref. 128]
- Inagaki, H., Y. Bessho, A. Koga and H. Hori. 1994. Expression of the tyrosinase-encoding gene in a colorless melanophore mutant of the medaka fish, *Oryzias latipes*. Gene, 150: 319-324. [ref. 129]
- Itakura, T., S. C. Mahata, M. Hirano, H. Ochi and Y. Ogino. 2002. Cytochrome P450 1A1 cDNA from freshwater teleost medaka (*Oryzias latipes*) and induction of mRNA by 3-methylcholanthrene in various organs. Environ. Sci., 9(6): 463-473. [ref. 142]
- Ito, M., H. Nakata, O. Saitoh and M. Sugimoto. 2002. Molecular cloning of RGS proteins from the medaka (*Oryzias latipes*). Zool. Sci., 19(12): 1436. [ref. 143]
- Iwamatsu, T., A. Yusa, H. Kobayashi, M. Yamashita and Y. Shibata. 2003. Experimental hybridization among *Oryzias* species. II. karyogamy and abnormality of chromosome separation in the cleavage of interspecific hybrids between *Oryzias latipes* and *O. javanicus*. Zool. Sci., 20(11): 1381-1387. [ref. 150]
- Kanamori, A., H. Hori, K. Naruse, H. Mitani and A. Shima. 2003. Genomic organization of ZP domain containing egg envelope genes in medaka (*Oryzias latipes*). Gene, 305(1): 35-45. [ref. 167]
- Kawakami, K., A. Shima and N. Kawakami. 2000. Identification of a functional transposase of the Tol2 element, an Ac-like element from the Japanese medaka fish, and its transposition in the zebrafish germ lineage. Proc. Natl. Acad. Sci. U.S.A., 97(21): 11403-11408. [ref. 177]
- Kawakami, K., K. Imanaka, M. Itoh and M. Taira. 2004. Excision of the Tol2 transposable element of the medaka fish *Oryzias latipes* in *Xenopus laevis* and *Xenopus tropicalis*. Gene, 338(1): 93-98. [ref. 178]
- Kawakami, K. and T. Noda. 2004. Transposition of the Tol2 element, an Ac-like element from the Japanese medaka fish *Oryzias latipes*, in mouse embryonic stem cells. Genetics, 166(2): 895-899. [ref. 179]
- Kawashima, K., A. Koga, H. Hori and A. Shima. 1998. Excision of the Tol2 transposable element of the medaka fish, *Oryzias latipes* in zebrafish, *Danio rerio*. Gene, 255(1-2): 17-22. [ref. 181]
- Kim, I-C., Y. J. Kim, Y-D. Yoon, Y-S. Lee, J-S. Lee and S. Kawamura. 2004. Cloning of cytochrome P450 1A (CYP1A) genes from the hermaphrodite fish *Rivulus marmoratus* and the Japanese medaka *Oryzias latipes*. Mar. Environ. Res., 58(2-5): 125-129. [ref. 184]
- Kim, I-S and E-H. Lee. 1992. New record of ricefish, *Oryzias latipes sinensis* (Pisces Oryziidae) from Korea. Korean Journal of

- Systematic Zoology, 8(2): 177-182. [ref. 185]
- Kimura, A., M. Shinohara, R. Ohkura and T. Takahashi. 2001. Expression and localization of transcripts of MT5-MMP and its related MMP in the ovary of the medaka fish *Oryzias latipes*. Biochem. Biophys. Acta., 1518(1-2): 115-123. [ref. 186]
- Kinoshita, M., M. Yamauchi, M. Sasanuma, Y. Ishikawa, T. Osada, Y. Wadamatsu, K. Ozato and K. Inoue. 2003. A transgene and its expression profile are stably transmitted to offspring in transgenic medaka generated by the particle gun method. Zool. Sci., 20(7): 869-875. [ref. 187]
- Kinoshita, M., S. Kani, K. Ozato and Y. Wakamatsu. 2000. Activity of the medaka translation elongation factor 1 α -A promoter examined using the GFP gene as a reporter. Dev. Growth Differ., 42(5): 469-478. [ref. 188]
- Kitagawa, S., H. Hori and A. Kanamori. 2002. Expression of an oogenesis-specific gene Fig α during sex differentiation in medaka. Zool. Sci., 19(12): 1438. [ref. 190]
- Koga, A., A. Shimada, M. Sakaizumi, H. Tachida and H. Hori. 2000. Evidence for recent invasion of the medaka fish genome by the Tol2 transposable element. Genetics, 155(1): 273-281. [ref. 195]
- Koga, A. and H. Hori. 1999. Homogeneity in the structure of the medaka fish transposable element Tol2. Genet. Res., 73: 7-14. [ref. 196]
- Koga, A. and H. Hori. 2000. Detection of de novo insertion of the medaka fish transposable element tol2. Genetics, 156(3): 1243-1247. [ref. 197]
- Koga, A. and H. Hori. 2001. The Tol2 transposable element of the medaka fish: an active DNA-based element naturally occurring in a vertebrate genome. (Review), Genes. Genet. Syst., 76(1): 1-8. [ref. 198]
- Koga, A., H. Hori and M. Sakaizumi. 2002. Transposable elements in medaka fish. Zool. Sci., 19(1): 1-6. [ref. 199]
- Koga, A., H. Hori and M. Sakaizumi. 2002. Gene transfer and cloning of flanking chromosomal regions using the medaka fish Tol2 transposable element. Mar. Biotechnol., 4(1): 6-11. [ref. 200]
- Koga, A., H. Inagaki, Y. Bessho and H. Hori. 1994. Body color mutants and atransposable element affecting the tyrosinase gene of the medaka fish, *Oryzias latipes*. (Review Article), The Fish Biology Journal MEDAKA, 6: 7-15. [ref. 201]
- Koga, A., H. Inagaki, Y. Bessho and H. Hori. 1995. Insertion of a novel transposable element in the tyrosinase gene is responsible for an albino mutation in the medaka fish, *Oryzias latipes*. Mol. Gen. Genet., 249: 400-405. [ref. 202]
- Koga, A., M. Suzuki, Y. Maruyama, M. Tsutsumi and H. Hori. 1999. Amino acid sequence of a putative transposase protein of the medaka fish transposable element Tol2 deduced from mRNA nucleotide sequences. FEBS Lett., 461(3): 295-298. [ref. 203]
- Koga, A., Y. Wakamatsu, J. Kurosawa and H. Hori. 1999. Oculocutaneous albinism in the i6 mutant of the medaka fish is associated with a deletion in the tyrosinase gene. Pigment Cell Res., 12: 252-258. [ref. 204]
- Kurita, J., T. Oshiro, F. Takashima and M. Sakaizumi. 1995. Cytogenetic studies on diploid and triploid oogenesis in interspecific hybrid fish between *Oryzias latipes* and *O. ocurvinotus*. J. Exp. Zool., 273(3): 234-241. [ref. 211]
- Kurosawa, G., K. Yamada, H. Ishiguro and H. Hori. 1999. Hox gene complexity in medaka fish may be similar to that in pufferfish rather than zebrafish. Biochem. Biophys. Res. Comm., 260: 66-77. [ref. 213]
- Kurosawa, G., N. Takamatsu, H. Hori, K. Yamada, H. Ishiguro, M. Matuda, S.

- Asakawa, N., Shimizu and Y. Kohara. 2002. The genome structure and the genomic comparison of medaka hox cluster. Zool. Sci., 19(12): 1418. [ref. 214]
- Lee, C., J. G. NA, K-C. Lee and K. Park. 2002. Choriogenin mRNA induction in male medaka, *Oryzias latipes* as a biomarker of endocrine disruption. Aquatic Toxicol., 61(3-4): 233-241. [ref. 217]
- Lee, C., S. H. Jeon, N. A. J. Gyun and K. Park. 2002. Sequence analysis of choriogenin H gene of medaka (*Oryzias latipes*) and mRNA expression. Environ. Toxicol. Chem., 21(8): 1709-1714. [ref. 218]
- Martinez-Morales, J. R., K. Naruse, H. Mitani, A. Shima and J. Wittbrodt. 2004. Rapid chromosomal assignment of medaka mutants by bulked segregant analysis. Gene, 329: 159-165. [ref. 224]
- Maruyama, K., S. Yasumasu and I. Iuchi. 2002. Analysis of globin gene expression in the medaka, *Oryzias latipes*. Zool. Sci., 19(12): 1499. [ref. 227]
- Maruyama, K., S. Yasumasu, I. Iuchi and K. Naruse. 2004. Globin gene organization in the medaka, *Oryzias latipes*. Zool. Sci., 18(Supplement): 27. [ref. 228]
- Maruyama, K., S. Yasumasu, K. Naruse, H. Mitani, A. Shima and I. Iuchi. 2004. Genomic organization and developmental expression of globin genes in the teleost *Oryzias latipes*. Gene, 335: 89-100. [ref. 229]
- Matsuda, M. 2002. Identification of sex determining gene in the medaka-DMY is a Y-specific DM-domain gene required for male development. Zool. Sci., 19(12): 1421. [ref. 231]
- Matsuda, M., C. Matsuda, Y. Nagahama, S. Hamaguchi and M. Sakaizumi. 2001. Positional cloning of the sex-determining region of medaka using a Y congenic strain. Zool. Sci., 18(Supplement): 27. [ref. 232]
- Matsuda, M., H. Yonekawa, S. Hamaguchi and M. Sakaizumi. 1997. Geographic variation and diversity in the mitochondrial DNA of the medaka, *Oryzias latipes*, as determined by restriction endonuclease analysis. Zool. Sci., 14: 517-526. [ref. 233]
- Matsuda, M., N. Kawamoto, S. Asakawa, N. Shimizu, Y. Nagahama, S. Hamaguchi, M. Sakaizumi and H. Hori. 2001. Construction of a BCA library derived from the inbred Hd-Rr strain of the teleost fish, *Oryzias latipes*. Genes Genet. Syst., 76(1): 61-63. [ref. 234]
- Matsuda, M., T. Yamagishi, M. Sakaizumi and S-R. Joen. 1997. Mitochondrial DNA variation in the Korean wild population of medaka, *Oryzias latipes*. Korean J. Limnol., 30(2): 119-128. [ref. 235]
- Matsuda, M., Y. Nagahama, A. Shinomiya, S. Hamaguchi, M. Sakaizumi, N. Shibata, S. Asakawa, N. Shimizu and H. Hori. 2002. DMY is a Y-specific DM-domain gene required for male development in the medaka fish. Nature, 417(6888): 559-563. [ref. 236]
- Matsuda, M., Y. Nagahama, T. Sato, Y. Toyozaki, S. Hamaguchi and M. Sakaizumi. 2003. *Oryzias curvinotus* has DMY, a gene that is required for male development in the Medaka, *O. latipes*. Zool. Sci., 20(2): 159-161. [ref. 237]
- Matsuzawa, T. and J. Hamilton. 1973. Polymorphism in lactate dehydrogenase of skeletal muscle associated with YY sex chromosomes in medaka (*Oryzias latipes*). Proc. Soc. Exp. Biol. Med., 142(1): 232-236. [ref. 238]
- Nakatsuru, Y., N. Nemoto, K. Nakagawa, P. Masahito and T. Ishikawa. 1987. O6-methylguanine DNA methyltransferase activity in liver from various fish species. Carcinogenesis, 8(8): 1123-1127. [ref. 250]
- Nanda, I., M. Kondo, U. Hornung, C. Winkler, M. Schartl, S. Asakawa, Z. Shan, T. Haaf and A. Shima. 2002. A duplicated copy of DMRT1 in the sex-determining region of the Y chromosome of the medaka, *Oryzias*

- latipes*. Proc. Natl. Acad. Sci. U.S.A., 99(18): 11778-11783. [ref. 253]
- Naruse, K., H. Hori, N. Shimizu, Y. Kohara and H. Takeda. 2004. Medaka genomics: a bridge between mutant phenotype and gene function. Mech. Dev., 121(7-8): 619-628. [ref. 254]
- Naruse, K., H. Mitani, A. Shima, M. Tanaka and K. Mita. 2002. Medaka gene map and the vertebrate genome evolution. Zool. Sci., 19(12): 1496. [ref. 255]
- Naruse, K., H. Mitani, S. Kondo, A. Shima and M. Tanaka. 2001. Medaka EST analysis, gene mapping and the vertebrate genome evolution. Zool. Sci., 18(Supplement): 27. [ref. 256]
- Naruse, K., K. Ijiri, A. Shima and N. Egami. 1985. The production of cloned fish in the medaka (*Oryzias latipes*). J. Exp. Zool., 236(3): 335-341. [ref. 257]
- Ogawa, N. 1965. A phenogenetic study of the vertebral fused (f) in the medaka, *Oryzias latipes*. Embryologia (Nagoya), 9(1): 13-33. [ref. 267]
- Ogiwara, K., M. Shinohara and T. Takahashi. 2004. Expression of proprotein convertase 2 mRNA in the ovarian follicles of the medaka, *Oryzias latipes*. Gene, 337: 79-89. [ref. 270]
- Ohtsuka, M., N. Kikuchi, K. Ozato, H. Inoko and M. Kimura. 2004. Comparative analysis of a 229-kb medaka genomic region, containing the zic1 and zic4 genes, with Fugu, human, and mouse. Genomics, 83(6): 1063-1071. [ref. 272]
- Okubo, K., K. Naruse, K. Aida, H. Mitani, M. Kondo, A. Shima, M. Tanaka, S. Asakawa and N. Shimizu. 2002. Structural characterization of GnRH loci in the medaka genome. Gene, 293(1-2): 181-189. [ref. 278]
- Philipp, D. and G. Whitt. 1977. Patterns of gene expression during teleost embryogenesis: lactate dehydrogenase isozyme ontogeny in the medaka (*Oryzias latipes*). Dev. Biol., 59(2): 183-197. [ref. 291]
- Rotchell, J. M., G. K. Ostrander, E. Ulnal and R. J. Van Beneden. 2001. Retinoblastoma gene mutations in chemically induced liver tumor samples of Japanese medaka (*Oryzias latipes*). Mar. Biotechnol., 3(Supplement): S44-S49. [ref. 298]
- Rotchell, J. M., G. K. Ostrander, J. B. Brair, J.-K. Shin and W. E. Hawkins. 2001. Cloning of the retinoblastoma cDNA from the Japanese medaka (*Oryzias latipes*) and preliminary evidence of mutational alterations in chemically induced retinoblastomas. Gene, 263(1-2): 231-237. [ref. 299]
- Sakaizumi, M., K. Moriwaki and N. Egami. 1983. Allozymic variation and regional differentiation in wild population of the fish *Oryzias latipes*. Copeia, 1983: 311-318. [ref. 305]
- Sakaizumi, M., M. Takeda, H. Otake, K. Togashi, A. Shinomiya, S. Hamaguchi, M. Matsuda and Y. Nagahama. 2002. Mutations affecting sex determination or differentiation from wild populations of the medaka, *Oryzias latipes*. Zool. Sci., 19(12): 1497. [ref. 306]
- Sato, T., M. Sakaizumi and S. Hamaguchi. 2002. Genetic analysis of a secondary sexual character-less mutant, scl, the medaka, *Oryzias latipes*. Zool. Sci., 19(12): 1488. [ref. 309]
- Schartl, M. 2004. A comparative view on sex determination in medaka. Mech. Dev., 121(7-8): 639-645. [ref. 310]
- Scholz, S., C. Kordes, J. Hamann and H. O. Gutzeit. 2004. Induction of vitellogenin in vivo and in vitro in the model teleost medaka (*Oryzias latipes*): comparison of gene expression and protein levels. Mar. Environ. Res., 57(3): 235-244. [ref. 312]
- Shima, A. and A. Shimada. 1988. Induction of mutations in male of the fish *Oryzias latipes* at a specific locus after gamma-irradiation. Mutat. Res., 198(1): 93-98. [ref. 317]

- Shimada, A and A. Shima. 2004. Transgenerational genomic instability as revealed by a somatic mutation assay using the medaka fish. *Mutat. Res.*, 552(1-2): 119-124. [ref. 322]
- Shimada, A., A. Shima, S. Fukumachi, Y. Wakamatsu and K. Ozato. 2002. Induction and characterization of mutations at the b locus of the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(4): 411-417. [ref. 323]
- Shinomiya, A., H. Otake, K. Togashi, S. Hamaguchi and M. Sakaizumi. 2004. Field survey of sex-reversals in the medaka, *Oryzias latipes*; genotypic sexing of wild populations. *Zool. Sci.*, 21(6): 613-619. [ref. 327]
- Shinomiya, A., S. Hamaguchi, M. Sakaizumi, M. Matsuda and Y. Nagahama. 2002. A strain having a ZZ/ZW sex determination system derived from a naturally occurring recessive mutation in the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(12): 1497. [ref. 328]
- Sun, H., R. Kondo, A. Shima, K. Naruse, H. Hori and S. I. Chigusa. 1999. Evolutionary analysis of putative olfactory receptor genes of medaka fish, *Oryzias latipes*. *Gene*, 231(1-2): 137-145. [ref. 335]
- Suzuki, A., N. Shibata, T. Sato, M. Sakaizumi and S. Hamaguchi. 2002. Development of gonads and genital ducts in secondary sexual character less mutant of medaka. *Zool. Sci.*, 19(12): 1488. [ref. 338]
- Takai, A., N. Kagawa and K. Fujikawa. 2004. Does- and time-dependent responses for micronucleus induction by X-rays and fast neutrons in gill cells of medaka. *Environ. Mol. Mutagen.*, 44(2): 108-112. [ref. 342]
- Takai, A., N. Kagawa and K. Fujikawa. 2004. Susceptibility of male and female medaka (*Oryzias latipes*) fish to spontaneous and X-ray induced micronucleus formation in gill cells. *Mutat. Res.*, 558(1-2): 131-136. [ref. 343]
- Takehana, Y., S. Uchiyama, M. Sakaizumi, M. Matsuda and S-R. Jeon. 2004. Geographic variation and diversity of the cytochrome b gene in wild population of medaka (*Oryzias latipes*) from Korea and China. *Zool. Sci.*, 21(4): 483-491. [ref. 344]
- Takemoto, T., H. Kimura, M. Nonaka, A. Shimada, A. Shima and S. Yanagi. 2001. Functional investigation of the DRP/CRAM family in medaka fish, *Oryzias latipes*. *Zool. Sci.*, 18(Supplement): 70. [ref. 345]
- Takenaka, Y. and M. Sakaizumi. 2001. Geographic variation in the cytochrome b gene of the Japanese medaka, *Oryzias latipes*. *Zool. Sci.*, 18(Supplement): 29. [ref. 347]
- Takenaka, Y. and M. Sakaizumi. 2002. Phylogeography of medaka (*Oryzias latipes*) based on mitochondrial cytochrome b DNA sequences. *Zool. Sci.*, 19(12): 1497. [ref. 348]
- Takeuchi, K. 1960. A study of the mutant (wavy) in the medaka, *Oryzias latipes*. *Annot. Zool. Japon.*, 33: 124-131. [ref. 349]
- Takeuchi, K. 1966. 'Wavy-fused' mutants in the Medaka, *Oryzias latipes*. *Nature*, 211(51): 866-867. [ref. 350]
- Tong, Y., T. Shan, Y. K. Poh, T. Yan, H. Wang, S. H. Lam and Z. Gong. 2004. Molecular cloning of zebrafish and medaka vitellogenin genes and comparison of their expression in response to 17beta-estradiol. *Gene*, 328: 25-36. [ref. 364]
- Toyozaki, Y., M. Sakaizumi and S. Hamaguchi. 2002. Three species closely related to *Oryzias latipes* have a male-heterogametic (XX-XY) sex-determining system. *Zool. Sci.*, 19(12): 1496. [ref. 365]
- Walter, R., H-Y. Li, S. Kazianis, G. W. Intano and C. A. Walter. 2002. Absence of global genomic cytosine methylation pattern erasure during medaka (*Oryzias latipes*) early embryo development. *Comp. Biochem. Physiol. B.*, 133B(4): 597-607. [ref. 383]
- Winkler, C., U. Hornung, M. Kondo, C. Neuner, J. Duschl, A. Shima and M. Schartl. 2004.

- Developmentally regulated and non-sex-specific expression of autosomal dmrt genes in embryos of the Medaka fish (*Oryzias latipes*). *Mech. Dev.*, 121(7-8): 997-1005. [ref. 392]
- Yamamoto, T. 1964. The problem of viability of YY zygotes in the medaka, *Oryzias latipes*. *Genetics*, 50: 45-58. [ref. 404]
- Yamamoto, T. 1964. Linkage map of sex chromosomes in the medaka, *Oryzias latipes*. *Genetics*, 50: 59-64. [ref. 405]
- Yamamoto, T. 1965. Estriol-induced XY females of the medaka (*Oryzias latipes*) and their progenies. *Gen. Comp. Endocrinol.*, 5(5): 527-533. [ref. 406]
- Yamamoto, T. 1967. Estrone-induced white YY females and mass production of white YY males in the medaka, *Oryzias latipes*. *Genetics*, 55(2): 329-336. [ref. 407]
- 安藤正典・西村哲治・神野透人・鈴木玲央・青木康展・有蘭幸司・石橋弘志・今川正良・中室克彦. 2004. 水域環境における内分泌かく乱化学物質の次世代への影響評価法確立に関する分子遺伝学的研究. 環境保全研究成果集, 2002 (1): 11.1-11.20. [ref. 422]
- 青木 宙・隆島史夫・平野哲也. 1997. 魚類のDNA-分子遺伝学的アプローチ. 恒星社厚生閣, 467pp. [ref. 425]
- 土井敏男. 2001. 閉鎖的ビオトープで見られたメダカの体型変異. 兵庫陸水生物, (53): 73-79. [ref. 433]
- 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp. [ref. 436]
- 江上信雄・山上健次郎・嶋 昭紘. 1990. メダカの生物学. 東京大学出版会, 315pp. [ref. 437]
- 平山 真. 2003. 透明メダカ現る!. 化学と生物, 41(2): 110-111. [ref. 461]
- 猪早敬二・工藤 明. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅠ. パターン形成と器官形成骨形成モデルとしてのメダカ. 蛋白質核酸 酶素, 45(17): 2745-2751. [ref. 469]
- 石川祐二・荒木和男. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカにおけるENUおよびX線による誘発突然変異体スクリーニング. 蛋白質核酸 酶素, 45(17): 2820-2828. [ref. 470]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]
- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp. [ref. 479]
- 古賀章彦. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝進化の理解を目指してⅣ・ゲノム・進化・種分化 メダカ属でのトランスポゾンの進化. 蛋白質核酸 酶素, 45(17): 2900-2908. [ref. 501]
- 松田 勝・長浜嘉孝. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカBACライブラリーの構築. 蛋白質核酸 酶素, 45(17): 2880-2885. [ref. 510]
- 三谷啓志. 2004. メダカから脊椎動物ゲノムの痕跡をたどる フグやゼブラフィッシュとの比較から何が見えてくるか?. 化学と生物, 42(4): 221-223. [ref. 512]
- 成瀬 清. 1985. 人為雌性発生によるクローンメダカの作成. 生物の科学 遺伝, 39(8): 22-24. [ref. 527]
- 成瀬 清・近藤 周・三谷啓志・嶋 昭紘・深町昌司・近藤真理子. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカ連鎖地図からみえるもの ポジショナルクローニング・比較マッピングからゲノム進化まで. 蛋白質核酸 酶素, 45(17): 2844-2852. [ref. 528]
- 大塚正人・木村 穂. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカ背腹構造突然変異体Daの原因遺伝子のポジショナルクローニング. 蛋白質核酸 酶素, 45(17): 2864-2871. [ref. 540]

- 尾里健二郎・若松佑子・依田欣哉. 1996. メダカと遺伝子. 平成8年度社会人公開実験講座96 “バイオサイエンスを体験する” 報告書, pp.12-32. [ref. 541]
- 酒泉 満. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してIV. ゲノム・進化・種分化 メダカの系統と種内構造. 蛋白質 核酸 酶素, 45(17): 2909-2917. [ref. 547]
- 酒泉 満. 2000. 分子情報に基づくメダカの種内構造の解析. 日本植物分類学会会報, 15(2): 27-36. [ref. 548]
- 酒泉 満. 2002. メダカの遺伝的地域差と地域個体群の保全. 日本水環境学会セミナー, 43: 90-100. [ref. 550]
- 嶋 昭紘. 2004. メダカの生物学研究最前線. 生物の化学 遺伝, 58(5): 89-92. [ref. 557]
- 嶋 昭紘・桑原義和・江口星雄. 2003. メダカ雄性殖細胞における γ 線誘発ゲノム損傷排除機構の解析. 共用設備管理部門年報, 30: 21-24. [ref. 558]
- 島田敦子・嶋 昭紘. 2000. メダカを用いた生殖細胞突然変異成立機構の解析 生殖系列における遺伝的不安定性. 放射線生物研究, 35(4): 370-382. [ref. 563]
- 島田敦子・嶋 昭紘. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してIII. 突然変異体作製・マッピング・クローニング メダカにおける自然および誘发生殖細胞突然変異率とその系統差. 蛋白質 核酸 酶素, 45(17): 2886-2892. [ref. 564]
- 田口泰子. 1985. メダカの近交系の作出とその応用. 生物の科学 遺伝, 39(8): 18-21. [ref. 572]
- 竹花佑介・酒泉 満. 2002. 日本産希少淡水魚の保護 メダカの遺伝的多様性の危機. 生物の科学 遺伝, 56(6): 66-71. [ref. 576]
- 田中 実・木下政人. 2003. メダカの細胞ひとつひとつを光らせて生殖腺形成機構を解明する 生殖細胞系列特異的にRNA翻訳制御を指示するエレメントがわかった. 化学と生物, 41(5): 278-280. [ref. 584]
- 田中 実・友永健士・木下政人・若松祐子. 2003. 遺伝子機能解析のためのポストゲノム実験動物メダカ (生きたまま内臓(細胞)を見られる「透明・遺伝子導入」メダカ). 生物物理, 43(2): 93-96. [ref. 585]
- 椿 宜高. 2000. 現代社会と遺伝学 生物多様性の危機と遺伝学. 生物の科学 遺伝, 54(10): 58-64. [ref. 590]
- 若松祐子・丹羽勝利・可児修一・尾里健二郎. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してV. 新技術の展開 メダカの核移植. 蛋白質 核酸 酶素, 45(17): 2962-2966. [ref. 606]
- 保全 (Conservation)
- Hata, K. 2002. Field experiment on the migration of fishes to paddy field with a small fishway. JARQ., 36(4): 219-225. [ref. 85]
- Kashiwaba, S., H. Ishikawa, N. Miyamoto, Y. Ohnishi and Y. Magara. 2002. Fish test for endocrine-disruption and estimation of water quality of Japanese rivers. Water Res., 36(8): 2161-2166. [ref. 171]
- Kobza, R. M., J. C. Trexler, W. F. Loftus and S. A. Perry. 2004. Community structure of fishes inhabiting aquatic refuges in a threatened Karst wetland and its implications for ecosystem management. Biol. Conserv., 116(2): 153-165. [ref. 194]
- Taneda, K., T. Nishimoto and R. Abe. 1996. The sensory cue for making school in Medaka, *Oryzias latipes*. Mem. Fac. Sci. Kochi Univ. Series d Biology, 16-17(0): 69-75. [ref. 357]
- 青井 透. 2001. 研究現場からの最新報告 (6) 水の循環利用と豊かな生態系復元を目的とするビオトープ創り. 環境施設, (83): 81-87. [ref. 423]
- 大日本水産会. 2000. 藤沢メダカの学校をつくる会. SAKANA, Vol.10. [ref. 429]
- 大日本水産会. 2001. 藤沢メダカの学校をつくる会. SAKANA, Vol.11. [ref. 430]
- 遠藤 徹. 2004. メダカの里「家根合地区」－水田生態系保全の取り組みと地域づくり－.

- ほ場と土壤, 36(2): 29-34. [ref. 440]
- 福田博道. 2001. メダカを川に戻したい. FRONT, 7月号. [ref. 443]
- 端 憲二. 1999. 近自然工法4 小さな魚道による休耕田への魚類遡上試験. 農業土木学会誌, 67(5): 497-502. [ref. 451]
- 端 憲二. 2000. 魚道の評価 田圃につける小さな魚道. 応用生態工学, 3(2): 231-234. [ref. 452]
- 端 憲二・本間新哉・中 達男. 2001. 流速増大時のメダカの避難行動. 農業工学関係研究成果情報, 2000: 13-14. [ref. 453]
- 堀 由紀子. 1998. 水族館のはなし. 岩波書店, 200pp. [ref. 465]
- 細谷和海. 2000. メダカの生息状況と保護. 水環境学会誌, 23(3): 7-11. [ref. 466]
- 市川憲平. 2004. 放棄田ビオトープによる里の自然再生とタガメやその他の水生動物の定着. ホシザキグリーン財団研究報告, (7): 137-150. [ref. 468]
- 岩切 稔・神野展光・長尾泰之・小寺智暉. 1994. 福岡県宗像市釣川水系産魚類IV. 宗像地域におけるメダカの分布. 福岡教育大学紀要 第3分冊 数学・理科・技術科編, (43): 65-74. [ref. 471]
- JICA. 2000. タイメダカ公開セレモニー. JICAフロンティア, 7月. [ref. 481]
- 城条義興. 2001. 藤沢メダカの学校をつくる会アドバイザー. 小学館, 小四教育技術, 7-8号. [ref. 482]
- 神奈川県公害センター. 1984. 神奈川県の水生生物. 神奈川県公害センター. [ref. 483]
- 神奈川県立生命の星・地球博物館. 1995. 神奈川県レッドデータ生物調査報告書, 神奈川県立博物館調査研究報告, 第7号, 神奈川県レッドデータ生物調査団編集. [ref. 484]
- 環境庁. 1999. 汽水・淡水魚類のレッドリストの見直しについて. 環境庁自然保護局野生生物課, pp.1-6. [ref. 487]
- 環境庁自然保護局. 1997. 第4回自然環境保全基礎調査(1993年度版), 動植物分布調査報告書, 淡水魚類. pp.146, 342. [ref. 488]
- 環境教育部会. 2001. 水のしらべ—藤沢の水と奏でる環境教育. 藤沢市教育文化センター. [ref. 489]
- 環境省. 2003. 改訂・日本の絶滅のおそれのある野生生物 - レッドデータブック (汽水・淡水魚類編). 自然環境研究センター, pp.162-163. [ref. 490]
- 環境省 自然環境局. 2004. 農林水産省と環境省の連携による「田んぼの生きもの調査2003」の結果について. かんきょう(東京), 29(6): 38-39. [ref. 491]
- 川地啓文. 2001. 藤沢メダカと環境教育. ふじさわ教育, 第126号. 藤沢市教育文化センター. [ref. 493]
- 建設省. 1998. 平成2~7年度河川水辺の国勢調査(河川版), 生物調査結果総括. 建設省河川局河川環境課, pp.25-26. [ref. 494]
- 建設省. 1998. 平成2~7年度河川水辺の国勢調査(河川版), 生物調査結果総括(参考資料). 建設省河川局河川環境課. [ref. 495]
- 菊池久登. 1996. 再発見された藤沢メダカ. ふじさわ教育, 第108号. 藤沢市教育文化センター. [ref. 496]
- 上月康則・村上仁士・森 裕行・佐藤陽一. 1998. 都市近郊河川におけるメダカを中心とした魚類相調査. 土木学会年次学術講演会公演概要集, 53: 492-493. [ref. 504]
- 松本清志. 1995. メダカのダイアローグ. ツバサ企画. [ref. 511]
- 森下郁子. 2000. 淡水生態系の保護と展望. 水環境学会誌, 23(3): 20-23. [ref. 516]
- 森下郁子. 2000. メダカを考える 淡水生態系の保護の展望. 水環境学会誌, 23(3): 148-151. [ref. 517]
- 森下郁子. 2001. 生態系の保全と回復に関する下水道の役割について(1). 月刊下水道, 24(7): 52-55. [ref. 518]
- 鍋島靖信・平松和也・上原一彦. 2003. 魚類の生息・繁殖に対する親水水路整備の効果. 大阪食とみどりの新技術, 2002: 21-22. [ref. 519]
- 中川雄三. 1998. 富士のすそ野のメダカの学校(科学で環境探検). 大日本図書, 55pp. [ref. 523]
- 中村滝男. 1999. 空とぶメダカ 絶滅危惧種メダカの不思議. ポプラ社, 79pp. [ref. 524]
- 中村滝男. 2000. メダカ保護活動の新たな展開.

- 水環境学会誌, 23(3): 12-15. [ref. 525]
- 中村滝男. 2000. メダカを考える メダカ保護活動の新たな展開. 水環境学会誌, 23(3): 140-143. [ref. 526]
- 西原重行. 1998. メダカと共に生きて. 電算印刷株式会社, 140pp. [ref. 529]
- 西岡健太郎・倉田健悟・上月康則・佐良家康・村上仁士・佐藤陽一・福田 守. 2000. 都市近郊用水路網内のメダカの生息環境に関する統計解析. 土木学会年次学術講演会公演概要集, 55: 18-19. [ref. 530]
- 沖津由季. 2000. 希少魚が暮らせる川づくり奮戦記6 小田原メダカがいる水田環境. 養殖, 37(8): 60-61. [ref. 533]
- 小澤祥司. 1999. 生きもの豊かな水田を再び. 水情報, 19(4): 8-11. [ref. 542]
- 小澤祥司. 2000. メダカが消える日－自然の再生をめざして-. 岩波書店, 224pp. [ref. 543]
- 酒泉 満. 2000. メダカを考える メダカとはどういう生きものか. 水環境学会誌, 23(3): 130-134. [ref. 549]
- 酒泉 満. 2002. メダカの遺伝的地域差と地域個体群の保全. 日本水環境学会セミナー, 43: 90-100. [ref. 550]
- 篠原亮太. 2000. メダカ保護の意味するところ. 水環境学会誌, 23(3): 1. [ref. 565]
- 白井康子. 2004. 香川県のため池の現状 - 希少動植物保護に向けた取り組み-. 国立環境研究所研究報告, (183): 16-25. [ref. 566]
- 水産庁. 1998. 日本の希少な野生水生生物に関するデータブック. 日本水産資源保護協会, pp.172-173. [ref. 569]
- 鈴木正貴・水谷正一・後藤 章. 2000. 環境に配慮した新技術4 水田生態系保全のための 小規模水田魚道の開発. 農業土木学会誌, 68(12): 1263-1266,1241(1). [ref. 570]
- 高木勇夫. 1992. 地図に刻まれた歴史と景観[藤沢市]. 新人物往来社, 158pp. [ref. 574]
- 竹田純一. 2002. 生物多様性の保全 新・生物多様性国家戦略の実践 里地里山と生物多様性. かんきょう (東京), 27(5): 15-16. [ref. 575]
- 竹内哲郎. 1988. めだかのお話. めだかの学校 株式会社, 18pp. [ref. 579]
- 多紀保彦. 2000. メダカの学校 (タイメダカと藤沢メダカ). 食器と容器, 6月.[ref. 580]
- 田中正彦. 2000. 都市の生物 生物からみた都市の姿 メダカはどこへいった. 生物の科学 遺伝, 54(8): 27-31. [ref. 583]
- 辻井要介・上田哲行. 2003. コンクリート化された水路におけるメダカの分布とそれに影響を及ぼす環境要因について. 環動昆, 14(3): 179-192. [ref. 593]
- 渡辺かほり. 2000. “藤沢メダカ”復活作戦. 養殖, 6月. [ref. 608]
- 渡辺かほり. 2000. メダカと環境教育「藤沢メダカの学校をつくる会」の活動から. 水環境学会誌, 23(3): 16-19. [ref. 609]
- 渡辺かほり. 2001. 藤沢の自然No.26, 鶴沼の自然. 藤沢市環境部みどり課. [ref. 610]
- 吉家世洋・森上義考. 1987. そだててみよう4, メダカ. あかね書房. [ref. 619]
- 吉崎正巳・太田一男. 1983. どきどき・自然, めだか. 福音館書店, 27pp. [ref. 622]

保 存 (Preservation)

- Aoki, K., M. Okamoto, K. Tatsumi and Y. Ishikawa. 1997. Cryopreservation of medaka spermatozoa. Zool. Sci., 14(4): 641-644. [ref. 7]
- Davis, C., M. Okihiro and D. Hinton. 2002. Effects of husbandry practices, gender, and normal physiological variation on growth and reproduction of Japanese medaka, *Oryzias latipes*. Aquat. Toxicol., 60(3-4): 185-201. [ref. 33]
- Egami, N. 1954. Effect of artificial photoperiodicity on time of oviposition in the fish *Oryzias latipes*. Annot. Zool. Japon., 27: 57-62. [ref. 38]
- Egami, N. 1956. How to rear the Japanese killifish, *Oryzias latipes*. In “Handbook of Genetics”, pp.851-852. [ref. 39]
- Egami, N. 1959. Record of the number of eggs obtained from a single pair of *Oryzias latipes*, kept in laboratory aquarium. J. Fac.

- Sci., Tokyo Univ. IV, 8: 521-538. [ref. 41]
- Shima, A., A. Shimada, J. Komura, K. Isa, K. Naruse, M. Sakaizumi and N. Egami. 1985. The preservation and utilization of wild populations of the medaka *Oryzias latipes*. Medaka, (3): 1-4. [ref. 320]
- Tiersch, T. R. 2001. Cryopreservation in aquarium fishes. Mar. Biotechnol., 3(Supplement1): S212-S223. [ref. 361]
- Tsukuda, H. and T. Katayama. 1957. The temperature adaptation in fishes I. The influence of rearing temperature on temperature tolerance, growth rate and body form. Physiol. and Ecol., 7: 113-122. [ref. 371]
- 相沢直樹・氏平政伸・島本敦介・谷下一夫. 1995. 生体組織凍結過程の熱的評価. 日本機械学会通常総会講演会講演論文集, 72(Pt1): 324-325. [ref. 419]
- 古川昭雄・椿 宜高・高村健二・永田尚志・可知直毅・高橋慎司・安岡善文・田村正行・渡辺信. 1995. 地球環境研究総合推進費による研究 野生生物種の減少に関する研究. 国立環境研究所地球環境研究年報, 1994: 79-88. [ref. 445]
- 太田克明・吉田元一・小野珠乙・山岸 哲. 1994. 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成5年度研究成果報告集2, pp.270-273.[ref. 537]
- 小澤祥司. 2000. メダカが消える日ー自然の再生をめざしてー. 岩波書店, 224pp. [ref. 543]
- 勝呂尚之. 2002. 日本産希少淡水魚の保護 水産試験場による希少淡水魚の生息地の復元. 生物の科学 遺伝, 56(6): 76-79, 12(2)-12(3). [ref. 568]
- 田口泰子. 1990. 海洋生物のジーンバンク 系統保存・凍結保存 メダカの系統保存について. 月刊海洋, 22(3): 142-148. [ref. 573]
- 椿 宜高・高村健二・永田尚志・富山清升・太田克明・吉田元一・山岸 哲・河田雅圭・土田浩治. 1996. 野生生物の種の減少 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成7年度終了研究成果報告集4, pp.271-287. [ref. 591]
- 椿 宜高・高村健二・富山清升・太田克明・吉田元一・山岸 哲・粕谷栄一・河田雅圭・伊沢雅子. 1995. 野生生物の種の減少 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成6年度研究成果報告集2, pp.280-284. [ref. 592]
- 氏平政伸・島本敦介・青木勝敏・相沢直樹・谷下一夫. 1995. メダカ受精卵の細胞外凍結における有効熱伝導率の測定. 日本機械学会熱工学部門講演会講演論文集, 1995: 117-119. [ref. 596]
- 氏平政伸・島本敦介・岡浩太郎・谷下一夫. 1996. 生体凍結過程における有効熱伝導率測定. バイオエンジニアリング学術講演会講演論文集, 7: 162-163. [ref. 597]
- 氏平政伸・山口 亮・相沢直樹・谷下一夫. 1995. mmオーダの生体組織の細胞外凍結による損傷. 日本機械学会論文集, 61(588): 3066-3074. [ref. 598]
- 氏平政伸・山口 亮・谷下一夫. 1995. mmオーダの生体組織の解凍過程に関する研究: 解凍後の顕微鏡観察によるメダカ受精卵の形態保存. 日本伝熱シンポジウム講演論文集, 32(1): 99-100. [ref. 599]
- 和田克彦・小林敬典・細谷和海・名古屋博之・前川光司・木曾克裕・小野里 坦・高田啓介・柴田直樹. 1995. 希少魚の個体群維持機構の解明と保存技術開発に関する予備的研究. 地球環境研究総合推進費 平成6年度研究成果報告集2, pp.344-348. [ref. 605]
- 若松佑子・尾里健二郎. 1998. メダカの地域集団 近交系 突然変異系. 蛋白質 核酸酵素, 43(11): 1487-1491. [ref. 607]
- 渡辺克己. 1994. 神奈川県産野生メダカの系統維持と教材化. 神奈川県立教育センター. [ref. 611]
- 山口 亮・氏平政伸・相沢直樹・谷下一夫.

1995. 生体組織の細胞外凍結による損傷の評価. 日本伝熱シンポジウム講演論文集, 32(1): 97-98. [ref. 614]

生態 (Ecology)

- Awaji, M. and I. Hanyu. 1989. Seasonal changes in ovarian response to photoperiods in orange-red type medaka. *Zool. Sci.* (Tokyo), 6(5): 943-950. [ref. 12]
- Awaji, M. and I. Hanyu. 1989. Temperature-photoperiod conditions necessary to begin the spawning season in wild type medaka. *Nippon Suisan Gakkaishi*, 55(4): 747. [ref. 13]
- Breder, C. M. Jr. 1954. Equations descriptive of fish schools and other animal aggregations. *Ecology*, 35: 361-370. [ref. 19]
- Bryant, M. J. and J. W. A. Grant. 1995. Resource defense, monopolization and variation of fitness in groups of female Japanese medaka depend on the synchrony of food arrival. *Anim. Behav.*, 49(6): 1469-1479. [ref. 21]
- Chan, K. K. 1976. A photosensitive daily rhythm in the female medaka, *Oryzias latipes*. *Can. J. Zool.*, 54(6): 852-856. [ref. 28]
- Dhanumkumari, C., R. K. Hanumantha and K. Shyamasundari. 1991. The life cycle of *Echinochasmus bagulai* (Trematoda: Echinostomatidae). *Int. J. Parasitol.*, 21: 259-263. [ref. 34]
- Dodd, J. M. 1960. Genetic and environmental aspects of sex determination in cold blooded vertebrates. *Mem. Soc. Endocrinol.*, 7: 17-44. [ref. 36]
- Egami, N. 1954. Effect of artificial photoperiodicity on time of oviposition in the fish *Oryzias latipes*. *Annot. Zool. Japon.*, 27: 57-62. [ref. 38]
- Egami, N. 1959. Effect of exposure to low temperature on the time of oviposition and the growth of the oocytes in the fish, *Oryzias latipes*. *J. Fac. Sci., Tokyo Univ.*

IV., 8: 539-548. [ref. 42]

Egami, N. 1969. Life span data for the small fish, *Oryzias latipes*. *Exp. Gerontol.*, 4: 127-129. [ref. 43]

Egami, N. and M. Nambu. 1961. Factors initiating mating behavior and oviposition in the fish, *Oryzias latipes*. *J. Fac. Sci., Tokyo Univ.*, IV, 9: 263-278. [ref. 44]

Grant, J. W. A. and L. D. Green. 1996. Mate copying versus preference for actively courting males by female Japanese medaka (*Oryzias latipes*). *Behav. Ecology*, 7(2): 165-167. [ref. 67]

Grant, J. W. A., M. J. Bryant and C. E. Soos. 1995. Operational sex ratio, mediated by synchrony of female arrival, alters the variance of male mating success in Japanese medaka. *Anim. Behav.*, 49(2): 367-375. [ref. 68]

Grant, J. W. A., P. C. Casey, M. J. Bryant and A. Shahsavaran. 1995. Mate choice by Japanese medaka (Pisces, Oryziidae). *Anim. Behav.*, 50(5): 1425-1428. [ref. 69]

Hamilton, J. B., R. O. Walter, R. M. Daniel and G. E. Mestler. 1969. Competition for mating between ordinary and supermale Japanese medaka fish. *Anim. Behav.*, 17(1): 168-176. [ref. 80]

Hata, K. 2002. Field experiment on the migration of fishes to paddy field with a small fishway. *JARQ.*, 36(4): 219-225. [ref. 85]

Hayashi, S. and T. Ueno. 2002. Quantification of male sexual behavior of the medaka fish (*Oryzias latipes*). *Neurosci. Res.*, (Supplement 26): S84. [ref. 90]

Hiramatsu, K. and S. Shikasho. 2002. Mathematical modeling of preference intensity of Japanese medaka for environmental factors in streamflow. *J. Fac. Agric. Kyusyu Univ.*, 47(1): 109-127. [ref. 102]

Hiramatsu, K., S. Shikasho and K. Mori. 2000. Mathematical modeling of fish schooling of Japanese Medaka using basic behavioral

- patterns. J. Fac. Agric. Kyusyu Univ., 45(1): 237-253. [ref. 103]
- Hosoda, K. 1958. The effect of water temperature upon the reward-type latent learning in *Oryzias latipes*. Ann. Animal Psychol., 8: 71-76. [ref. 117]
- Hosoda, K. 1958. The effect of water temperature on the latent learning in medakas and guppies. Japan. J. Psychol., 29: 48-50. [ref. 118]
- Howard, R. D., R. S. Martens, S. A. Innis, J. M. Drnevich and J. Hale. 1998. Mate choice and mate competition influence male body size in Japanese medaka. Anim. Behav., 55(5): 1151-1163. [ref. 120]
- Inoue, K. and Y. Takagi. 2001. Seawater adaptability of medaka (*Oryzias latipes*) and three related species. Zool. Sci., 18(Supplement): 115. [ref. 134]
- Inoue, K. and Y. Takei. 2002. Diverse Adaptability in *Oryzias* Species to High Environmental Salinity. Zool. Sci., 19(7): 727-734. [ref. 135]
- Ito, S. 1959. On the spawning time of the teleostean fishes. Japanese J. Ecol., 9: 3. [ref. 144]
- Kanda, T. and Y. Itazawa. 1986. When does the group effect on oxygen consumption appear in fish? J. Fac. Agric. Kyushu Univ., 30(4): 197-207. [ref. 169]
- Kobza, R. M., J. C. Trexler, W. F. Loftus and S. A. Perry. 2004. Community structure of fishes inhabiting aquatic refuges in a threatened Karst wetland and its implications for ecosystem management. Biol. Conserv., 116(2): 153-165. [ref. 194]
- Kuroiwa, T., A. Ohki, K. Naka and S. Maeda. 1994. Biomethylation and biotransformation of arsenic in a freshwater food chain: Green alga (*Chlorella vulgaris*) fwdarw shrimp (*Neocaridina denticulata*) fwdarw killifish (*Oryzias latipes*). Applied Organometallic Chemistry, 8(4): 325-333. [ref. 212]
- Magnuson, J. J. 1962. An analysis of the aggressive behavior, growth, and competition for food and space in medaka (*Oryzias latipes* (Pisces, Cyprinodontidae)). Can. J. Zool., 40: 313-363. [ref. 222]
- Mori, A. 1998. Prey-handling behavior of three species of homalopsine snakes: features associated with piscivory and Duvernoy's glands. Journal of Herpetology, 32(1): 40-50. [ref. 243]
- Mori, S. 1970. The influence of the water volume of containers upon the fighting behavior in *Oryzias latipes*. Ann. Animal Psychol., 20: 109-114. [ref. 244]
- Ogawa, K. 1957. Social behavior in the 3-individuals group of *Oryzias latipes*. Ann. Animal Psychol., 7: 72-78. [ref. 266]
- Onchi, R., Y. Koya and H. Munehara. 2002. Sperm-emission behavior by sneaker and its reproductive success in medaka fish. Zool. Sci., 19(12): 1501. [ref. 279]
- Ono, Y. and T. Uematsu. 1957. Mating ethogram in *Oryzias latipes*. J. Fac. Sci., Hokkaido Univ., VI, 13: 197-202. [ref. 282]
- Ono, Y. and T. Uematsu. 1958. Seasonal variation of social behavior in *Oryzias latipes*. Ann. Animal Psychol., 8: 63-69. [ref. 283]
- Ostrand, K. G. and G. R. Wilde. 2001. Temperature, dissolved oxygen, and salinity tolerances of five prairie stream fishes and their role in explaining fish assemblage patterns. Trans. Am. Fish Soc., 130(5): 742-749. [ref. 286]
- Ruzzante, D. E. and R. W. Doyle. 1991. Rapid behavioral changes in medaka (*Oryzias latipes*) caused by selection for competitive and non competitive growth. Evolution, 45: 1936-1946. [ref. 300]
- Ruzzante, D. E. and R. W. Doyle. 1993. Evolution of social behavior in a resource-rich, structured environment: selection experiments with medaka (*Oryzias latipes*). Evolution, 47: 456-470. [ref. 301]
- Sakaizumi, M. 1984. Rigid isolation between the northern population and the southern

- population of the Medaka, *Oryzias latipes*. Zool. Sci., 1: 795-800. [ref. 304]
- Sakaizumi, M., Y. Shimizu and S. Hamaguchi. 1992. Electrophoretic studies of meiotic segregation in inter- and intraspecific hybrids among east Asian species of the genus *Oryzias latipes* (Pisces: Oryziatidae). J. Exp. Zool., 246: 85-92. [ref. 307]
- Takeuchi, T. and E. Manabe. 1990. Statistical analysis on the sexual behavior of the medaka, *Oryzias latipes*. Research Bulletin of Shujitsu Women's College, 20: 1-32. [ref. 355]
- Taneda, K., T. Nishimoto and R. Abe. 1996. The sensory cue for making school in Medaka, *Oryzias latipes*. Mem. Fac. Sci. Kochi Univ. Series d Biology, 16-17(0): 69-75. [ref. 357]
- Taneda, K., T. Nishimoto and T. Matsuoka. 1991. The influence of neighbor fish on avoiding direction by sudden startling in medaka *Oryzias latipes*. Mem. Fac. Sci. Kochi Univ. Series d Biology, 12: 73-80. [ref. 358]
- Tatsuzawa, S., Y. Kano and M. Sakaizumi. 2001. The first record of two freshwater fishes, the Medaka *Oryzias latipes* and the loach *Misgurnus anguillicaudatus* from Mage-shima island: ecological, genetical, and historical views for conservation. Biogeography, 3: 89-100. [ref. 359]
- Trexler, J. C. and J. Travis. 1992. Effects of habitat and body size on mortality rates of *Poecilia latipinna*. Ecology, 73: 2224. [ref. 366]
- Tsukuda, H. and T. Katayama. 1957. The temperature adaptation in fishes I. The influence of rearing temperature on temperature tolerance, growth rate and body form. Physiol. and Ecol., 7: 113-122. [ref. 371]
- Tsuneki, K. and T. Sasaki. 1957. Fixation of poison by populations of *Oryzias latipes*. Japanese J. Ecol., 6: 153. [ref. 372]
- Walter, R. and J. Hamilton. 1970. Head-up movements as an indicator of sexual unreceptivity in female medaka, *Oryzias latipes*. Anim. Behav., 18: 125-127. [ref. 384]
- Weber, D. N. and R. E. Spieler. 1987. Effects of the light-dark cycle and scheduled feeding on behavioral and reproductive rhythms of the cyprinodont fish, Medaka, *Oryzias latipes*. Experientia, 43(6): 621-624. [ref. 387]
- Weis, J. S. and P. Weis. 2003. Is the invasion of the common reed, *Phragmites australis*, into tidal marshes of the eastern US an ecological disaster? Mar. Pollut. Bull., 46(7): 816-820. [ref. 388]
- Yamamoto, T. 1961. Progenies of sex-reversal females mated with sex-reversal males in the medaka, *Oryzias latipes*. J. Exp. Zool., 146: 163-179. [ref. 403]
- Yokota, T. and T. Oishi. 1992. Seasonal change in the locomotor activity rhythm of the medaka, *Oryzias latipes*. Int. J. Biometeorol., 36: 39-44. [ref. 414]
- Yoshioka, H. 1962. On the effects of environmental factors upon the reproduction of fishes. I. The effects of day-length on the reproduction of the Japanese killifish, *Oryzias latipes*. Bull. Fac. Fish., Hokkaido Univ., 13: 123-136. [ref. 415]
- Yoshioka, H. 1966. On the effects of environmental factors upon the reproduction of fishes. III. The occurrence and regulation of refractory period in the photoperiodic response of medaka, *Oryzias latipes*. J. Hokkaido Univ. Educ., 17: 23-33. [ref. 416]
- 安芸昌彦. 2004. 栗林公園の淡水魚類. 香川生物, 31: 13-22. [ref. 420]
- 秋月岩魚. 1999. ブラックバスがメダカを食う -日本の生態系が危ない-. 宝島社新書, 222pp. [ref. 421]
- 青井 透. 2001. 研究現場からの最新報告(6)水の循環利用と豊かな生態系復元を目的とす

- るビオトープ創り. 環境施設, 83: 81-87. [ref. 423]
- 有薗幸司. 2003. 環境ホルモンの生態系への影響. NIRS-M, 169: 133-140. [ref. 427]
- 土井麻記子・高村嘉幸・深津和彦・鴻野雅一. 2004. アクアリウムを用いた生態系に対する排水の安全性評価に関する研究. 水環境学会誌, 27(5): 343-348. [ref. 432]
- 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp. [ref. 436]
- 江上信雄・山上健次郎・嶋 昭紘. 1990. メダカの生物学. 東京大学出版会, 315pp. [ref. 437]
- 遠藤 徹. 2004. メダカの里「家根合地区」水田生態系保全の取り組みと地域づくり, ほ場と土壤, 36(2): 29-34. [ref. 440]
- 福本紘一・宮下敏夫・南部和人. 2003. 水生植物の移植とビオトープ. 水草研究会誌, 79: 29-32. [ref. 444]
- 後出明子・坂 正臣・大西正健. 2003. 京都府下におけるメダカ生息調査と評価分析. 生物高分子, 3(1): 31-34. [ref. 450]
- 端 憲二・竹村武士・本間新哉・佐藤政良. 2001. 流れにおけるメダカの遊泳行動に関する実験的考察. 農業土木学会誌, 69(9): 987-992. [ref. 454]
- 平松和昭・四ヶ所四男美・福田信二. 2003. ファジイ推論によるメダカの環境応答モデルの開発. 農業土木学会論文集, 228: 789-796. [ref. 458]
- 平松和也・山本雅照. 1998. 生態系維持回復のための水際帯調査. 大阪府立淡水魚試験場業務報告, 30: 17-19. [ref. 459]
- 市川憲平. 1993. 雄が子守をする虫たち 5, メダカの生態(2). 海洋と生物, 15(6): 421-426. [ref. 467]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]
- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp. [ref. 479]
- 環境省 自然環境局. 2004. 農業水産省と環境省の連携による「田んぼの生きもの調査2003」の結果について. かんきょう(東京), 29(6): 38-39. [ref. 491]
- 北野 聰・山形哲也・市川 寛・小林 尚. 2002. 長野県北部の水田用水路における魚類群集及びメダカの推定個体数. 滋賀自然教育研究施設研究業績, 39: 17-20. [ref. 497]
- 北野 聰・山形哲也・市川 寛・小林 尚. 2003. 長野県北部の水田用水路に生息する淡水魚類: 耕作放棄 1年後の状況. 滋賀自然教育研究施設研究業績, 40: 29-32. [ref. 498]
- 小島 昭. 2001. 炭素繊維による藻場形成 棚名湖での挑戦. 水, 43(12): 27-31. [ref. 502]
- 小島 覚. 1999. 環境を考える・シリーズ10森林の減少と生物多様性(その3). 菌じん, 45(12): 32-38. [ref. 503]
- 草野慎司. 1996. メダカのくらし. あかね書房, 54pp. [ref. 506]
- 宮本敦史. 2001. 水生生物の分布, 生態調査 淡水魚類の分布と希少魚類の繁殖基礎試験. 三重県科学技術振興センター水産技術センター事業報告, 2000: 93-94. [ref. 513]
- 宮本敦史・藤吉利彦. 2003. 水生生物の分布, 生体調査-I 淡水魚類の分布と希少魚類の繁殖基礎試験. 三重県科学技術振興センター水産研究部事業報告, 2002: 98-99. [ref. 514]
- 森下郁子. 2000. メダカを考える 淡水生態系の保護の展望. 水環境学会誌, 23(3): 148-151. [ref. 517]
- 森下郁子. 2001. 生態系の保全と回復に関する下水道の役割について(1). 月刊下水道, 24(7): 52-55. [ref. 518]
- 西原重行. 1998. メダカと共に生きて. 電算印刷株式会社, 140pp. [ref. 529]
- 西 隆行・神崎たかし・谷口三紀生・立石 康彦・岩崎英二. 2001. ビオトープの構築に関する基礎研究. 日本環境衛生センター所報, 27: 43-50. [ref. 531]
- 西 隆行・谷口三紀生・立石康彦・宮原哲也. 2001. ビオトープ構築の試み. 日本環境衛生センター所報, 28: 98-103. [ref. 532]
- 沖津由季・勝呂尚之. 2001. メダカを中心とした小田原市桑原・鬼柳農業用水路の魚類. 神奈川自然誌資料, 22: 51-59. [ref. 534]
- 太田滋規・西森克浩・山中 治. 2003. 8.琵琶湖沿岸帶魚類等調査費1)琵琶湖および河川

- 下流の魚類等生息状況調査. 滋賀県水産試験場事業報告, 2002: 112-113. [ref. 538]
- 小澤祥司. 1999. 生きもの豊かな水田を再び. 水情報, 19(4): 8-11. [ref. 542]
- 小澤祥司. 2000. メダカが消える日—自然の再生をめざして-. 岩波書店, 224pp. [ref. 543]
- 酒泉 満・久保孝一. 1981. カラー自然シリーズ35, メダカ. 偕成社, 38pp. [ref. 551]
- 笹井隆邦. 2001. 藍那のため池の生物. 神戸常盤短期大学紀要, 22: 23-30. [ref. 552]
- 白井康子・木村正英. 2004. 香川県のため池の現状—希少動植物保護に向けた取り組み-. 国立環境研究所研究報告, 183: 16-25. [ref. 566]
- 勝呂尚之. 2002. 日本産希少淡水魚の保護 水産試験場による希少淡水魚の生息地の復元. 生物の科学 遺伝, 56(6): 76-79, 12(2)-12(3). [ref. 568]
- 鈴木正貴・水谷正一・後藤 章. 2000. 環境に配慮した新技術4 水田生態系保全のための小規模水田魚道の開発. 農業土木学会誌, 68(12): 1263-1266, 1241(1). [ref. 570]
- 竹田純一. 2002. 生物多様性の保全 新・生物多様性国家戦略の実践 里地里山と生物多様性. かんきょう(東京), 27(5): 15-16. [ref. 575]
- 竹村武士・奥島修二・山本勝利・小出水規行・端憲二. 2003. 小水路においてメダカが高い選好性を示す底質条件. 農業工学研究所研究成果情報, 2002: 17-18. [ref. 577]
- 竹内邦輔. 1981. 誰にでもできるメダカの実験. 新光印刷株式会社, 110pp. [ref. 578]
- 竹内哲郎. 1988. めだかのお話. めだかの学校 株式会社, 18pp. [ref. 579]
- 田中宏明・高橋明宏・東谷 忠・矢古宇靖子・斎藤正義. 2001. 下水道による水生生態系への影響に関する調査. 国土技術政策総合研究所資料, (10): 225-230. [ref. 581]
- 田中正治・平松和也. 2004. 淀川の生息魚類の分布およびその生態. 大阪府立食とみどりの総合技術センター水生生物センター研究報告, 12: 69. [ref. 582]
- 田中正彦. 2000. 都市の生物 生物からみた都市の姿 メダカはどこへいった. 生物の科学 遺伝, 54(8): 27-31. [ref. 583]
- 寺尾 修. 1985. 野生メダカの生態. 生物の科学 遺伝, 39(8): 47-50. [ref. 587]
- 辻井要介・上田哲行. 2003. コンクリート化された水路におけるメダカの分布とそれに影響を及ぼす環境要因について. 環動昆, 14(3): 179-192. [ref. 593]
- 上山尊士. 1998. 生態影響試験業務. TRC News, 62: 13-16. [ref. 595]
- 梅津 剛. 2002. 微生物からの生態系を考えた海水魚養殖装置の開発. 養殖, 39(7): 84-87. [ref. 600]
- 宇根 豊. 1999. 田んぼのエコロジー 田んぼの虫たちは, 自然の生きもの. 生物の科学 遺伝, 53(4): 26-30. [ref. 601]
- 宇根 豊. 2000. いのちが集まる・命が育む「田んぼの学校」入学編. 農村環境整備センター, pp.38-41. [ref. 602]
- 横山 正. 1987. メダカのそだちかた ポプラ社の実験観察シリーズ3. ポプラ社, 39pp. [ref. 618]
- 吉岡義正・小瀬洋喜・後藤幹保・ヒキノ ヒロシ. 1987. 主として水棲の動物及び植物を用いた生態影響試験法間の関係. 衛生化学, 33(1): 11-19. [ref. 621]
- 吉崎正巳・太田一男. 1983. めだか(どきどき・自然). 福音館書店, 27pp. [ref. 622]

行 動 (Behavior)

- Egami, N. and M. Nambu. 1961. Factors initiating mating behavior and oviposition in the fish, *Oryzias latipes*. J. Fac. Sci., Tokyo Univ., IV, 9: 263-278. [ref. 44]
- Grant, J. W. A. and L. D. Green. 1996. Mate copying versus preference for actively courting males by female Japanese medaka (*Oryzias latipes*). Behav. Ecology, 7(2): 165-167. [ref. 67]
- Grant, J. W. A., M. J. Bryant and C. E. Soos. 1995. Operational sex ratio, mediated by synchrony of female arrival, alters the variance of male mating success in

- Japanese medaka. Anim. Behav., 49(2): 367-375. [ref. 68]
- Grant, J. W. A., P. C. Casey, M. J. Bryant and A. Shahsavari. 1995. Mate choice by Japanese medaka (Pisces, Oryziidae). Anim. Behav., 50(5): 1425-1428. [ref. 69]
- Hamilton, J. B., R. O. Walter, R. M. Daniel and G. E. Mestler. 1969. Competition for mating between ordinary and supermale Japanese medaka fish. Anim. Behav., 17(1): 168-176. [ref. 80]
- Hata, K. 2002. Field experiment on the migration of fishes to paddy field with a small fishway. JARQ., 36(4): 219-225. [ref. 85]
- Hayashi, S. and T. Ueno. 2002. Quantification of male sexual behavior of the medaka fish (*Oryzias latipes*). Neurosci. Res., (Supplement 26): S84. [ref. 90]
- Hidaka, H. and R. Tatsukawa. 1985. Avoidance test of a fish, medaka (*Oryzias latipes*), to aquatic contaminants, with special reference to monochloramine. Arch. Environ. Contam. Toxicol., 14(5): 565-571. [ref. 97]
- Hidaka, H. and R. Tatsukawa. 1989. Avoidance by olfaction in a fish, medaka (*Oryzias latipes*), to aquatic contaminants. Environ. Pollut., 56(4): 299-309. [ref. 98]
- Hiramatsu, K., S. Shikasho and K. Mori. 2000. Mathematical modeling of fish schooling of Japanese Medaka using basic behavioral patterns. J. Fac. Agric. Kyusyu Univ., 45(1): 237-253. [ref. 102]
- Howard, R. D., R. S. Martens, S. A. Innis, J. M. Drnevich and J. Hale. 1998. Mate choice and mate competition influence male body size in Japanese medaka. Anim. Behav., 55(5): 1151-1163. [ref. 120]
- Magnuson, J. J. 1962. An analysis of the aggressive behavior, growth, and competition for food and space in medaka (*Oryzias latipes* (Pisces, Cyprinodontidae)). Can. J. Zool., 40: 313-363. [ref. 222]
- Mori, A. 1998. Prey-handling behavior of three species of homalopsine snakes: features associated with piscivory and Duvernoy's glands. Journal of Herpetology, 32(1): 40-50. [ref. 243]
- Mori, S. 1970. The influence of the water volume of containers upon the fighting behavior in *Oryzias latipes*. Ann. Animal Psychol., 20: 109-114. [ref. 244]
- Ogawa, K. 1957. Social behavior in the 3-individuals group of *Oryzias latipes*. Ann. Animal Psychol., 7: 72-78. [ref. 266]
- Onchi, R., Y. Koya and H. Munehara. 2002. Sperm-emission behavior by sneaker and its reproductive success in medaka fish. Zool. Sci., 19(12): 1501. [ref. 279]
- Ono, Y. and T. Uematsu. 1957. Mating ethogram in *Oryzias latipes*. J. Fac. Sci., Hokkaido Univ., VI, 13: 197-202. [ref. 282]
- Ono, Y. and T. Uematsu. 1958. Seasonal variation of social behavior in *Oryzias latipes*. Ann. Animal Psychol., 8: 63-69. [ref. 283]
- Packer, A. 2001. Medaka on the move. Nat. Genet., 28(4): 302. [ref. 288]
- Ruzzante, D. E. and R. W. Doyle. 1993. Evolution of social behavior in a resource-rich, structured environment: selection experiments with medaka (*Oryzias latipes*). Evolution, 47: 456-470. [ref. 301]
- Takeuchi, T. and E. Manabe. 1990. Statistical analysis on the sexual behavior of the medaka, *Oryzias latipes*. Research Bulletin of Shujitsu Women's College, 20: 1-32. [ref. 355]
- Taneda, K., T. Nishimoto and R. Abe. 1996. The sensory cue for making school in Medaka, *Oryzias latipes*. Mem. Fac. Sci. Kouchi Univ. Series d Biology, 16-17(0): 69-75. [ref. 357]
- Taneda, K., T. Nishimoto and T. Matsuoka. 1991. The influence of neighbor fish on avoiding direction by sudden startling in medaka *Oryzias latipes*. Mem. Fac. Sci. Kouchi Univ. Series d Biology, 12: 73-80.

- [ref. 358]
- Walter, R. and J. Hamilton. 1970. Head-up movements as an indicator of sexual unreceptivity in female medaka, *Oryzias latipes*. Anim. Behav., 18: 125-127. [ref. 384]
- Walter, R. and J. Hamilton. 1970. Supermales (YY sex chromosomes) and androgen-treated XY males: competition for mating with female killifish *Oryzias latipes*. Anim. Behav., 18: 128-131. [ref. 385]
- Weber, D. N. and R. E. Spieler. 1987. Effects of the light-dark cycle and scheduled feeding on behavioral and reproductive rhythms of the cyprinodont fish, Medaka, *Oryzias latipes*. Experientia, 43(6): 621-624. [ref. 387]
- 江口星雄. 1995. IML-2 “メダカの宇宙における交尾・産卵行動” 実験の概要. 東京大学原子力研究総合センター技術発表会技術報告, 4: 26-33. [ref. 439]
- 古川瑠美・野呂晋一・長尾和右・井尻憲一・江口星雄. 1999. メダカ稚魚の成長に伴う視覚の発達および微小重力下での挙動. 宇宙生物科学, 13(3): 164-165. [ref. 446]
- 古屋康則・恩地理恵. 2002. メダカの交接・産卵行動の観察法. 岐阜大学教育学部研究報告自然科学, 26(2): 19-22. [ref. 449]
- 端 憲二・本間新哉・中 達男. 2001. 流速増大時のメダカの避難行動. 農業工学関係研究成果情報, 2000: 13-14. [ref. 453]
- 端 憲二・竹村武士・本間新哉・佐藤政良. 2001. 流れにおけるメダカの遊泳行動に関する実験的考察. 農業土木学会誌, 69(9): 987-992. [ref. 454]
- 畠山 仁・狩野賢司. 2001. メダカ *Oryzias latipes* の雌の配偶者選択 雌の体サイズ・体重に対する好み. 日本獣医畜産大学研究報告, 50: 25-33. [ref. 455]
- 平松和昭・四ヶ所四男美・福田信二. 2003. フィジイ推論によるメダカの環境応答モデルの開発. 農業土木学会論文集, 228: 789-796. [ref. 458]
- 鈴木正貴・水谷正一・後藤 章. 2001. 水田水域における淡水魚の双方向移動を保証する小規模魚道の試作と実験. 応用生態工学, 4(2): 163-177. [ref. 570]
- 竹村武士・奥島修二・山本勝利・小出水規行・端憲二. 2003. 小水路においてメダカが高い選好性を示す底質条件. 農業工学研究所研究成果情報, 2002: 17-18. [ref. 577]
- 寺尾 修. 1985. 野生メダカの生態. 生物の科学 遺伝, 39(8): 47-50. [ref. 587]
- ビオトープ・水田 (Biotope and Paddy field)**
- Hata, K. 2002. Field experiment on the migration of fishes to a paddy field with a small fishway. JARQ, 36(4): 219-225. [ref. 85]
- Iguchi, K., M. Yoshiya, W. Wu and N. Shimizu. 1999. Biodiversity of freshwater fishes in the paddy water systems of Anji, China. National Res, Inst Fisheries Sci. 13: 27-36. [ref. 124]
- 青井 透. 2001. 研究現場からの最新報告 (6) 水の循環利用と豊かな生態系復元を目的とするビオトープ創り. 環境施設, 83: 81-87. [ref. 423]
- 伴 幸成. 1993. 田んぼの生き物たち. 大日本図書, 35pp. [ref. 428]
- 土井敏男. 閉鎖的ビオトープで見られたメダカの体型変異. 兵庫陸水生物, (53): 73-79. [ref. 433]
- 金子美智雄. 2000. 田んぼに行こう 里山探検をしよう, テーマ発見! 総合学習体験ブック. ほるぶ出版, 40pp. [ref. 486]
- 環境省 自然環境局. 2004. 農林水産省と環境省の連携による「田んぼの生きもの調査2003」の結果について. かんきょう(東京), 29(6): 38-39. [ref. 491]
- 小澤祥司. 1999. 生きもの豊かな水田を再び. 水情報, 19(4): 8-11. [ref. 542]
- 小澤祥司. 2000. メダカが消える日-自然の再生をめざして-. 岩波書店, 224pp. [ref. 543]
- 田部和博・安部伸治. 2003. 「自然再生推進法」の制定と「バイオマス・ニッポン総合戦略」の策定-2 生物多様性国家戦略と環境省の

- 自然再生事業について. 農業土木学会誌, 71(8): 705-708. [ref. 571]
- 竹田純一. 2002. 生物多様性の保全 新・生物多様性国家戦略の実践 里地里山と生物多様性. かんきょう, 27(5): 15-16. [ref. 575]
- 宇根 豊. 1999. 田んぼのエコロジー 田んぼの虫たちは、自然の生きもの. 生物の科学 遺伝, 53(4): 26-30. [ref. 601]
- 宇根 豊. 2000. いのちが集まる・命が育む「田んぼの学校」入学編. 農村環境整備センター, pp.38-41. [ref. 602]
- 牛野 正. 2004. 農業土木とランドスケープエコロジー-3 スイスの景観コンセプト. 農業土木学会誌, 72(1): 27-31, 1(1). [ref. 603]

毒 性 (Toxicology)

- Allinson, G. and M. Morita. 1995. Bioaccumulation and toxic effects of elevated of 3,3',4,4'-tetrachloroazobenzene (33'44'-TCAB) towards aquatic organisms. 2: Bioaccumulation and toxic effects of dietary 33'44'-TCAB on the Japanese medaka (*Oryzias latipes*). Chemosphere, 30(2): 223-232. [ref. 3]
- Allinson, G., M. Ueoka and M. Morita. 1994. Effect of dietary 1,3,6,8-tetrachlorodibenzo-p-dioxin on the Japanese freshwater fish *Oryzias latipes* (Medaka) and aquatic snail *Indoplanorbis exustus* (Indohiramakigai). Chemosphere, 28(7): 1369-1383. [ref. 4]
- Aoki, K. and H. Matsudaira. 1984. Factors influencing methylazoxymethanol acetate initiation of liver tumors in *Oryzias latipes*: carcinogen dosage and time of exposure. Natl. Cancer Inst. Monogr., 65: 345-351. [ref. 5]
- Belanger, S. E., D. S. Cherry and J. JR. Cairns. 1990. Functional and pathological impairment of Japanese Medaka (*Oryzias latipes*) by long-term asbestos exposure. Aquatic Toxicol., 17(2): 133-154. [ref. 14]
- Bentivegna, C. S. and T. Piatkowski. 1998.

- Effects of tributyltin on medaka (*Oryzias latipes*) embryos at different stages of development. Aquatic Toxicol., 44(1-2): 117-128. [ref. 15]
- Braunbeck, T. A., S. J. Ted, S. M. Lester and E. D. Hinton. 1992. Ultrastructural alterations in liver of medaka (*Oryzias latipes*) exposed to diethylnitrosamine. Toxicologic Pathology, 20(2): 179-196. [ref. 18]
- Brittelli, M. R., H. H. Chen and C. F. Muska. 1985. Induction of branchial (gill) neoplasms in the medaka fish (*Oryzias latipes*) by N-methyl-N'-nitrosoguanidine. Cancer Res., 45(7): 3209-3214. [ref. 20]
- Bunton, T. E. 1989. Diethylnitrosamine (DEN)-induced degenerative, proliferative and neoplastic lesions in the liver of the medaka (*Oryzias latipes*) following short-term exposure. Mar. Environ. Res., 28(1-4): 369-374. [ref. 22]
- Cantrell, S. M., L. H. Lutz, M. Hannink and D. E. Tillitt. 1996. Embryotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD): the embryonic vasculature is a physiological target for TCDD-induced DNA damage and apoptotic cell death in medaka (*Oryzias latipes*). Toxicol. Appl. Pharmacol., 141(1): 23-34. [ref. 27]
- Chulakasem, W., J. A. Nelson and J. J. Magnuson. 1989. Interaction between effects of low pH and low ion concentration on mortality during early development of medaka, *Oryzias latipes*. Can. J. Zool., 67(9): 2158-2168. [ref. 31]
- Cooper, K. R., H. Liu, P-A. Bergqvist and C. Rappe. 1991. Evaluation of baltic herring and Icelandic cod liver oil for embryo toxicity, using the Japanese medaka (*Oryzias latipes*) embryo larval assay. Environ. Toxicol. Chem., 10(6): 707-714. [ref. 32]
- Dial, N. A. 1978. Methylmercury: some effects on embryogenesis in the Japanese medaka,

- Oryzias latipes*. Teratology, 17(1): 83-91. [ref. 35]
- Etoh, H. and Y. Hyodo-Taguchi. 1983. Effects of tritiated water on germ cells in medaka embryos. Radiat. Res., 93(2): 332-339. [ref. 45]
- Fabacher, D. L., J. M. Besser, C. J. Schmitt, J. C. Harshbarger, P. H. Peterman and J. A. Lebo. 1991. Contaminated sediments from tributaries of the Great Lakes: chemical characterization and carcinogenic effects in medaka (*Oryzias latipes*). Arch. Environ. Contam. Toxicol., 21(1): 17-34. [ref. 47]
- Featherstone, D., C. D. Drewes, J. R. Coats and S. P. Bradbury. 1993. A non-invasive neurotoxicity assay using larval medaka. ASTM. Spec. Tech. Publ., 1216: 275-288. [ref. 49]
- Gonzalez-Doncel, M., C. Barrueco, E. De, La, Pena and D. E. Hinton. 2003. Stage sensitivity of medaka (*Oryzias latipes*) eggs and embryos to permethrin. Aquatic Toxicol., 62(3): 255-268. [ref. 65]
- Gray, M. A. and C. D. Metcalfe. 1999. Toxicity of 4-tert-octylphenol to early life stages of Japanese medaka (*Oryzias latipes*). Aquatic Toxicol., 46(2): 149-154. [ref. 70]
- Gronen, S., S. Manning, S. Barnes, D. Barnes, M. Brouwer and N. Denslow. 1999. Serum vitellogenin levels and reproductive impairment of male Japanese medaka (*Oryzias latipes*) exposed to 4-tert-octylphenol. Environ. Health Perspect., 107(5): 385-390. [ref. 71]
- Haasch, M. L. and A. W. Ford. 2004. Combined effects of ethanol and cinnamaldehyde in the Japanese medaka embryo-larval assay (MELA). Mar. Environ. Res., 58(2-5): 175-179. [ref. 74]
- Hatanaka, J., N. Doke, T. Harada, T. Aikawa and M. Enomoto. 1982. Usefulness and rapidity of screening for the toxicity and carcinogenicity of chemicals in medaka, *Oryzias latipes*. Jpn. J. Exp. Med., 52(5): 243-253. [ref. 86]
- Hawkins, W. E., J. W. Fournie, R. M. Overstreet and W. W. Walker. 1986. Intraocular neoplasms induced by methylazoxymethanol acetate in Japanese medaka (*Oryzias latipes*). J. Natl. Cancer Inst., 76(3): 453-465. [ref. 87]
- Hawkins, W. E., R. M. Overstreet, J. W. Fournie and W. W. Walker. 1985. Development of aquarium fish models for environmental carcinogenesis: tumor induction in seven species. J. Appl. Toxicol., 5(4): 261-264. [ref. 88]
- Hawkins, W. E., W. W. Walker, R. M. Overstreet, J. S. Lytle and T. F. Lytle. 1990. Carcinogenic effects of some polycyclic aromatic hydrocarbons on the Japanese medaka and guppy in waterborne exposures. Sci. Total Environ., 94(1-2): 155-167. [ref. 89]
- Heath, A. G., J. J. Cech, J. G. Zinkl and M. D. Steele. 1993. Sublethal effects of three pesticides on Japanese medaka. Arch. Environ. Contam. Toxicol., 25(4): 485-491. [ref. 92]
- Heisinger, J. F. and W. Green. 1975. Mercuric chloride uptake by eggs of the ricefish and resulting teratogenic effects. Bull. Environ. Contam. Toxicol., 14(6): 665-673. [ref. 93]
- Helmstetter, M. F. and R. W. III. Alden. 1995. Passive trans-chorionic transport of toxicants in topically treated Japanese medaka (*Oryzias latipes*) eggs. Aquatic Toxicol., 32(1): 1-13. [ref. 94]
- Helmstetter, M. F. and R. W. III. Alden. 1995. Toxic responses of Japanese medaka (*Oryzias latipes*) eggs following topical and immersion exposures to pentachlorophenol. Aquatic Toxicol., 32(1): 15-29. [ref. 95]
- Hidaka, H. and R. Tatsukawa. 1985. Avoidance test of a fish, medaka (*Oryzias latipes*), to aquatic contaminants, with special reference to monochloramine. Arch. Environ. Contam. Toxicol., 14(5): 565-571.

- [ref. 97]
- Hinton, D. E., R. C. Lantz and J. A. Hampton. 1984. Effect of age and exposure to a carcinogen on the structure of the medaka liver: a morphometric study. *Natl. Cancer Inst. Monogr.*, 65: 239-249. [ref. 101]
- Hiraoka, Y. and H. Okuda. 1983. Characteristics of vertebral abnormalities of medaka as a water pollution indicator. *Hiroshima J. Med. Sci.*, 32(3): 261-266. [ref. 104]
- Hiraoka, Y., S. Ishizawa, T. Kamada and H. Okuda. 1985. Acute toxicity of 14 different kinds of metals affecting medaka fry. *Hiroshima J. Med. Sci.*, 34(3): 327-330. [ref. 105]
- Hiraoka, Y., J. Tanaka and H. Okuda. 1989. Toxicity to medaka of solutions of fenitrothion degraded by strong alkali. *Environ. Pollut.*, 58(1): 35-42. [ref. 106]
- Hiraoka, Y., J. Tanaka and H. Okuda. 1990. Toxicity of fenitrothion degradation products to medaka (*Oryzias latipes*). *Bull. Environ. Contam. Toxicol.*, 44(2): 210-215. [ref. 107]
- Holcombe, G. W., D. A. Benoit, D. E. Hammermeister, E. N. Leonard and D. R. Johnson. 1995. Acute and long term effects of nine chemicals on Japanese medaka (*Oryzias latipes*). *Arch. Environ. Contam. Toxicol.*, 28(3): 287-297. [ref. 112]
- Hornung, M. W., P. M. Cook, K. M. Flynn, D. B. Lothenbach, R. D. Johnson and J. W. Nichols. 2004. Use of multi-photon laser-scanning microscopy to describe the distribution of xenobiotic chemicals in fish early life stages. *Aquat. Toxicol.*, 67(1): 1-11. [ref. 116]
- Hyodo-Taguchi, Y. and H. Matsudaira. 1987. Higher susceptibility to N-methyl-N'-nitro-N-nitrosoguanidine-induced tumorigenesis in an interstrain hybrid of the fish, *Oryzias latipes* (medaka). *Jpn. J. Cancer Res.*, 78(5): 487-493. [ref. 123]
- Inamori, Y., M. Kubo, H. Tsujibo, M. Ogawa, Y. Saito, Y. Miki and S. Takemura. 1987. The ichthyotoxicity and coronary vasodilator action of 3,3'-dihydroxy-alpha,beta-diethylstilbene. *Chem. Pharm. Bull (Tokyo)*, 35(2): 887-890. [ref. 130]
- Ishikawa, T., P. Masahito and S. Takayama. 1984. Usefulness of the medaka, *Oryzias latipes*, as a test animal: DNA repair processes in medaka exposed to carcinogens. *Natl. Cancer Inst. Monogr.*, 65: 35-43. [ref. 136]
- Ishikawa, T. and S. Takayama. 1979. Importance of hepatic neoplasms in lower vertebrate animals as a tool in cancer research. *J. Toxicol. Environ. Health*, 5 (2-3): 537-550. [ref. 137]
- Jacquet, C., V. Thermes, A. de, Luze, S. Puiseux-Dao, C. Bernard, J. S. Joly, F. Bourrat and M. Edery. 2004. Effects of microcystin-LR on development of medaka fish embryos (*Oryzias latipes*). *Toxicon*, 43(2): 141-147. [ref. 162]
- James, M., W. Hawkins and W. Walker. 1994. Phase 1 and phase 2 biotransformation and carcinogenicity of 2-acetylaminofluorene in medaka and guppy. *Aquatic Toxicol.*, 28 (1-2): 79-95. [ref. 166]
- Kashiwaba, S., H. Ishikawa, N. Miyamoto, Y. Ohnishi and Y. Magara. 2002. Fish test for endocrine-disruption and estimation of water quality of Japanese rivers. *Water Res.*, 36(8): 2161-2166. [ref. 171]
- Kaur, R., B. Buckley, K. R. Cooper, S. S. Park and Y. K. Kim. 1996. Toxicity test of nanji island landfill (Seoul, Korea) leachate using Japanese medaka (*Oryzias latipes*) embryo larval assay. *Bull. Environ. Contam. Toxicol.*, 57(1): 84-90. [ref. 175]
- Kiparissis, Y., T. L. Metcalfe, G. C. Balch and C. D. Metcalfe. 2003. Effects of antiandrogens, vi clozolin and cyproterone acetate on gonadal development in Japanese medaka (*Oryzias latipes*). *Aquatic Toxicol.*, 63(4): 391-403. [ref. 189]
- Klaunig, J. E., B. A. Barut and P. J. Goldblatt. 1984. Preliminary studies on the usefulness

- of medaka, *Oryzias latipes*, embryos in carcinogenicity testing. Natl. Cancer Inst. Monogr., 65: 155-161. [ref. 191]
- Kleaks, P. L., J. A. Nyman and S. Bhattacharyya. 2004. Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms. Environ. Pollut., 129(3): 345-353. [ref. 192]
- Koizumi, N., M. Miyajima and M. Susukida. 1992. Variation of zinc and copper in metallothionein-like protein in killifish (*Oryzias latipes*) exposed to cadmium. Chemosphere, 24(12): 1799-1803. [ref. 205]
- Kubota, Y., A. Shimada and A. Shima. 1992. Detection of γ -ray-induced DNA damages in malformed dominant lethal embryos of the Japanese medaka (*Oryzias latipes*) using AP-PCR fingerprinting. Mutat. Res., 283(4): 263-270. [ref. 210]
- Lauren, D. J., S. J. Teh and D. E. Hinton. 1990. Cytotoxicity phase of diethylnitrosamine-induced hepatic neoplasia in medaka. Cancer Res., 50(17): 5504-5514. [ref. 216]
- Lelewellyn, G. C., G. A. Stephenson and J. W. Hofman. 1977. Aflatoxin B1 induced toxicity and teratogenicity in Japanese Medaka eggs (*Oryzias latipes*). Toxicon, 15(6): 582-587. [ref. 219]
- Leung, T. S. and R. V. Bulkley. 1979. Effects of petroleum hydrocarbons on length of incubation and hatching success in the Japanese medaka. Bull. Environ. Toxicol., 23(1-2): 236-243. [ref. 220]
- Marty, G. D., J. M. Munoz, D. J. Lauren and D. E. Hinton. 1990. Age-dependent changes in toxicity of N-nitroso compounds to Japanese medaka (*Oryzias latipes*) embryos. Aquatic Toxicol., 17(1): 45-62. [ref. 225]
- Marty, G. D., S. Wetzlich, J. M. Nunez, A. Craigmill and D. E. Hinton. 1991. Fish-based biomonitoring to determine toxic characteristics of complex chemical mixtures: documentation of bioremediation at a pesticide disposal site. Aquat. toxicol. (Amst.), 19(4): 329-340. [ref. 226]
- Michibata, H. 1981. Effect of water hardness on the toxicity of cadmium to the egg of the teleost *Oryzias latipes*. Bull. Environ. Contam. Toxicol., 27(2): 187-192. [ref. 239]
- Michibata, H. and R. Hori. 1979. The accumulation of manganese from the environmental medium by the egg of *Oryzias latipes*. J. Cell Physiol., 98(1): 241-243. [ref. 240]
- Michibata, H., Y. Nojima and M. K. Kojima. 1987. Stage sensitivity of eggs of the teleost *Oryzias latipes* to cadmium exposure. Environ. Res. Sect. A., 42(2): 321-327. [ref. 241]
- Nagase, H., T. Hamasaki, T. Sato, H. Kito, Y. Yoshioka and Y. Ose. 1991. Structure-activity relationships for organotin compounds on the red killifish *Oryzias latipes*. Appl. Organomet. Chem., 5(2): 91-97. [ref. 247]
- Nakayama, K., Y. Oshima, T. Yamaguchi, Y. Tsuruda, I. J. Kang, N. Imada, T. Honjo and M. Kobayashi. 2004. Fertilization success and sexual behavior in male medaka, *Oryzias latipes*, exposed to tributyltin. Chemosphere, 55(10): 1331-1337. [ref. 251]
- Newsted, J. L., J. Nakanishi, I. Cousins, K. Werner and J. P. Giesy. 2002. Predicted distribution and ecological risk assessment of a "segregated" hydrofluoroether in the Japanese environment. Environ. Sci. Technol., 36(22): 4761-4769. [ref. 258]
- Nimrod, A. C. and W. H. Benson. 1998. Reproduction and development of Japanese medaka following an early life stage exposure to xenoestrogens. Aquatic Toxicol., 44(1-2): 141-156. [ref. 259]
- Nirmala, K., Y. Oshima, N. Imada, T. Honjo, K. Kobayashi and R. Lee. 1999. Transgenerational toxicity of tributyltin and its combined effects with polychlorinated biphenyls on reproductive processes in Japanese medaka

- (*Oryzias latipes*). Environ. Toxicol. Chem., 18(4): 717-721. [ref. 260]
- Oh, H. S., S. K. Lee, Y-H. Kim and J. K. Roh. 1991. Mechanism of selective toxicity of diazinon to killifish (*Oryzias latipes*) and loach (*Misgurnus anguillicaudatus*). ASTM. Spec. Tech. Publ., 1124: 343-353. [ref. 271]
- Onitsuka, S., Y. Kasai and K. Yoshimura. 1989. Quantitative structure-toxic activity relationship of fatty acids and the sodium salts to aquatic organisms. Chemosphere, 18(7-8): 1621-1631. [ref. 281]
- Ortego, L. S., A. C. Nimrod, G. R. Parsons, W. H. Benson and W. T. Brehm. 1994. Early life-stage effects in medaka (*Oryzias latipes*) following in ovo exposure to polyamine biosynthetic inhibitors. Ecotoxicol. Environ. Saf., 28(3): 329-339. [ref. 284]
- Owens, K. D. and K. N. Baer. 2000. Modifications of the topical Japanese Medaka (*Oryzias latipes*) embryo larval assay for assessing developmental toxicity of pentachlorophenol and p,p'-dichlorodiphenyltrichloroethane. Ecotoxicol. Environ. Saf., 47(1): 87-95. [ref. 287]
- Rao, S. S., T. A. Neheli, C. D. Metcalfe and B. Schmidt. 1997. Assessing the Toxicity of Environmental Contaminants with Early Life Stages of Japanese Medaka (*Oryzias latipes*). Environ. Toxicol. Water Qual., 12(4): 349-352. [ref. 292]
- Rice, P. J., C. D. Drewes, T. M. Klubertanz, J. R. Coats and S. P. Bradbury. 1997. Acute toxicity and behavioral effects of chlorpyrifos, permethrin, phenol, strychnine, and 2,4-dinitrophenol to 30-day-old Japanese medaka (*Oryzias latipes*). Environ. Toxicol. Chem., 16(4): 696-704. [ref. 293]
- Schlenk, D. and A. El-Alfy. 1998. Expression of branchial flavin-containing monooxygenase is directly correlated with salinity-induced aldicarb toxicity in the euryhaline fish (*Oryzias latipes*). Mar. Environ. Res., 46 (2-3) (Special Issue): 103-106. [ref. 311]
- Seki, M., H. Yokota, H. Matsubara, M. Maeda, H. Tadokoro and K. Kobayashi. 2004. Fish full life-cycle testing for androgen methyltestosterone on medaka (*Oryzias latipes*). Environ. Toxicol. Chem., 23(3): 774-781. [ref. 314]
- Shimura, R., Y. X. Ma, K. Ijiri, S. Nagaoka and M. Uchiyama. 2004. Nitrate toxicity on visceral organs of medaka fish, *Oryzias latipes*: aiming to raise fish from egg to egg in space. Biological Sciences in Space, 18(1): 7-12. [ref. 326]
- Shugart, L. R., J. F. McCarthy, S. J. D'surney, M. S. JR. Greeley and C. G. Hull. 1990. Molecular and cellular markers of toxicity in the Japanese Medaka *Oryzias latipes*. US. DOE. Rep., pp32. [ref. 329]
- Solomon, F. P. and E. M. Faustman. 1987. Developmental toxicity of four model alkylating agents on Japanese medaka fish (*Oryzias latipes*) embryos. Environ. Toxicol. Chem., 6(10): 747-753. [ref. 330]
- Solomon, H. M. and J. S. Weis. 1979. Abnormal circulatory development in medaka caused by the insecticides carbaryl, malathion and parathion. Teratology, 19(1): 51-62. [ref. 331]
- Spitsbergen, J. M. 1995. Cardiovascular toxicity of environmental contaminants to developing fish-molecular mechanisms. AD. Rep., pp27. [ref. 332]
- Tadokoro, H., M. Maeda, Y. Kawashima, M. Kitano, D. F. Hwang and T. Yoshida. 1991. Aquatic toxicity testing for multicomponent compounds with special reference to preparation of test solution. Ecotoxicol. Environ. Saf., 21(1): 57-67. [ref. 340]
- Takemoto, Y., Y. Ohshima, H. Yamada and J. Miyamoto. 1984. Fate of fenitrothion in several developmental stages of the killifish (*Oryzias latipes*). Arch. Environ. Contam. Toxicol., 13(5): 579-587. [ref. 346]
- Takimoto, Y., M. Ohshima and J. Miyamoto.

1987. Comparative metabolism of fenitrothion in aquatic organisms. I. Metabolism in the euryhaline fish, *Oryzias latipes* and *Mugil cephalus*. *Ecotoxicol. Environ. Saf.*, 13(1): 104-117. [ref. 356]
- Teather, K., M. Harris, J. Boswell and M. Gray. 2001. Effects of acrobat MZ(R) and tattoo C(R) on Japanese medaka (*Oryzias latipes*) development and adult male behavior. *Aquatic Toxicol.*, 51(4): 419-430. [ref. 360]
- Tsuda, T., S. Aoki, T. Inoue and M. Kojima. 1995. Accumulation of IBP, parathion and EPN by killifish: comparison of individual and mixed pesticides. *Comparative Biochemistry and Physiology C Pharmacology Toxicology and Endocrinology*, 111C(1): 19-22. [ref. 367]
- Tsuda, T., M. Kojima, H. Harada, A. Nakajima and S. Aoki. 1997. Relationships of bioconcentration factors of organophosphate pesticides among species of fish. *Comparative Biochemistry and Physiology C Pharmacology Toxicology and Endocrinology*, 116C(3): 213-218. [ref. 368]
- Tsuda, T., M. Kojima, H. Harada, A. Nakajima and S. Aoki. 1997. Acute toxicity, accumulation and excretion of isoprothiolane and its degradation products in killifish. *Water Res.*, 31(2): 323-327. [ref. 369]
- Tsuda, T., M. Kojima, H. Arada, A. Nakajima and S. Aoki. 1997. Acute toxicity, accumulation and excretion of organophosphorous insecticides and their oxidation products in killifish. *Chemosphere*, 35(5): 939-949. [ref. 370]
- Walker, W. W., C. S. Heard, K. Lotz, T. F. Lytle, W. E. Hawkins, D. H. Barnes and R. M. Overstreet. 1989. Tumorigenic, growth, reproductive, and developmental effects in medaka exposed to bis (tri-n-butyltin) oxide. *Oceans*, 1989(2): 516-524. [ref. 381]
- Walker, W. W., C. S. Manning, R. M. Overstreet and W. E. Hawkins. 1985. Development of aquarium fish models for environmental carcinogenesis: an intermittent-flow exposure system for volatile, hydrophobic chemicals. *J. Appl. Toxicol.*, 5(4): 255-260. [ref. 382]
- Wester, P. W. 1991. Histopathological effects of environmental pollutants β -HCH and methyl mercury on reproductive organs in freshwater fish. *Comp. Biochem. Physiol. C*, 100(1-2): 237-239. [ref. 389]
- Wester, P. W. and J. H. Canton. 1986. Histopathological study of *Oryzias latipes* (medaka) after long-term β -hexachlorocyclohexane exposure. *Aquatic Toxicol.*, 9(1): 21-45. [ref. 390]
- Wester, P. W., J. H. Canton, A. A. J. Van Iersel, E. I. Krajnc and H. M. A. G. Vaessen. 1990. The toxicity of bis (tri-n-butyltin) oxide (TBTO) and di-n-butyltindichloride (DBTC) in the small fish species *Oryzias latipes* (medaka) and *Poecilia reticulata* (guppy). *Aquatic Toxicol.*, 16(1): 53-72. [ref. 391]
- Wisk, J. D. and K. R. Cooper. 1990. Comparison of the toxicity of several polychlorinated dibenzo-p-dioxins and 2, 3, 7, 8-tetrachlorodibenzofuran in embryos of the Japanese medaka (*Oryzias latipes*). *Chemosphere*, 20(3-4): 361-377. [ref. 393]
- Wisk, J. D. and K. R. Cooper. 1990. The stage specific toxicity of 2, 3, 7, 8-tetrachlorodibenzo-p-dioxin in embryos of the Japanese medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.*, 9(9): 1159-1169. [ref. 394]
- Wisk, J. D. and K. R. Cooper. 1992. Effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on benzo(a)pyrene hydroxylase activity in embryos of the Japanese medaka (*Oryzias latipes*). *Arch. Toxicol.*, 66(4): 245-249. [ref. 395]
- Zhu, J. and L. Li. 1997. Acute toxicity of DMAH to *Oryzias latipes* as modified by some influential factors. *J. Environ. Sci. (China)*, 9(1): 20-26. [ref. 418]

- 安藤正典・西村哲治・神野透人・鈴木玲央・青木康展・有薦幸司・石橋弘志・今川正良・中室克彦. 2004. 水域環境における内分泌かく乱化学物質の次世代への影響評価法確立に関する分子遺伝学的研究. 環境保全研究成果集, 2002 (1) : 11.1-11.20. [ref. 422]
- 新家淳治・中川喜明・荒木恵一・早川修二・金丸豪. 1989. 化学物質の環境汚染に対する研究 水中微生物分解性試験及び魚類(ヒメダカ)急性毒性試験. 三重県環境科学センター研究報告, (9): 53-60. [ref. 426]
- 有薦幸司. 2003. 環境ホルモンの生態系への影響. NIRS-M, 169: 133-140. [ref. 427]
- 土井麻記子・深津和彦・篠原亮太・花嶋正孝・樋口壯太郎・友田啓二郎. 2004. 廃棄物埋立地浸出水の短期曝露によるヒメダカ(*Oryzias latipes*)への影響. 廃棄物学会論文誌, 15(4): 294-301. [ref. 431]
- 土井麻記子・高村嘉幸・深津和彦・鴻野雅一. 2004. アクアリウムを用いた生態系に対する排水の安全性評価に関する研究. 水環境学会誌, 27(5): 343-348. [ref. 432]
- 藤田肖子・石嶋直之・西内康浩. 1987. 水田除草剤に対する淡水ヒメダカと海水順化ヒメダカの感受性の相違. 農薬検査所報告, (27): 32-36. [ref. 441]
- 藤原茂樹・山田一裕・西村 修・須藤隆一. 1999. ヒメダカに対する界面活性剤とアンモニアの複合影響. 用水と排水, 41(7): 598-602. [ref. 442]
- 日高秀雄・八反田 誠・立川 涼. 1984. メダカによる農薬の忌避. 日本農芸化学会誌, 58(2): 145-151. [ref. 456]
- 日野隆信・藤曲正登・福田秀生. 1981. 六価クロムの生体影響に関する研究 ヒメダカを対象とした慢性毒性試験. 千葉県衛星研究所研究報告, 5: 58-60. [ref. 457]
- 堀尾 刚. 2003. 農業生態系の持つ自然循環機能に基づいた食料と環境の安全性の確保 メダカを使って化学物質の内分泌かく乱作用を簡易に検出する. 農業環境研究成果情報, 19: 28-29. [ref. 463]
- 春日清一. 1980. メダカにおけるカドミウム急性毒性の雌雄差. 日本水産学会誌, 46(9): 1073-1076. [ref. 492]
- 中川久機・石尾真弥. 1991. 重金属イオンに対するメダカの卵, 仔魚および成魚の感受性の比較. 水産増殖, 39(4): 435-440. [ref. 522]
- ビー・エス・ティー. 2004. メダカセンサー. クリモト技報, (50): 118-120. [ref. 544]
- 佐藤英毅. 1988. カダヤシ, メダカ, グッピーおよびフナにおける台所用洗剤に対する感受性. 川崎市衛生研究所年報, 23(1987): 81-82. [ref. 554]
- 茂岡忠義・山県民次・蓑田妙子・山内文雄. 1988. クロロフェノール類のヒメダカへの急性毒性と胚のふ化阻害性及び構造との相関. 衛生化学, 34(4): 343-349. [ref. 556]
- 島田敦子. 1985. メダカを用いた環境汚染の検定. 生物の科学 遺伝, 39(8): 34-36. [ref. 562]
- 富森聰子・吉川宏幸・長屋祐一・谷山鉄郎. 1994. ヒメダカにおよぼすゴルフ場使用農薬の複合致死性. 人間と環境, 20(3): 162-167, 169-171. [ref. 588]
- 上田成一・宮崎憲明・白井玄爾・山口道雄. 1995. 化学物質の魚類急性毒性および藻類生長阻害. 長崎県衛生公害研究所報, (40): 134, 135, 135(1). [ref. 594]
- 渡辺 志. 1992. モノアルキルフォスフェイト(MAP)のヒメダカ魚毒性に及ぼすアルミニウムイオンの影響. 合成洗剤研究会誌, 16(1): 13-14. [ref. 612]
- 山本義和・井上雅代・小林けい子・永井京子. 1982. ヒメダカに対するカドミウムの急性毒性に及ぼす水温の影響. 生態化学, 5(1): 31-35. [ref. 616]
- 山本義和・井上雅代・小林けい子・永井京子. 1982. ヒメダカに対するカドミウムの急性毒性に及ぼす水温の影響 特にへい死個体について. 生態化学, 5(2): 15-20. [ref. 617]
- 吉岡敏行・萩野泰夫・森 忠繁. 1995. 水生生物に対する界面活性剤および衣料洗剤の毒性評価. 全国公害研会誌, 20(3): 138-143. [ref. 620]

- 生殖・発生(Reproduction and Embryology)
- Aketa, K. 1962. Cytochemical observations on disappearance of ribonucleic acid-rich granules upon fertilization in egg of the medaka (*Oryzias latipes*). *Exp. Cell Res.*, 28: 254-259. [ref. 1]
- Asami, K., T. Hanai and A. Irimajiri. 1987. Development of medaka egg as monitored by their dielectric behavior. *Bull. Inst. Chem. Res. Kyoto Univ.*, 64(5-6): 339-343. [ref. 11]
- Awaji, M. and I. Hanyu. 1989. Seasonal changes in ovarian response to photoperiods in orange-red type medaka. *Zool. Sci. (Tokyo)*, 6(5): 943-950. [ref. 12]
- Awaji, M. and I. Hanyu. 1989. Temperature-photoperiod conditions necessary to begin the spawning season in wild type medaka. *Nippon Suisan Gakkaishi*, 55(4): 747. [ref. 13]
- Bentivegna, C. S. and T. Piatkowski. 1998. Effects of tributyltin on medaka (*Oryzias latipes*) embryos at different stages of development. *Aquatic Toxicol.*, 44(1-2): 117-128. [ref. 15]
- Billard, R. 1984. Ultrastructural change in the spermatogonia and spermatocytes of *Poecilia reticulata* during spermatogenesis. *Cell Tissue Res.*, 237(2): 219-226. [ref. 17]
- Cameron, I. L., K. E. Hunter and W. D. Winters. 1985. Retardation of embryogenesis by extremely low frequency 60Hz electromagnetic fields. *Physiol. Chem. Phys. Med. NMR.*, 17(1): 135-138. [ref. 25]
- Cameron, I. L., W. C. Lawrence and J. B. Lum. 1985. Medaka eggs as a model system for screening potential teratogens. *Prog. Clin. Biol. Res.*, 163C: 239-243. [ref. 26]
- Cheer, A., J. P. Vincent, R. Nuccitelli and G. Oster. 1987. Cortical activity in vertebrate eggs. I: The activation waves. *J. Theor. Biol.*, 124(4): 377-404. [ref. 29]
- Davis, C., M. Okihiro and D. Hinton. 2002.

- Effects of husbandry practices, gender, and normal physiological variation on growth and reproduction of Japanese medaka, *Oryzias latipes*. *Aquat. Toxicol.*, 60(3-4): 185-201. [ref. 33]
- Egami, N. 1954. Effect of artificial photoperiodicity on time of oviposition in the fish *Oryzias latipes*. *Annot. Zool. Japon.*, 27: 57-62. [ref. 38]
- Egami, N. 1959. Record of the number of eggs obtained from a single pair of *Oryzias latipes*, kept in laboratory aquarium. *J. Fac. Sci., Tokyo Univ. IV*, 8: 521-538. [ref. 41]
- Egami, N. 1959. Effect of exposure to low temperature on the time of oviposition and the growth of the oocytes in the fish, *Oryzias latipes*. *J. Fac. Sci., Tokyo Univ. IV*, 8: 539-548. [ref. 42]
- Evans, J. P., L. Zane, S. Francescato and A. Pilastro. 2003. Directional postcopulatory sexual selection revealed by artificial insemination. *Nature (Lond)*, 421(6921): 360-363. [ref. 46]
- Fluck, R. A. 1982. Localization of acetylcholinesterase activity in young embryos of the medaka *Oryzias latipes*, a teleost. *Comp. Biochem. Physiol. C.*, 72(1): 59-64. [ref. 50]
- Fluck, R. A., K. L. Krok, B. A. Bast, S. E. Michaud and C. E. Kim. 1998. Gravity influences the position of dorsoventral axis in medaka fish embryos (*Oryzias latipes*). *Dev. Growth Differ.*, 40(5): 509-518. [ref. 52]
- Fluck, R. A. and L. F. Jaffe. 1988. Electrical currents associated with rhythmic contractions of the blastoderm of the medaka, *Oryzias latipes*. *Comp. Biochem. Physiol. A.*, 89(4): 609-613. [ref. 53]
- Fukuda, S., Y. Nagahama, N. Sakai and S. Adachi. 1994. Steroidogenesis in the ovarian follicle of medaka (*Oryzias latipes*, a daily spawner) during oocyte maturation. *Dev. Growth Differ.*, 36(1): 81-88. [ref. 58]
- Gilkey, J. C. 1983. Roles of calcium and pH in

- activation of eggs of the medaka fish, *Oryzias latipes*. J. Cell Biol., 97(3): 669-678. [ref. 64]
- Hamaguchi, S. 1982. A light- and electron-microscopic study on the migration of primordial germ cells in the teleost, *Oryzias latipes*. Cell Tissue Res., 227(1): 139-151. [ref. 75]
- Hamaguchi, S. 1983. Asymmetrical development of the gonads in the embryos and fry of the fish, *Oryzias celebensis*. Dev. Growth Differ., 25(6): 553-561. [ref. 76]
- Hamaguchi, S. 1985. Changes in the morphology of the germinal dense bodies in primordial germ cells of the teleost, *Oryzias latipes*. Cell Tissue Res., 240(3): 669-673. [ref. 77]
- Hartley, W. R., A. Thiagarajah, M. B. Anderson, M. W. Broxson, S. E. Major and S. I. Zell. 1998. Gonadal development in Japanese Medaka (*Oryzias latipes*) exposed to 17,β-estradiol. Mar. Environ. Res., 46 (1-5): 145-148. [ref. 82]
- Hishida, T. and E. Nakano. 1954. Respiratory metabolism during fish development. Embryologia, 2: 67-79. [ref. 111]
- Hong, Y., T. Lui, H. Zhao, H. Xu, W. Wang, R. Lui, T. Chen, J. Deng and J. Gui. 2004. Establishment of a normal medakafish spermatogonial cell line capable of sperm production in vitro. Proc. Natl. Acad. Sci. U.S.A., 101(21): 8011-8016. [ref. 113]
- Hori, R. and K. Sato. 1973. The incorporation of 45 Ca into the egg of the medaka, *Oryzias latipes*. Experientia, 29(2): 144-145. [ref. 114]
- Hori, R. and T. Kono. 1974. On the content of sodium and potassium of the unfertilized egg of the medaka, *Oryzias latipes* and its changes accompanying fertilization. Protoplasma, 80(1): 149-153. [ref. 115]
- Hyodo, M., A. Aoki, C. Ando, M. Katsumata, S. Nyui, N. Motegi, T. Morozumi and M. Matsuhashi. 1996. Essential role of the yolk syncytial layer for the development of isolated blastoderms from medaka embryos. Dev. Growth Differ., 38(4): 383-392. [ref. 21]
- Hyodo-Taguchi, Y. and H. Etoh. 1983. The fecundity and fertility of medaka exposed to chronic γ-radiation in their embryonic stages. J. Radiat. Res., 24(3): 270-277. [ref. 122]
- Ijiri, K. 1980. Gamma-ray irradiation of the sperm of the fish *Oryzias latipes* and induction of gynogenesis. J. Radiat. Res., 21(3-4): 263-270. [ref. 126]
- Ijiri, K. 1998. Development of space-fertilized eggs and formation of primordial germ cells in the embryos of Medaka fish. Adv. Space Res., 21(8-9): 1155-1158. [ref. 127]
- Inoue, K. and Y. Takei. 2002. Diverse adaptability in *Oryzias* species to high environmental salinity. Zool. Sci., 19(7): 727-734. [ref. 135]
- Ito, S. 1959. On the spawning time of the teleostean fishes. Japanese J. Ecol., 9: 3. [ref. 144]
- Iwamatsu, T. 1965. On fertilizability of pre-ovulation eggs in the medaka, *Oryzias latipes*. Embryologia (Nagoya), 61: 327-336. [ref. 145]
- Iwamatsu, T. 1965. Effect of acetone on the cortical changes at fertilization of the egg of the medaka, *Oryzias latipes*. Embryologia (Nagoya), 9(1): 1-12. [ref. 146]
- Iwamatsu, T. 1994. Medaka oocytes rotate within the ovarian follicle during oogenesis. Dev. Growth Differ., 36(2): 177-186. [ref. 147]
- Iwamatsu, T. 1998. Studies on fertilization in the teleost, I .Dynamic responses of fertilized medaka eggs. Dev. Growth Differ., 40(5): 475-483. [ref. 148]
- Iwamatsu, T., A. Inden, K. Hirata and H. Uwa. 1984. Experiments on interspecific hybridization between *Oryzias latipes* and *Oryzias celebensis*. Zool. Sci., 1(4): 653-663. [ref. 149]
- Iwamatsu, T. and I. Ota. 1974. Cleavage initiating activities of sperm fractions

- injected into the egg of the medaka, *Oryzias latipes*. J. Exp. Zool., 187(1): 3-15. [ref. 151]
- Iwamatsu, T., T. Mori and R. A. Fluck. 1993. Mechanical dechorionation of fertilized eggs for experimental embryology in the medaka. Zool. Sci., 10(6): 945-951. [ref. 154]
- Iwamatsu, T., T. Ohta, E. Oshima and N. Sakai. 1988. Oogenesis in the medaka *Oryzias latipes* - Stages of oocyte development. Zool. Sci., 5(2): 353-373. [ref. 155]
- Iwamatsu, T., T. Ohta, E. Oshima and T. Sugiura. 1985. Requirement of extracellular calcium ions for early fertilization events in the medaka egg. Dev. Growth Differ., 27(6): 751-762. [ref. 156]
- Iwamatsu, T., Y. Tada and Y. Nishiyama. 1982. On sperm nuclei and development in polyspermic eggs of the teleost, *Oryzias latipes*. Annot. Zool. Jpn., 55(2): 91-99. [ref. 157]
- Iwata, A., M. Iwata and E. Nakano. 1986. Distribution of fibronectin, microtubules and microfilaments in the melanophore of the medaka, *Oryzias latipes*. Cell Struct. Funct., 11(2): 99-107. [ref. 160]
- Jacobson, A. G. 1988. Somitomeres: mesodermal segments of vertebrate embryos. Development, 104(Suppl): 209-220. [ref. 161]
- Kanamori, A., Y. Nagahama and N. Egami. 1985. Development of the tissue architecture in gonads of the medaka *Oryzias latipes*. Zool. Sci., 2(5): 695-706. [ref. 168]
- Kawahara, T. and I. Yamashita. 2000. Estrogen-independent ovary formation in the medaka fish, *Oryzias latipes*. Zool. Sci., 17(1): 65-68. [ref. 176]
- Kobayashi, H. and T. Iwamatsu. 2000. Development and fine structure of the yolk nucleus of previtellogenic oocytes in medaka *Oryzias latipes*. Dev. Growth Differ., 42(6): 623-631. [ref. 193]
- Koya, Y. and A. Iwase. 2004. Annual reproductive cycle and rate of the spermatogenic process in male mosquitofish *Gambusia affinis*. Ichthyol. Res., 51(2): 131-136. [ref. 207]
- Koya, Y., M. Inoue, T. Naruse and S. Sawaguchi. 2000. Dynamics of oocyte and embryonic development during ovarian cycle of the viviparous mosquitofish *Gambusia affinis*. Fish Sci., 66(1): 63-70. [ref. 208]
- Kuwahara, Y., A. Shimada, H. Mitani and A. Shima. 2002. A Critical stage in spermatogenesis for radiation-induced cell death in Medaka Fish, *Oryzias latipes*. Radiat. Res., 157(4): 386-392. [ref. 215]
- Martindale, M. Q., S. Meier and A. G. Jacobson. 1987. Mesodermal metamерism in the teleost, *Oryzias latipes* (the medaka). J. Morphol., 193(3): 241-252. [ref. 223]
- Masuda, K., I. Iuchi and K. Yamagami. 1991. Analysis of hardening of the egg envelope (chorion) of the fish, *Oryzias latipes*. Dev. Growth Differ., 33(1): 75-83. [ref. 230]
- Michibata, H. and R. Hori. 1979. The accumulation of manganese from the environmental medium by the egg of *Oryzias latipes*. J. Cell Physiol., 98(1): 241-243. [ref. 240]
- Nakamoto, M. and N. Shibata. 2003. Cloning and expression of daxi in Medaka gonad. Zool. Sci., 20(12): 1559. [ref. 248]
- Nuccitelli, R. 1980. The electrical changes accompanying fertilization and cortical vesicle secretion in the medaka egg. Dev. Biol., 76(2): 483-498. [ref. 263]
- Nuccitelli, R. 1980. The fertilization potential is not necessary for the block to polyspermy or the activation of development in the medaka egg. Dev. Biol., 76(2): 499-504. [ref. 264]
- Oba, Y., M. Yoshikuni, Y. Nagahama and M. Mita. 1997. Inhibitory guanine-nucleotide-binding-regulatory protein α subunits in

- medaka (*Oryzias latipes*) oocytes. cDNA cloning and decreased expression of proteins during oocytes maturation. Eur. J. Biochem., 249(3): 846-853. [ref. 265]
- Ogiwara, K., M. Shinohara and T. Takahashi. 2002. Identification of the follicles digestion factor using the medaka fish in vitro ovulation system. Zool. Sci., 19(12): 1547. [ref. 269]
- Patyna, P. J., K. R. Cooper, R. A. Davi, T. F. Parkerton and R. P. Brown. 1999. A proposed multigeneration protocol for Japanese medaka (*Oryzias latipes*) to evaluate effects of endocrine disruptors. Sci. Total Environ., 233(1-3): 211-220. [ref. 290]
- Ridgway, E. B., J. C. Gilkey and L. F. Jaffe. 1977. Free calcium increases explosively in activating medaka eggs. Proc. Natl. Acad. Sci. U.S.A., 74(2): 623-627. [ref. 294]
- Saiki, A., M. Tamura, M. Matsumoto, J. Katowgi, A. Watanabe and K. Onitake. 1997. Establishment of in vitro spermatogenesis form spermatocytes in the medaka, *Oryzias latipes*. Dev. Growth Differ., 39(3): 337-344. [ref. 302]
- Sakai, N., T. Iwamatsu, K. Yamauchi and Y. Nagahama. 1987. Development of the steroidogenic capacity of medaka (*Oryzias latipes*) ovarian follicles during vitellogenesis and oocyte maturation. Gen. Comp. Endocrinol., 66(3): 333-342. [ref. 303]
- Sguigna, C., R. Fluck and B. Barber. 1988. Calcium dependence of rhythmic contractions of the *Oryzias latipes* blastoderm. Comp. Biochem. Physiol. C., 89(2): 369-374. [ref. 315]
- Shibata, N. and S. Hamaguchi. 1988. Evidence for the sexual bipotentiality of spermatogonia in the fish, *Oryzias latipes*. J. Exp. Zool., 245(1): 71-77. [ref. 316]
- Shima, A. and A. Shimada. 1991. Development of a possible nonmammalian test system for radiation-induced germ-cell mutagenesis using a fish, the Japanese medaka (*Oryzias latipes*). Proc. Natl. Acad. Sci. U.S.A., 88(6): 2545-2549. [ref. 318]
- Shima, A. and A. Shimada. 2001. The medaka as a model for studying germ-cell mutagenesis and genomic instability. Mar. Biotechnol., 3(Supplement 1): S162-S167. [ref. 319]
- Shima, A. and N. Egami. 1978. Absence of systematic polyploidization of hepatocyte nuclei during the ageing process of the male medaka, *Oryzias latipes*. Exp. Gerontol., 13(1-2): 51-55. [ref. 321]
- Shimada, Y. and N. Egami. 1984. The unique responses of the primordial germ cells in the fish *Oryzias latipes* to gamma-rays. Int. J. Radiat. Biol. Relat. Stud. Phys. Chem. Med., 45(3): 227-235. [ref. 324]
- Srivastava, P. N. 1966. Effect of ionizing radiation on the ovaries of Japanese medaka, *Oryzias latipes* (t. et s.). Acta. Anat (Basel), 63(3): 434-444. [ref. 333]
- Suyama, I., H. Etoh, T. Maruyama, Y. Kato and R. Ichikawa. 1981. Effects of ionizing radiation on the early development of *Oryzias* eggs. J. Radiat. Res., 22(1): 125-133. [ref. 336]
- Suzuki, A. and N. Shibata. 2004. Developmental process of genital ducts in the medaka, *Oryzias latipes*. Zool. Sci., 21(4): 397-406. [ref. 337]
- Suzuki, A., N. Shibata, T. Sato, M. Sakaizumi and S. Hamaguchi. 2003. Early development of gonads and genital ducts in secondary sexual character less mutant. Scl. of the Medaka. Zool. Sci., 20(12): 1604. [ref. 339]
- Takahashi, H. and Y. Iwasaki. 1973. Histochemical demonstration of delta 5-3 beta-hydroxysteroid dehydrogenase activity in the testis of the medaka, *Oryzias latipes*. Endocrinol. Jpn., 20(5): 529-533. [ref. 341]
- Takeuchi, K. 1967. Discrimination of genetic sex in embryos of d-rR strain of the medaka (*Oryzias latipes*). Experientia, 23(7): 569-

570. [ref. 353]
- Urasaki, H. 1976. The role of pineal and eyes in the photoperiodic effect on the gonad of the medaka, *Oryzias latipes*. Chronobiologia, 3(3): 228-234. [ref. 375]
- Uwa, H. 1965. Gynogenetic haploid embryos of the medaka (*Oryzias latipes*). Embryologia (Nagoya), 9(1): 40-48. [ref. 377]
- Watanabe, A., E. Kobayashi, T. Ogawa and K. Onitake. 1998. Fibroblast growth factor may regulate the initiation of oocytes growth in the developing ovary of the medaka, *Oryzias latipes*. Zool. Sci., 15(4): 531-536. [ref. 386]
- Yamagami, K. 1975. Relationship between two kinds of hatching enzymes in the hatching liquid of the medaka, *Oryzias latipes*. J. Exp. Zool., 192(1): 127-132. [ref. 398]
- Yamamoto, K. 1962. Origin of the yearly crop of eggs in the medaka, *Oryzias latipes*. Annot. Zool. Japon., 35: 156-161. [ref. 399]
- Yamamoto, T. 1956. The physiology of fertilization in the medaka (*Oryzias latipes*). Exp. Cell Res., 10(2): 387-393. [ref. 400]
- Yamamoto, T. 1958. Artificial induction of functional sex-reversal in genotypic females of the medaka (*Oryzias latipes*). J. Exp. Zool., 137(2): 227-263. [ref. 401]
- Yamamoto, T. 1959. The effects of estrone dosage level upon the percentage of sex-reversals in genetic male (XY) of the medaka (*Oryzias latipes*). J. Exp. Zool., 141: 133-153. [ref. 402]
- Yoshioka, H. 1962. On the effects of environmental factors upon the reproduction of fishes. I. The effects of day-length on the reproduction of the Japanese killifish, *Oryzias latipes*. Bull. Fac. Fish., Hokkaido Univ., 13: 123-136. [ref. 415]
- Yoshioka, H. 1966. On the effects of environmental factors upon the reproduction of fishes. III. The occurrence and regulation of refractory period in the photoperiodic response of medaka, *Oryzias latipes*. J. Hokkaido Univ. Educ. 17: 23-33. [ref. 416]
- 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp. [ref. 436]
- 江上信雄・山上健次郎・嶋 昭経. 1990. メダカの生物学. 東京大学出版会, 315pp. [ref. 437]
- 古屋康則・石川早苗・沢口小有美. 2004. カダヤシ *Gambusia affinis* の卵巣周期に及ぼす水温および光周期の影響. 魚類学雑誌, 51(1): 43-50. [ref. 448]
- 堀尾 剛・伊藤文成. 2001. 成育初期段階のメダカに対する内分泌かく乱物質作用に関する研究. 環境化学討論会予稿集, 10: 358-359. [ref. 463]
- 岩松鷹司. 1981. 異種精子によるメダカ (*Oryzias latipes*) 卵の発生. 愛知教育大学研究報告 自然科学, 30: 141-151. [ref. 474]
- 岩松鷹司. 1985. メダカ卵の実験的操作. 生物の科学 遺伝, 39(8): 42-46. [ref. 475]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]
- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 岩松鷹司. 1993. メダカ卵における第一卵割面とはい軸について. 愛知教育大学研究報告 自然科学, 42: 79-86. [ref. 477]
- 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp. [ref. 479]
- 金森 章. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解をめざしてⅣ. ゲノム, 進化, 種分化 メダカを用い脊椎動物の生殖巣の性分化を探る 蛋白質 核酸 酶素, 45(17): 2949-2953. [ref. 485]
- 小林重光. 2002. メダカはいにおける脈管系の発生について. 医学検査, 51(6): 915-919. [ref. 499]
- 小林哲邦・菱田富雄. 1989. 卵膜除去したメダカ胚に対するメチルテストステロンの影響 特に生殖細胞数の変化に関連して. 朝日大学教養部研究報告, 15: 113-121. [ref. 500]
- Lin B-L・東海明宏・吉田喜久雄・米沢義堯・富永 衛・荻野 哲・笠島通夫・芦田昭二・

- 岩松鷹司. 2003. メダカ (*Oryzias latipes*) 精巣卵の定量的検出のための新手法（小片化法）. 水環境学会誌, 26(11): 725-730. [ref. 507]
- 長尾和右・井尻憲一・西脇優子. 1999. 3Dクリノスタットによる疑似微小重力下におけるメダカはい発生 網膜および軟骨形成. 宇宙生物科学, 13(3): 162-163. [ref. 520]
- 小村潤一郎. 1985. メダカの細胞培養. 生物の科学 遺伝, 39(8): 28-30. [ref. 535]
- 大沢一爽. 1982. メダカの実験 - 33章. 共立出版, 160pp. [ref. 536]
- 太田忠之・岩松鷹司. 1981. 精子ペん毛微細管による魚卵の卵割誘導. 生物の科学 遺伝, 35(6): 43-45. [ref. 539]
- 佐々木 昆. 1993. 小さないのち—メダカの誕生. フレーベル館, 32pp. [ref. 553]
- 嶋 昭紘・島田敦子. 1995. メダカ生殖細胞の突然変異. 共用設備管理部門総合研究部門年報, 22: 31-34. [ref. 559]
- 嶋 昭紘・島田敦子. 1996. メダカ生殖細胞の突然変異. 共用設備管理部門総合研究部門年報, 23: 27-30. [ref. 560]
- 嶋 昭紘・島田敦子. 1997. メダカ生殖細胞の突然変異. 共用設備管理部門総合研究部門年報, 24: 21-24. [ref. 561]
- 竹内邦輔. 1981. 誰にでもできるメダカの実験. 新光印刷株式会社, 110pp. [ref. 578]
- 生 理 (Physiology)
- Aoki, K. and H. Umeura. 1970. Cell types in the pituitary of the medaka, *Oryzias latipes*. Endocrinol. Jpn., 17(1): 45-55. [ref. 5]
- Anraku, E., T. Iwamatsu and T. Ohta. 1989. Effects of forskolin on fine structure of medaka follicles. Dev. Growth Differ., 31(1): 45-53. [ref. 8]
- Asahina, K., A. Urabe, T. Sakai, H. Hirose and H. Hibiya. 1989. Effects of various androgens on the formation of papillary processes on the anal fin rays in the female medaka *Oryzias latipes*. Bull. Jpn. Soc. Sci. fish, 55(10): 1871. [ref. 10]
- Cameron, I. L. and K. E. Hunter. 1984. Regulation of the permeability of the medaka fish embryo chorion by exogenous sodium and calcium ions. J. Exp. Zool., 231(3): 447-454. [ref. 24]
- Chan, K. K. 1976. A photosensitive daily rhythm in the female medaka, *Oryzias latipes*. Can. J. Zool., 54(6): 852-856. [ref. 28]
- Davis, C., M. Okihiro and D. Hinton. 2002. Effects of husbandry practices, gender, and normal physiological variation on growth and reproduction of Japanese medaka, *Oryzias latipes*. Aquat. Toxicol., 60(3-4): 185-201. [ref. 33]
- Featherstone, D., C. D. Drewes and J. R. Coats. 1991. Noninvasive detection of electrical events during the startle response in larval medaka. J. Exp. Biol., 158: 583-589. [ref. 48]
- Fluck, R. A., A. L. Miller, V. C. Abraham and L. F. Jaffe. 1994. Calcium buffer injections inhibit ooplasmic segregation in medaka eggs. Biol. Bull (Woods Hole), 186(3): 254-262. [ref. 51]
- Ghoneum, M., N. Egami and K. Ijiri. 1988. Effect of oestrogen treatment on the survival of irradiated fish of the species *Oryzias latipes*. Folia. Morphol (praha)., 36(1): 25-31. [ref. 63]
- Gilkey, J. C. 1983. Roles of calcium and pH in activation of eggs of the medaka fish, *Oryzias latipes*. J. Cell Biol., 97(3): 669-678. [ref. 64]
- Gonzalez-doncel, M., C. Barrueco, E. De, La, Pena and D. E. Hinton. 2003. Stage sensitivity of medaka (*Oryzias latipes*) eggs and embryos to permethrin. Aquatic Toxicol., 62(3): 255-268. [ref. 65]
- Gu, Y., S. Yasumasu and I. Iuchi. 2003. 3D-structure of hce, one of Medaka hatching enzymes, and analysis of the mechanism of its egg envelope digestion using metated recombinant hce. Biochim. Biophys. Acta., 1618(2): 95-105. [ref. 72]
- Ha, C. R. and I. Iuchi. 1996. Participation of a

- metalloprotease in the fertilization associated conversion of the egg envelope (chorion) of the fish, *Oryzias latipes*. Dev. Growth Differ., 38(5): 383-396. [ref. 73]
- Hamaguchi, S. and M. Sakaizumi. 1992. Sexually differentiated mechanisms of sterility in interspecific hybrids between *Oryzias latipes* and *O. curvinotus*. J. Exp. Zool., 263(3): 323-329. [ref. 79]
- Hara, T., S. Hagino and S. Hosokawa. 2004. Quantification of vitellogenin in several developmental stages of medaka (*Oryzias latipes*) S-rR strain. Environ. Sci. (Tokyo), 11(4): 221-230. [ref. 81]
- Hishida, T. 1965. Accumulation of estrone-16-C-14 and diethylstilbestrolmonoethyl-1-C-14 in larval gonads of the Medaka, *Oryzias latipes*, and determination of the minimum dosage of estrogen for sex reversal. Gen. Comp. Endocrinol., 56: 137-144. [ref. 110]
- Hong, Y., T. Liu, H. Zhao, H. Xu, W. Wang, R. Liu, T. Chen, J. Deng and J. Gui. 2004. Establishment of a normal medakafish spermatogonial cell line capable of aperm production in vitro. Proc. Natl. Acad. Sci. U.S.A., 101(21): 8011-8016. [ref. 113]
- Inoue, K., K. Naruse, N. Suzuki and Y. Takei. 2003. Four distinct C-type natriuretic peptides found in Medaka reveal new evolutionary history of the natriuretic peptide family. Zool. Sci., 20(12): 1599. [ref. 133]
- Inoue, K. and Y. Takagi. 2001. Seawater adaptability of medaka (*Oryzias latipes*) and three related species. Zool. Sci., 18(Supplement): 115. [ref. 134]
- Inoue, K. and Y. Takei. 2002. Diverse Adaptability in *Oryzias* Species to High Environmental Salinity. Zool. Sci., 19(7): 727-734. [ref. 135]
- Ishikawa, Y. and Y. Hyodo-Taguchi. 1994. Cranial nerves and brain fiber systems of the medaka fry as observed by a whole-mount staining method. Neurosci. Res., 19(4): 379-386. [ref. 141]
- Iwamatsu, T., S. Y. Takahashi, N. Sakai, Y. Nagahama and Y. Onitake. 1987. Induction and inhibition of in vitro oocyte maturation and production of steroids in fish follicles by forskolin. J. Exp. Zool., 241(1): 101-111. [ref. 153]
- Iwamatsu, T., Y. Yoshimoto and Y. Hiramoto. 1988. Cytoplasmic Ca²⁺ release induced by microinjection of Ca²⁺ and effects of microinjected divalent cations on Ca²⁺ sequestration and exocytosis of cortical alveoli in the medaka egg. Dev. Biol., 125(2): 451-457. [ref. 158]
- Jaffe, L. F. 1980. Calcium explosions as triggers of development. Ann. NY. Acad. Sci., 339: 86-101. [ref. 163]
- Jaffe, L. F. 1983. Sources of calcium in egg activation: a review and hypothesis. Dev. Biol., 99(2): 265-276. [ref. 164]
- Ka-Sing, Chan, K. 1977. Effect of synthetic luteinizing hormone-releasing hormone (LH-RH) on ovarian development in Japanese medaka, *Oryzias latipes*. Can. J. Zool., 55(1): 155-160. [ref. 172]
- Kawamoto, N. 1969. Effects of androstenedione, 19-norethynodrel, progesterone, and medaka, *Oryzias latipes*. Dev. Growth Differ., 11(2): 89-103. [ref. 180]
- Kubota, Y. and A. Shima. 1991. Comparative study of embryonic thermoresistance of two inbred strains of the medaka (*Oryzias latipes*). J. Comp. Physiol. B. Biochem. Stst. Environ. Physiol., 160(6): 621-625. [ref. 209]
- Mihoko, Y. and M. Sugimoto. 2002. Is reduction of size required for the apoptosis in fish melanophores? Zool. Sci., 19(12): 1473. [ref. 242]
- Morishita, F. 1987. Responses of the melanophores of the medaka, *Oryzias latipes*, to adrenergic drugs: evidence for involvement of alpha 2 adrenergic receptors mediating melanin aggregation.

- Comp. Biochem. Physiol. C., 88(1): 69-74. [ref. 245]
- Murshall, W. S. 2003. Rapid regulation of NaCl secretion by estuarine teleost fish: coping strategies for short-duration freshwater exposures. *Biochim. Biophys. Acta.*, 1618(2): 95-105. [ref. 246]
- Nakamura, H., K. Ozato, Y. Wakamatsu and T. Iwamatsu. 2003. Normal growth of the "see-through" medaka. *Zool. Sci.*, 20(5): 607-615. [ref. 249]
- Namoto, S. 1987. A subtype of adenosine receptors mediating pigment dispersion in leucophores of the medaka: evidence for an A₂-receptor. *Comp. Biochem. Physiol. C.*, 88(1): 75-81. [ref. 252]
- Niwa, Hs. 1965. Effects of castration and administration of methyl testosterone on the nuptial coloration of the Medaka (*Oryzias latipes*). *Embryologia* (Nagoya), 61: 289-298. [ref. 261]
- Niwa, Hs. 1965. Inhibition by estradiol of methyl testosterone-induced nuptialcoloration the Medaka (*Oryzias latipes*). *Embryologia* (Nagoya), 61: 299-307. [ref. 262]
- Ohya, T. and S. Hayashi. 2003. Subnuclei structure of vasotocinergic (VT) neurons preopticus of the mature Medaka: presence of sexual dimorphism. *Neurosci. Res.*, 46(Supplement): 78. [ref. 273]
- Oikawa, T. 1971. Histochemical and physiological study of chrmaffin cells in the skin of the medaka, *Oryzias latipes*. *Dev. Growth Differ.*, 13(2): 125-130. [ref. 274]
- Okamoto, H. and J. Y. Kuwada. 1991. Outgrowth by fin motor axons in wildtype and a finless mutant of the Japanese medaka fish. *Dev. Biol.*, 146(1): 49-61. [ref. 275]
- Okamoto, H. and J. Y. Kuwada. 1991. Alteration of pectoral fin nerves following ablation of fin buds and by ectopic fin buds in the Japanese medaka fish. *Dev. Biol.*, 146(1): 62-71. [ref. 276]
- Okubo, K., K. Naruse, K. Aida, H. Mitani, M. Kondo and A. Shima. 2002. Structural and functional characterization of gonadotropin-releasing hormones and their receptors in the medaka *Oryzias latipes*. *Zool. Sci.*, 19(12): 1419-1420. [ref. 277]
- Onitake, K. and T. Iwamatsu. 1986. Immunocytochemical demonstration of steroid hormones in the granulose cells of the medaka, *Oryzias latipes*. *J. Exp. Zool.*, 239(1): 97-103. [ref. 280]
- Oshima, N., H. Kasukawa, R. Fujii, B. C. Wilkes, V. J. Hruby and M. E. Hadley. 1986. Action of melanin-concentrating hormone (MCH) on teleost chromatophores. *Gen. Comp. Endocrinol.*, 64(3): 381-388. [ref. 285]
- Sakaizumi, M., M. Takeda, H. Otake, K. Togashi, A. Shinomiya, S. Hamaguchi, M. Matsuda and Y. Nagahama. 2002. Mutations affecting sex determination or differentiation from wild populations of the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(12): 1497. [ref. 306]
- Sato, T., M. Sakaizumi and S. Hamaguchi. 2002. Endocrine mechanisms on onset of vitellogenesis in spring in the viviparous teleost, *Gambusia affinis*. *Zool. Sci.*, 19(12): 1487. [ref. 308]
- Schoots, A. F., R. C. Meijer and J. M. Denude. 1983. Dopaminergic regulation of hatching in fish embryos. *Dev. Biol.*, 100(1): 59-63. [ref. 313]
- Seki, M., H. Yokota, H. Matsumura, M. Maeda, H. Tadokoro and K. Kobayashi. 2004. Fish full life-cycle testing for androgen methyltestosterone on Medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.*, 23(3): 774-781. [ref. 314]
- Shimizu, K., S. Sawaguchi, Y. Koya and A. Shimizu. 2002. Endocrine mechanisms on onset of vitellogenesis in spring in the viviparous teleost, *Gambusia affinis*. *Zool. Sci.*, 19(12): 1487. [ref. 325]
- Sugimoto, M. 1993. Morphological color changes in the medaka, *Oryzias latipes*, after

- prolonged background adaptaion- I . Changes in the population morphology of melanophores. Comparative biochemistry and physiology A Comparative Physiology, 104(3): 513-518. [ref. 334]
- Suzuki, A. and N. Shibata. 2004. Developmental process of genital ducts in the medaka, *Oryzias latipes*. Zool. Sci., 21(4): 397-406. [ref. 337]
- Takeuchi, K. 1967. Large tooth formation in female medaka, *Oryzias latipes*, given methyl testosterone. J. Dent., 46(4): 750. [ref. 351]
- Tohya, S., A. Mochizuki and Y. Iwasa. 2000. A cell-sorting model can explain cone mosaic pattern of Medaka. Proc. 1st. Int. Conf. Syst., 2000: 239-244. [ref. 362]
- Toledo, C., J. Hendricks, P. Loveland, J. Wilcox and G. Bailey. 1987. Metabolism and DNA-binding in vivo of aflatoxin B1 in medaka (*Oryzias latipes*). Comp. Biochem. Physiol. C., 87(2): 275-281. [ref. 363]
- Ueda, H., G. Young and Y. Nagahama. 1983. Immunocytochemical identification of thyrotropin (TSH)-producing cells in pituitary glands of several species of teleosts with antiserum to human TSH beta subunit. Cell Tissue Res., 231(1): 199-204. [ref. 373]
- Uematsu, K. 1990. An analysis of sufficient stimuli for the oviposition in the medaka *Oryzias latipes*. J. Fac. Appl. Biol. Sci. Hiroshima Univ., 29(2): 109-116. [ref. 374]
- Utida, S., S. Hatai, T. Hirano and F. I. Kamemoto. 1971. Effect of prolactin on survival and plasma sodium levels in hypophysectomized medaka *Oryzias latipes*. Gen. Comp. Endocrinol., 16(3): 566-573. [ref. 376]
- Uwa, H. 1968. Hormonal inhibitions of ethisterone-induced process formation in adult females of the medaka, *Oryzias latipes*. Embryologia (Nagoya), 10(2): 147-154. [ref. 378]
- Wyban, J. A. 1982. Solube peptidase isozymes of the Japanese Medaka (*Oryzias latipes*): tissue distributions and substrate specificities. Biochem. Genet., 20(9-10): 849-858. [ref. 396]
- Yamamoto, T. 1968. Effects of 17-alpha-hydroxyprogesterone and androstenedione upon sex differentiation in the medaka, *Oryzias latipes*. Gen. Comp. Endocrinol., 10(1): 8-13. [ref. 408]
- Yamamoto, T., K. Takeuchi and M. Takai. 1968. Male-inducing action of androsterone and testosterone propionate upon XX zygotes in the medaka, *Oryzias latipes*. Embryologia (Nagoya), 10(2): 116-125. [ref. 409]
- Yamamoto, T. and N. Matsuda. 1963. Effects of estradiol and some alkycarbonyl androstanes upon sex differentiation in the medaka. *Oryzias latipes*. Gen. Comp. Endocrinol., 3: 101-110. [ref. 410]
- Yasumasu, S. 2001. Active hatching enzyme, HCE, of medaka prepared by an E. coli expression system. Zool. Sci., 18(Supplement): 67. [ref. 412]
- 安藤正典・西村哲治・神野透人・鈴木玲央・青木康展・有薗幸司・石橋弘志・今川正良・中室克彦. 2004. 水域環境における内分泌かく乱化学物質の次世代への影響評価法確立に関する分子遺伝学的研究. 環境保全研究成果集, 2002 (1) : 11.1-11.20. [ref. 422]
- 有薗幸司. 2003. 環境ホルモンの生態系への影響. NIRS-M, 169: 133-140. [ref. 427]
- 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp. [ref. 436]
- 江上信雄・山上健次郎・鷗 昭絃. 1990. メダカの生物学. 東京大学出版会, 315pp. [ref. 437]
- 古田寿昭. 2002. 生体系の非破壊分析(2)光で生理活性をオン-オフできるスイッチBhcケージド化合物の設計、合成そして応用. 現代化学, 378: 24-31. [ref. 447]
- 岩松鷹司. 1993. メダカ学. サイエンティスト社, 324pp. [ref. 476]

- 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp. [ref. 478]
- 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp. [ref. 479]
- 工藤 明. 2002. 重力研究に適した実験動物、メダカの骨形成とヒレ再生. 宇宙ステーション利用計画ワークショップ予稿集, 24(2): 133-134. [ref. 505]
- Lin B-L・東海明宏・吉田喜久雄・米沢義堯・富永 衛・荻野 哲・籠島通夫・芦田昭二・岩松鷹司. 2004. メダカ(*Oryzias latipes*)第二次性徴による量的内分泌搅乱影響評価のための有効な手法(分節計数法). 水環境学会誌, 27(1): 47-51. [ref. 508]
- たたら迫典久. 2003. メダカを用いた内分泌搅乱化学物質の評価法. ファルマシア, 39(9): 881-883. [ref. 586]
- 豊泉龍児・小笠原 強. 2001. 尾(尾鰭)から肢(他の鰭)への分化転換個体の作出を目的とした、有尾両生類の肢ならびに魚類の鰭の再生現象に関する研究. 神奈川大学総合理学研究所年報, 2000: 179-181. [ref. 589]
- 山本時男 (編). 1975. メダカの生物学と品種. 佑学社. [ref. 615]
4. Allinson, G., M. Ueoka and M. Morita. 1994. Effect of dietary 1,3,6,8-tetrachlorodibenzo-p-dioxin on the Japanese freshwater fish *Oryzias latipes* (Medaka) and aquatic snail *Indoplanorbis exustus* (Indohiramakigai). Chemosphere, 28(7): 1369-1383.
5. Aoki, K. and H. Matsudaira. 1984. Factors influencing methylazoxymethanol acetate initiation of liver tumors in *Oryzias latipes*: carcinogen dosage and time of exposure. Natl. Cancer Inst. Monogr., 65: 345-351.
6. Aoki, K. and H. Umeura. 1970. Cell types in the pituitary of the medaka, *Oryzias latipes*. Endocrinol. Jpn., 17(1): 45-55.
7. Aoki, K., M. Okamoto, K. Tatsumi and Y. Ishikawa. 1997. Cryopreservation of medaka spermatozoa. Zool. Sci., 14(4): 641-644.
8. Anraku, E., T. Iwamatsu and T. Ohta. 1989. Effects of forskolin on fine structure of medaka follicles. Dev. Growth Differ., 31(1): 45-53.
9. Araki, K., H. Okamoto, I. Nakayama, H. Nagoya and A. C. Graveson. 2001. Analysis of haploid development based on expression patterns of developmental genes in the medaka *Oryzias latipes*. Dev. Growth Differ., 43(5): 591-599.
10. Asahina, K., A. Urabe, T. Sakai, H. Hirose and H. Hibiya. 1989. Effects of various androgens on the formation of papillary processes on the anal fin rays in the female medaka *Oryzias latipes*. Bull. Jpn. Soc. Sci. fish, 55(10): 1871.
11. Asami, K., T. Hanai and A. Irimajiri. 1987. Development of medaka egg as monitored by their dielectric behavior. Bull. Inst. Chem. Res. Kyoto Univ., 64(5-6): 339-343.
12. Awaji, M. and I. Hanyu. 1989. Seasonal changes in ovarian response to photoperiods in orange-red type medaka. Zool. Sci (Tokyo), 6(5): 943-950.
13. Awaji, M. and I. Hanyu. 1989. Temperature-photoperiod conditions necessary to begin

総合目録 (General list)

1. Aketa, K. 1962. Cytochemical observations on disappearance of ribonucleic acid-rich granules upon fertilization in egg of the medaka (*Oryzias latipes*). Exp. Cell Res., 28: 254-259.
2. Ali, M. Y. and C. C. Lindsey. 1974. Heritable and temperature-induced meristic in the medaka, *Oryzias latipes*. Can. J. Zool., 52(8): 959-976.
3. Allinson, G. and M. Morita. 1995. Bioaccumulation and toxic effects of elevated of 3,3',4,4'-tetrachloroazobenzene (33'44'-TCAB) towards aquatic organisms. 2: Bioaccumulation and toxic effects of dietary 33'44'-TCAB on the Japanese medaka (*Oryzias latipes*). Chemosphere, 30(2): 223-232.
4. Asahina, K., A. Urabe, T. Sakai, H. Hirose and H. Hibiya. 1989. Effects of various androgens on the formation of papillary processes on the anal fin rays in the female medaka *Oryzias latipes*. Bull. Jpn. Soc. Sci. fish, 55(10): 1871.
5. Asami, K., T. Hanai and A. Irimajiri. 1987. Development of medaka egg as monitored by their dielectric behavior. Bull. Inst. Chem. Res. Kyoto Univ., 64(5-6): 339-343.
6. Awaji, M. and I. Hanyu. 1989. Seasonal changes in ovarian response to photoperiods in orange-red type medaka. Zool. Sci (Tokyo), 6(5): 943-950.
7. Awaji, M. and I. Hanyu. 1989. Temperature-photoperiod conditions necessary to begin

- the spawning season in wild type medaka. Nippon Suisan Gakkaishi, 55(4): 747.
14. Belanger, S. E., D. S. Cherry and J. JR. Cairns. 1990. Functional and pathological impairment of Japanese Medaka (*Oryzias latipes*) by long-term asbestos exposure. Aquatic Toxicol., 17(2): 133-154.
 15. Bentivegna, C. S. and T. Piatkowski. 1998. Effects of tributyltin on medaka (*Oryzias latipes*) embryos at different stages of development. Aquatic Toxicol., 44(1-2): 117-128.
 16. Benton, M. J., S. A. Diamond and S. I. Guttman. 1994. A genetic and morphometric comparison of *helisoma trivolvis* and *gambusia holbrookii* from clean and contaminated habitats. Ecotoxicol. Environ. Saf., 29(1): 20-37.
 17. Billard, R. 1984. Ultrastructural change in the spermatogonia and spermatocytes of *Poecilia reticulata* during spermatogenesis. Cell Tissue Res., 237(2): 219-226.
 18. Brauneck, T. A., S. J. Ted, S. M. Lester and E. D. Hinton. 1992. Ultrastructural alterations in liver of medaka (*Oryzias latipes*) exposed to diethylnitrosamine. Toxicologic Pathology, 20(2): 179-196.
 19. Breder, C. M. Jr. 1954. Equations descriptive of fish schools and other animal aggregations. Ecology, 35: 361-370.
 20. Brittelli, M. R., H. H. Chen and C. F. Muska. 1985. Induction of branchial (gill) neoplasms in the medaka fish (*Oryzias latipes*) by N-methyl-N'-nitrosoguanidine. Cancer Res., 45(7): 3209-3214.
 21. Bryant, M. J. and J. W. A. Grant. 1995. Resource defense, monopolization and variation of fitness in groups of female Japanese medaka depend on the synchrony of food arrival. Anim. Behav., 49(6): 1469-1479.
 22. Bunton, T. E. 1989. Diethylnitrosamine (DEN)-induced degenerative, proliferative and neoplastic lesions in the liver of the medaka (*Oryzias latipes*) following short-term exposure. Mar. Environ. Res., 28(1-4): 369-374.
 23. Bunton, T. E. 1991. Ultrastructure of hepatic hemangiopericytoma in the medaka (*Oryzias latipes*). Exp. Mol. Pathol., 54(2): 87-98.
 24. Cameron, I. L. and K. E. Hunter. 1984. Regulation of the permeability of the medaka fish embryo chorion by exogenous sodium and calcium ions. J. Exp. Zool., 231(3): 447-454.
 25. Cameron, I. L., K. E. Hunter and W. D. Winters. 1985. Retardation of embryogenesis by extremely low frequency 60Hz electromagnetic fields. Physiol. Chem. Phys. Med. NMR., 17(1): 135-138.
 26. Cameron, I. L., W. C. Lawrence and J. B. Lum. 1985. Medaka eggs as a model system for screening potential teratogens. Prog. Clin. Biol. Res., 163C: 239-243.
 27. Cantrell, S. M., L. H. Lutz, M. Hannink and D. E. Tillitt. 1996. Embryotoxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD): the embryonic vasculature is a physiological target for TCDD-induced DNA damage and apoptotic cell death in medaka (*Oryzias latipes*). Toxicol. Appl. Pharmacol., 141(1): 23-34.
 28. Chan, K. K. 1976. A photosensitive daily rhythm in the female medaka, *Oryzias latipes*. Can. J. Zool., 54(6): 852-856.
 29. Cheer, A., J. P. Vincent, R. Nuccitelli and G. Oster. 1987. Cortical activity in vertebrate eggs. I: The activation waves. J. Theor. Biol., 124(4): 377-404.
 30. Chen Y. R., H. Uwa and X. L. Chu. 1989. Taxonomy and distribution of the genus *Oryzias* in Yunnan China Cyprinodontiforms Oryziidae. Acta. Zootaxonomica. Sinica., 14: 239-246.
 31. Chulakasem, W., J. A. Nelson and J. J. Magnuson. 1989. Interaction between effects of low pH and low ion concentration on mortality during early development of

- medaka, *Oryzias latipes*. Can. J. Zool., 67(9): 2158-2168.
32. Cooper, K. R., H. Liu, P-A. Bergovist and C. Rappe. 1991. Evaluation of baltic herring and Icelandic cod liver oil for embryo toxicity, using the Japanese medaka (*Oryzias latipes*) embryo larval assay. Environ. Toxicol. Chem., 10(6): 707-714.
33. Davis, C., M. Okihiro and D. Hinton. 2002. Effects of husbandry practices, gender, and normal physiological variation on growth and reproduction of Japanese medaka, *Oryzias latipes*. Aquat. Toxicol., 60(3-4): 185-201.
34. Dhanumkumari, C., R. K. Hanumantha and K. Shyamasundari. 1991. The life cycle of *Echinochasmus bagulai* (Trematoda: Echinostomatidae). Int. J. Parasitol., 21: 259-263.
35. Dial, N. A. 1978. Methylmercury: some effects on embryogenesis in the Japanese medaka, *Oryzias latipes*. Teratology, 17(1): 83-91.
36. Dodd, J. M. 1960. Genetic and environmental aspects of sex determination in cold blooded vertebrates. Mem. Soc. Endocrinol., 7: 17-44.
37. Egami, N. 1954. Geographical variation in the male characters of the fish *Oryzias latipes*. Annot. Zool. Japon., 27: 7-12.
38. Egami, N. 1954. Effect of artificial photoperiodicity on time of oviposition in the fish *Oryzias latipes*. Annot. Zool. Japon., 27: 57-62.
39. Egami, N. 1956. How to rear the Japanese killifish, *Oryzias latipes*. In "Handbook of Genetics", pp.851-852.
40. Egami, N. 1959. Note on sexual difference in the shape of the body in the fish, *Oryzias latipes*. Annot. Zool. Japon., 32: 59-64.
41. Egami, N. 1959. Record of the number of eggs obtained from a single pair of *Oryzias latipes*, kept in laboratory aquarium. J. Fac. Sci., Tokyo Univ. IV., 8: 521-538.
42. Egami, N. 1959. Effect of exposure to low temperature on the time of oviposition and the growth of the oocytes in the fish, *Oryzias latipes*. J. Fac. Sci., Tokyo Univ. IV., 8: 539-548.
43. Egami, N. 1969. Life span data for the small fish, *Oryzias latipes*. Exp. Gerontol., 4: 127-129.
44. Egami, N. and M. Nambu. 1961. Factors initiating mating behavior and oviposition in the fish, *Oryzias latipes*. J. Fac. Sci., Tokyo Univ., IV, 9: 263-278.
45. Etoh, H. and Y. Hyodo-Taguchi. 1983. Effects of tritiated water on germ cells in medaka embryos. Radiat. Res., 93(2): 332-339.
46. Evans, J. P., L. Zane, S. Francescato and A. Pilastro. 2003. Directional postcopulatory sexual selection revealed by artificial insemination. Nature (Lond), 421(6921): 360-363.
47. Fabacher, D. L., J. M. Besser, C. J. Schmitt, J. C. Harshbarger, P. H. Peterman and J. A. Lebo. 1991. Contaminated sediments from tributaries of the Great Lakes: chemical characterization and carcinogenic effects in medaka (*Oryzias latipes*). Arch. Environ. Contam. Toxicol., 21(1): 17-34.
48. Featherstone, D., C. D. Drewes and J. R. Coats. 1991. Noninvasive detection of electrical events during the startle response in larval medaka. J. Exp. Biol., 158: 583-589.
49. Featherstone, D., C. D. Drewes, J. R. Coats and S. P. Bradbury. 1993. A non-invasive neurotoxicity assay using larval medaka. ASTM Spec. Tech. Publ., 1216: 275-288.
50. Fluck, R. A. 1982. Localization of acetylcholinesterase activity in young embryos of the medaka *Oryzias latipes*, a teleost. Comp. Biochem. Physiol. C., 72(1): 59-64.
51. Fluck, R. A., A. L. Miller, V. C. Abraham and L. F. Jaffe. 1994. Calcium buffer

- injections inhibit ooplasmic segregation in medaka eggs. *Biol. Bull. (Woods Hole)*, 186(3): 254-262.
52. Fluck, R. A., K. L. Krok, B. A. Bast, S. E. Michaud and C. E. Kim. 1998. Gravity influences the position of dorsoventral axis in medaka fish embryos (*Oryzias latipes*). *Dev. Growth Differ.*, 40(5): 509-518.
53. Fluck, R. A. and L. F. Jaffe. 1988. Electrical currents associated with rhythmic contractions of the blastoderm of the medaka, *Oryzias latipes*. *Comp. Biochem. Physiol. A.*, 89(4): 609-613.
54. Foscarini, R. 1989. A comparative study of the skin and gill structure in oviparous and viviparous fresh water fish larvae. *J. Fish. Biol.*, 34(1): 31-40.
55. Fukamachi, S., A. Shimada and A. Shima. 2002. Positional cloning of the responsible gene for medaka orangered variants. *Zool. Sci.*, 19(12): 1420-1421.
56. Fukamachi, S., A. Shimada, K. Naruse and A. Shima. 2001. Genomic analysis of γ -ray-induced germ-cell mutations at the b locus recovered from the medaka specific-locus test. *Mutat. Res.*, 458(1-2): 19-29.
57. Fukamachi, S., H. Mitani, A. Shima and M. Sugimoto. 2004. Somatolactin selectively regulates proliferation and morphogenesis of neural-crest derived pigment cells in medaka. *Proc. Natl. Acad. Sci. U.S.A.*, 101(29): 10661-10666.
58. Fukuda, S., Y. Nagahama, N. Sakai and S. Adachi. 1994. Steroidogenesis in the ovarian follicle of medaka (*Oryzias latipes*, a daily spawner) during oocyte maturation. *Dev. Growth Differ.*, 36(1): 81-88.
59. Funayama, T., H. Mitani and A. Shima. 1993. Ultraviolet-induced DNA damage and its photorepair in tail fin cells of the medaka, *Oryzias latipes*. *Photochemistry and Photobiology*, 58(3): 380-385.
60. Furutani, M. and H. Kondoh. 2002. A genome-wide screen for mutations affecting embryonic pattern formation in the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(12): 1422.
61. Furutani, M. and J. Wittbrodt. 2004. Medaka and zebrafish, an evolutionary twin study. *Mech. Dev.*, 121(7-8): 629-637.
62. Ghoneum, M. and N. Egami. 1982. Age related changes in morphology of the thymus of the fish, *Oryzias latipes*. *Exp. Gerontol.*, 17: 33-40.
63. Ghoneum, M., N. Egami and K. Ijiri. 1988. Effect of oestrogen treatment on the survival of irradiated fish of the species *Oryzias latipes*. *Folia. Morphol (praha)*, 36(1): 25-31.
64. Gilkey, J. C. 1983. Roles of calcium and pH in activation of eggs of the medaka fish, *Oryzias latipes*. *J. Cell Biol.*, 97(3): 669-678.
65. Gonazalez-doncel, M., C. Barrueco, E. De La, Pena and D. E. Hinton. 2003. Stage sensitivity of medaka (*Oryzias latipes*) eggs and embryos to permethrin. *Aquatic Toxicol.*, 62(3): 255-268.
66. Grabher, C., T. Henrich, A. Arenz, J. Wittbrodt, T. Sasado and M. Frutani. 2003. Transposon-mediated enhancer trapping in medaka. *Gene*, 322: 57-66.
67. Grant, J. W. A. and L. D. Green. 1996. Mate copying versus preference for actively courting males by female Japanese medaka (*Oryzias latipes*). *Behav. Ecology*, 7(2): 165-167.
68. Grant, J. W. A., M. J. Bryant and C. E. Soos. 1995. Operational sex ratio, mediated by synchrony of female arrival, alters the variance of male mating success in Japanese medaka. *Anim. Behav.*, 49(2): 367-375.
69. Grant, J. W. A., P. C. Casey, M. J. Bryant and A. Shahsavari. 1995. Mate choice by Japanese medaka (Pisces, Oryziidae). *Anim. Behav.*, 50(5): 1425-1428.

70. Gray, M. A. and C. D. Metcalfe. 1999. Toxicity of 4-tert-octylphenol to early life stages of Japanese medaka (*Oryzias latipes*). *Aquatic Toxicol.*, 46(2): 149-154.
71. Gronen, S., S. Manning, S. Barnes, D. Barnes, M. Brouwer and N. Denslow. 1999. Serum vitellogenin levels and reproductive impairment of male Japanese medaka (*Oryzias latipes*) exposed to 4-tert-octylphenol. *Environ. Health Perspect.*, 107(5): 385-390.
72. Gu, Y., S. Yasumasu and I. Iuchi. 2003. 3D-structure of hce, one of Medaka hatching enzymes, and analysis of the mechanism of its egg envelope digestion using mutated recombinant hce. *Biochim. Biophys. Acta.*, 1618(2): 95-105.
73. Ha, C. R. and I. Iuchi. 1996. Participation of a metalloproteinase in the fertilization associated conversion of the egg envelope (chorion) of the fish, *Oryzias latipes*. *Dev. Growth Differ.*, 38(5): 383-396.
74. Haasch, M. L. and A. W. Ford. 2004. Combined effects of ethanol and cinnamaldehyde in the Japanese medaka embryo-larval assay (MELA). *Mar. Environ. Res.*, 58(2-5): 175-179.
75. Hamaguchi, S. 1982. A light- and electron-microscopic study on the migration of primordial germ cells in the teleost, *Oryzias latipes*. *Cell Tissue Res.*, 227(1): 139-151.
76. Hamaguchi, S. 1983. Asymmetrical development of the gonads in the embryos and fry of the fish, *Oryzias celebensis*. *Dev. Growth Differ.*, 25(6): 553-561.
77. Hamaguchi, S. 1985. Changes in the morphology of the germinal dense bodies in primordial germ cells of the teleost, *Oryzias latipes*. *Cell Tissue Res.*, 240(3): 669-673.
78. Hamaguchi, S. 1996. Bilaterally asymmetrical testes in fishes of the genus *Oryzias*. *Zool. Sci.*, 13(5): 757-763.
79. Hamaguchi, S. and M. Sakaizumi. 1992. Sexually differentiated mechanisms of sterility in interspecific hybrids between *Oryzias latipes* and *O. curvinotus*. *J. Exp. Zool.*, 263(3): 323-329.
80. Hamilton, J. B., R. O. Walter, R. M. Daniel and G. E. Mestler. 1969. Competition for mating between ordinary and supermale Japanese medaka fish. *Anim. Behav.*, 17(1): 168-176.
81. Hara, T., S. Hagino and S. Hosokawa. 2004. Quantification of vitellogenin in several developmental stages of medaka (*Oryzias latipes*) S-rR strain. *Environ. Sci. (Tokyo)*, 11(4): 221-230.
82. Hartley, W. R., A. Thiagarajah, M. B. Anderson, M. W. Broxson, S. E. Major and S. I. Zell. 1998. Gonadal development in Japanese Medaka (*Oryzias latipes*) exposed to 17β -estradiol. *Mar. Environ. Res.*, 46(1-5): 145-148.
83. Harumi, T. and T. Watanabe. 2003. Expression of membrane-bound and soluble guanylyl cyclase mRNAs in embryonic and adult retina of the medaka fish *Oryzias latipes*. *Zool. Sci.*, 20(2): 133-140.
84. Harumi, T., T. Watanabe, T. Yamamoto, Y. Tanabe and N. Suzuki. 2002. Expression of particulate and soluble guanylyl cyclase mRNA in the medaka retina. *Zool. Sci.*, 19(12): 1436.
85. Hata, K. 2002. Field experiment on the migration of fishes to paddy field with a small fishway. *JARQ.*, 36(4): 219-225.
86. Hatanaka, J., N. Doke, T. Harada, T. Aikawa and M. Enomoto. 1982. Usefulness and rapidity of screening for the toxicity and carcinogenicity of chemicals in medaka, *Oryzias latipes*. *Jpn. J. Exp. Med.*, 52(5): 243-253.
87. Hawkins, W. E., J. W. Fournie, R. M. Overstreet and W. W. Walker. 1986. Intraocular neoplasms induced by methylazoxymethanol acetate in Japanese medaka (*Oryzias latipes*). *J. Natl. Cancer Inst.*

- Inst., 76(3): 453-465.
88. Hawkins, W. E., R. M. Overstreet, J. W. Fournie and W. W. Walker. 1985. Development of aquarium fish models for environmental carcinogenesis: tumor induction in seven species. *J. Appl. Toxicol.*, 5(4): 261-264.
89. Hawkins, W. E., W. W. Walker, R. M. Overstreet, J. S. Lytle and T. F. Lytle. 1990. Carcinogenic effects of some polycyclic aromatic hydrocarbons on the Japanese medaka and guppy in waterborne exposures. *Sci. Total Environ.*, 94(1-2): 155-167.
90. Hayashi, S. and T. Ueno. 2002. Quantification of male sexual behavior of the medaka fish (*Oryzias latipes*). *Neurosci. Res.*, (Supplement 26): S84.
91. Hayashida, Y., T. Kawamura, R. Horie and I. Yamashita. 2004. Retionic acid and its receptors are required for expression of aryl hydrocarbon receptor mRNA and embryonic development of blood vessel and bone in the medaka fish, *Oryzias latipes*. *Zool. Sci.*, 21(5): 541-551.
92. Heath, A. G., J. J. Cech, J. G. Zinkl and M. D. Steele. 1993. Sublethal effects of three pesticides on Japanese medaka. *Arch. Environ. Contam. Toxicol.*, 25(4): 485-491.
93. Heisinger, J. F. and W. Green. 1975. Mercuric chloride uptake by eggs of the ricefish and resulting teratogenic effects. *Bull. Environ. Contam. Toxicol.*, 14(6): 665-673.
94. Helmstetter, M. F. and R. W. III. Alden. 1995. Passive trans-chorionic transport of toxicants in topically treated Japanese medaka (*Oryzias latipes*) eggs. *Aquatic Toxicol.*, 32(1): 1-13.
95. Helmstetter, M. F. and R. W. III. Alden. 1995. Toxic responses of Japanese medaka (*Oryzias latipes*) eggs following topical and immersion exposures to pentachlorophenol. *Aquatic Toxicol.*, 32(1): 15-29.
96. Hendreson, V. 1983. Scale morphologic response to aromatic hydrocarbons. *Bull. Environ. Contam. Toxicol.*, 31(3): 315-321.
97. Hidaka, H. and R. Tatsukawa 1985. Avoidance test of a fish, medaka (*Oryzias latipes*), to aquatic contaminants, with special reference to monochloramine. *Arch. Environ. Contam. Toxicol.*, 14(5): 565-571.
98. Hidaka, H. and R. Tatsukawa. 1989. Avoidance by olfaction in a fish, medaka (*Oryzias latipes*), to aquatic contaminants. *Environ. Pollut.*, 56(4): 299-309.
99. Higuchi, T., S. Kitagawa, H. Hori and A. Kanamori. 2002. Cloning and expression of a meiosis-specific gene, DMC1, in medaka. *Zool. Sci.*, 19(12): 1438.
100. Hinton, D. E., E. R. Walker, C. A. Pinkstaff and E. M. Zuchelkowski. 1984. Morphological survey of teleost organs important in carcinogenesis with attention to fixation. *Natl. Cancer Inst. Monogr.*, 65: 291-320.
101. Hinton, D. E., R. C. Lantz and J. A. Hampton. 1984. Effect of age and exposure to a carcinogen on the structure of the medaka liver: a morphometric study. *Natl. Cancer Inst. Monogr.*, 65: 239-249.
102. Hiramatsu, K. and S. Shikasho. 2002. Mathematical modeling of preference intensity of Japanese medaka for environmental factors in streamflow. *J. Fac. Agric. Kyusyu Univ.*, 47(1): 109-127.
103. Hiramatsu, K., S. Shikasho and K. Mori. 2000. Mathematical modeling of fish schooling of Japanese Medaka using basic behavioral patterns. *J. Fac. Agric. Kyusyu Univ.*, 45(1): 237-253.
104. Hiraoka, Y. and H. Okuda. 1983. Characteristics of vertebral abnormalities of medaka as a water pollution indicator. *Hiroshima J. Med. Sci.*, 32(3): 261-266.

105. Hiraoka, Y., S. Ishizawa, T. Kamada and H. Okuda. 1985. Acute toxicity of 14 different kinds of metals affecting medaka fry. Hiroshima J. Med. Sci., 34(3): 327-330.
106. Hiraoka, Y., J. Tanaka and H. Okuda. 1989. Toxicity to medaka of solutions of fenitrothion degraded by strong alkali. Environ. Pollut., 58(1): 35-42.
107. Hiraoka, Y., J. Tanaka and H. Okuda. 1990. Toxicity of fenitrothion degradation products to medaka (*Oryzias latipes*). Bull. Environ. Contam. Toxicol., 44(2): 210-215.
108. Hirose, K. 1972. The ultrastructure of the ovarian follicle of medaka, *Oryzias latipes*. Z. Zellforsch. Mikrosk. Anat., 123(3): 316-329.
109. Hirshfield, M., G. A. Kaufman and R. J. Beyers. 1972. Relationships of weight, length and body composition in the medaka, *Oryzias latipes*. Am. Midl. Nat., 88: 239-245.
110. Hishida, T. 1965. Accumulation of estrone-16-C-14 and diethylstilbestrolmonoethyl-1-C-14 in larval gonads of the Medaka, *Oryzias latipes*, and determination of the minimum dosage of estrogen for sex reversal. Gen. Comp. Endocrinol., 56: 137-144.
111. Hishida, T. and E. Nakano. 1954. Respiratory metabolism during fish development. Embryologia, 2: 67-79.
112. Holcombe, G. W., D. A. Benoit, D. E. Hammermeister, E. N. Leonard and D. R. Johnson. 1995. Acute and long term effects of nine chemicals on Japanese medaka (*Oryzias latipes*). Arch. Environ. Contam. Toxicol., 28(3): 287-297.
113. Hong, Y., T. Liu, H. Zhao, H. Xu, W. Wang, R. Liu, T. Chen, J. Deng and J. Gui. 2004. Establishment of a normal medakafish spermatogonial cell line capable of sperm production in vitro. Proc. Natl. Acad. Sci. U.S.A., 101(21): 8011-8016.
114. Hori, R. and K. Sato. 1973. The incorporation of 45 Ca into the egg of the medaka, *Oryzias latipes*. Experientia, 29(2): 144-145.
115. Hori, R. and T. Kono. 1974. On the content of sodium and potassium of the unfertilized egg of the medaka, *Oryzias latipes* and its changes accompanying fertilization. Protoplasma, 80(1): 149-153.
116. Hornung, M. W., P. M. Cook, K. M. Flynn, D. B. Lothenbach, R. D. Johnson and J. W. Nichols. 2004. Use of multi-photon laser-scanning microscopy to describe the distribution of xenobiotic chemicals in fish early life stages. Aquat. Toxicol., 67(1): 1-11.
117. Hosoda, K. 1958. The effect of water temperature upon the reward-type latent learning in *Oryzias latipes*. Ann. Animal Psychol., 8: 71-76.
118. Hosoda, K. 1958. The effect of water temperature on the latent learning in medakas and guppies. Japan. J. Psychol., 29: 48-50.
119. Howard, R. D., J. A. Dewoody and W. M. Muir. 2004. Transgenic male mating advantage provides opportunity for Trojan gene effect in a fish. Proc. Natl. Acad. Sci. U.S.A., 101(9): 2934-2938.
120. Howard, R. D., R. S. Martens, S. A. Innis, J. M. Drnevich and J. Hale. 1998. Mate choice and mate competition influence male body size in Japanese medaka. Anim. Behav., 55(5): 1151-1163.
121. Hyodo, M., A. Aoki, C. Ando, M. Katsumata, S. Nyui, N. Motegi, T. Morozumi and M. Matsuhashi. 1996. Essential role of the yolk syncytial layer for the development of isolated blastoderms from medaka embryos. Dev. Growth Differ., 38(4): 383-392.
122. Hyodo-Taguchi, Y. and H. Etoh. 1983. The fecundity and fertility of medaka exposed

- to chronic γ -radiation in their embryonic stages. *J. Radiat. Res.*, 24(3): 270-277.
123. Hyodo-Taguchi, Y. and H. Matsudaira. 1987. Higher susceptibility to N-methyl-N'-nitro-N-nitrosoguanidine-induced tumorigenesis in an interstrain hybrid of the fish, *Oryzias latipes* (medaka). *Jpn. J. Cancer Res.*, 78(5): 487-493.
124. Iguchi, K., M. Yoshioka, W. Wu and N. Shimizu. 1999. Biodiversity of freshwater fishes in the paddy water systems of Anji, China. *National Res. Inst. Fisheries Sci.*, 13: 27-36.
125. Iida, A., H. Inagaki, M. Suzuki, Y. Wakamatsu, H. Hori and A. Koga. 2004. The tyrosinase gene of the i(b) albino mutant of the medaka fish carries a transposable element insertion in the promoter region. *Pigment Cell Res.*, 17(2): 158-164.
126. Ijiri, K. 1980. Gamma-ray irradiation of the sperm of the fish *Oryzias latipes* and induction of gynogenesis. *J. Radiat. Res.*, 21(3-4): 263-270.
127. Ijiri, K. 1998. Development of space-fertilized eggs and formation of primordial germ cells in the embryos of Medaka fish. *Adv. Space Res.*, 21(8-9): 1155-1158.
128. Inagaki, H., A. Koga, Y. Bessho and H. Hori. 1998. The tyrosinase gene from medaka fish; transgenic expression rescues albino mutation. *Pigment Cell Res.*, 11(5): 283-290.
129. Inagaki, H., Y. Bessho, A. Koga and H. Hori. 1994. Expression of the tyrosinase-encoding gene in a colorless melanophore mutant of the medaka fish, *Oryzias latipes*. *Gene*, 150: 319-324.
130. Inamori, Y., M. Kubo, H. Tsujibo, M. Ogawa, Y. Saito, Y. Miki and S. Takemura. 1987. The ichthyotoxicity and coronary vasodilator action of 3,3'-dihydroxy-alpha,beta-diethylstilbene. *Chem. Pharm. Bull. (Tokyo)*, 35(2): 887-890.
131. Inohaya, K. and A. Kudo. 2002. Analysis of the vertebral ligament formation in living transgenic medaka. *Zool. Sci.*, 19(12): 1450.
132. Inohaya, K., A. Kudo and T. Suzuki. 2003. Analysis of the vertebral ligament formation in the teleost, *Oryzias latipes*. *Zool. Sci.*, 20(12): 1559.
133. Inoue, K., K. Naruse, N. Suzuki and Y. Takei. 2003. Four distinct C-type natriuretic peptides found in Medaka reveal new evolutionary history of the natriuretic peptide family. *Zool. Sci.*, 20(12): 1599.
134. Inoue, K. and Y. Takagi. 2001. Seawater adaptability of medaka (*Oryzias latipes*) and three related species. *Zool. Sci.*, 18(Supplement): 115.
135. Inoue, K. and Y. Takei. 2002. Diverse Adaptability in *Oryzias* Species to High Environmental Salinity. *Zool. Sci.*, 19(7): 727-734.
136. Ishikawa, T., P. Masahito and S. Takayama. 1984. Usefulness of the medaka, *Oryzias latipes*, as a test animal: DNA repair processes in medaka exposed to carcinogens. *Natl. Cancer Inst. Monogr.*, 65: 35-43.
137. Ishikawa, T. and S. Takayama. 1979. Importance of hepatic neoplasms in lower vertebrate animals as a tool in cancer research. *J. Toxicol. Environ. Health*, 5 (2-3): 537-550.
138. Ishikawa, Y. 1990. Development of muscle nerve in the teleost fish, medaka. *Neurosci. Res.*, Suppl(13): S152-S156.
139. Ishikawa, Y. 1994. Innervation of lateral line system in the medaka, *Oryzias latipes*. *The Fish Biology journal MEDAKA*, 6: 17-24.
140. Ishikawa, Y., T. Yasuda, K. Maruyama, T. Kage, N. Yamamoto, M. Yoshimoto and H. Ito. 2004. Axonogenesis in the medaka embryonic brain. *J. Neurochemistry*, 43(2-3): 481.

141. Ishikawa, Y. and Y. Hyodo-Taguchi. 1994. Cranial nerves and brain fiber systems of the medaka fry as observed by a whole-mount staining method. *Neurosci. Res.*, 19(4): 379-386.
142. Itakura, T., S. C. Mahata, M. Hirano, H. Ochi and Y. Ogino. 2002. Cytochrome P450 1A1 cDNA from freshwater teleost medaka (*Oryzias latipes*) and induction of mRNA by 3-methylcholanthrene in various organs. *Environ. Sci.*, 9(6): 463-473.
143. Ito, M., H. Nakata, O. Saitoh and M. Sugimoto. 2002. Molecular cloning of RGS proteins from the medaka (*Oryzias latipes*). *Zool. Sci.*, 19(12): 1436.
144. Ito, S. 1959. On the spawning time of the teleostean fishes. *Japanese J. Ecol.*, 9: 3.
145. Iwamatsu, T. 1965. On fertilizability of pre-ovulation eggs in the medaka, *Oryzias latipes*. *Embryologia (Nagoya)*, 61: 327-336.
146. Iwamatsu, T. 1965. Effect of acetone on the cortical changes at fertilization of the egg of the medaka, *Oryzias latipes*. *Embryologia (Nagoya)*, 9(1): 1-12.
147. Iwamatsu, T. 1994. Medaka oocytes rotate within the ovarian follicle during oogenesis. *Dev. Growth Differ.*, 36(2): 177-186.
148. Iwamatsu, T. 1998. Studies on fertilization in the teleost, I .Dynamic responses of fertilized medaka eggs. *Dev. Growth Differ.*, 40(5): 475-483.
149. Iwamatsu, T., A. Inden, K. Hirata and H. Uwa. 1984. Experiments on interspecific hybridization between *Oryzias latipes* and *Oryzias celebensis*. *Zool. Sci.*, 1(4): 653-663.
150. Iwamatsu, T., A. Yusa, H. Kobayashi, M. Yamashita and Y. Shibata. 2003. Experimental hybridization among *Oryzias* species. II . karyogamy and abnormality of chromosome separation in the cleavage of interspecific hybrids between *Oryzias latipes* and *O. javanicus*. *Zool. Sci.*, 20(11): 1381-1387.
151. Iwamatsu, T. and I. Ota. 1974. Cleavage initiating activities of sperm fractions injected into the egg of the medaka, *Oryzias latipes*. *J. Exp. Zool.*, 187(1): 3-15.
152. Iwamatsu, T., K. Onitake, K. Matsuyama, M. Satoh and S. Yukawa. 1997. Effect of micropylar morphology and size on rapid sperm entry into the eggs of the medaka. *Zool. Sci.*, 14(4): 623-628.
153. Iwamatsu, T., S. Y. Takahashi, N. Sakai, Y. Nagahama and Y. Onitake. 1987. Induction and inhibition of in vitro oocyte maturation and production of steroids in fish follicles by forskolin. *J. Exp. Zool.*, 241(1): 101-111.
154. Iwamatsu, T., T. Mori and R. A. Fluck. 1993. Mechanical dechorionation of fertilized eggs for experimental embryology in the medaka. *Zool. Sci.*, 10(6): 945-951.
155. Iwamatsu, T., T. Ohta, E. Oshima and N. Sakai. 1988. Oogenesis in the medaka *Oryzias latipes* - Stages of oocyte development. *Zool. Sci.*, 5(2): 353-373.
156. Iwamatsu, T., T. Ohta, E. Oshima and T. Sugiura. 1985. Requirement of extracellular calcium ions for early fertilization events in the medaka egg. *Dev. Growth Differ.*, 27(6): 751-762.
157. Iwamatsu, T., Y. Tada and Y. Nishiyama. 1982. On sperm nuclei and development in polyspermic eggs of the teleost, *Oryzias latipes*. *Annot. Zool. Jpn.*, 55(2): 91-99.
158. Iwamatsu, T., Y. Yoshimoto and Y. Hiramoto. 1988. Cytoplasmic Ca^{2+} release induced by microinjection of Ca^{2+} and effects of microinjected divalent cations on Ca^{2+} sequestration and exocytosis of cortiacal alveoli in the medaka egg. *Dev. Biol.*, 125(2): 451-457.
159. Iwata, A. and H. Uwa. 1981. Maintenance and growth of the horny processes of the medaka, *Oryzias latipes*, in vitro. *Dev.*

- Growth Differ., 23(3): 245-248.
160. Iwata, A., M. Iwata and E. Nakano. 1986. Distribution of fibronectin, microtubules and microfilaments in the melanophore of the medaka, *Oryzias latipes*. Cell Struct. Funct., 11(2): 99-107.
161. Jacobson, A. G. 1988. Somitomeres: mesodermal segments of vertebrate embryos. Development, 104(Suppl): 209-220.
162. Jacquet, C., V. Thermes, A. de, Luze, S. Puiseux-Dao, C. Bernard, J. S. Joly, F. Bourrat and M. Edery. 2004. Effects of microcystin-LR on development of medaka fish embryos (*Oryzias latipes*). Toxicon, 43(2): 141-147.
163. Jaffe, L. F. 1980. Calcium explosions as triggers of development. Ann. NY. Acad. Sci., 339: 86-101.
164. Jaffe, L. F. 1983. Sources of calcium in egg activation: a review and hypothesis. Dev. Biol., 99(2): 265-276.
165. Jaffe, L. F. 1986. Calcium and morphogenetic fields. Ciba. Found. Symp., 122: 271-288.
166. James, M., W. Hawkins and W. Walker. 1994. Phase 1 and phase 2 biotransformation and carcinogenicity of 2-acetylaminofluorene in medaka and guppy. Aquatic Toxicol., 28(1-2): 79-95.
167. Kanamori, A., H. Hori, K. Naruse, H. Mitani and A. Shima. 2003. Genomic organization of ZP domain containing egg envelope genes in medaka (*Oryzias latipes*). Gene, 305(1): 35-45.
168. Kanamori, A., Y. Nagahama and N. Egami. 1985. Development of the tissue architecture in gonads of the medaka *Oryzias latipes*. Zool. Sci., 2(5): 695-706.
169. Kanda, T. and Y. Itazawa. 1986. When does the group effect on oxygen consumption appear in fish? J. Fac. Agric. Kyushu Univ., 30(4): 197-207.
170. Kaneko, T. and S. Hasegawa. 1999. Application of laser scanning microscopy to morphological observations on drinking in freshwater medaka larvae and those exposed to 80% seawater. Fish Sci., 65(3): 492-493.
171. Kashiwaba, S., H. Ishikawa, N. Miyamoto, Y. Ohnishi and Y. Magara. 2002. Fish test for endocrine-disruption and estimation of water quality of Japanese rivers. Water Res., 36(8): 2161-2166.
172. Ka-Sing, Chan, K. 1977. Effect of synthetic luteinizing hormone-releasing hormone (LH-RH) on ovarian development in Japanese medaka, *Oryzias latipes*. Can. J. Zool., 55(1): 155-160.
173. Katoh, F. and T. Kaneko. 2002. Effects of environmental Ca '2+' levels on branchial chloride cell morphology in freshwater-adapted killifish *Fundulus heteroclitus*. Fish Sci., 68(2): 347-355.
174. Kaufman, G. A. and R. J. Beyers. 1972. Relationships of weight, length and body composition in the medaka, *Oryzias latipes*. Am. Midl. Nat., 88: 239-245.
175. Kaur, R., B. Buckley, K. R. Cooper, S. S. Park and Y. K. Kim. 1996. Toxicity test of nanji island landfill (Seoul, Korea) leachate using Japanese medaka (*Oryzias latipes*) embryo larval assay. Bull. Environ. Contam. Toxicol., 57(1): 84-90.
176. Kawahara, T. and I. Yamashita. 2000. Estrogen-independent ovary formation in the medaka fish, *Oryzias latipes*. Zool. Sci., 17(1): 65-68.
177. Kawakami, K., A. Shima and N. Kawakami. 2000. Identification of a functional transposase of the Tol2 element, an Ac-like element from the Japanese medaka fish, and its transposition in the zebrafish germ lineage. Proc. Natl. Acad. Sci. U.S.A., 97(21): 11403-11408.
178. Kawakami, K., K. Imanaka, M. Itoh and M. Taira. 2004. Excision of the Tol2 transposable element of the medaka fish *Oryzias latipes* in *Xenopus laevis* and

- Xenopus tropicalis*. Gene, 338(1): 93-98.
179. Kawakami, K. and T. Noda. 2004. Transposition of the Tol2 element, an Ac-like element from the Japanese medaka fish *Oryzias latipes*, in mouse embryonic stem cells. Genetics, 166(2): 895-899.
180. Kawamoto, N. 1969. Effects of androstenedione, 19-norethynyltestosterone, progesterone, and medaka, *Oryzias latipes*. Dev. Growth Differ., 11(2): 89-103.
181. Kawashima, K., A. Koga, H. Hori and A. Shima. 1998. Excision of the Tol2 transposable element of the medaka fish, *Oryzias latipes* in zebrafish, *Danio rerio*. Gene, 255(1-2): 17-22.
182. Khoo, G., V. P. E. Phang and T. M. Lim. 1992. The confocal laser scanning microscope as a tool for studying xanthophores of the swordtail (*Xiphophorus helleri*). Zool. Sci., 9(3): 665-669.
183. Kikuchi, S. and A. Imaizumi. 1983. A note on morphological changes in the thymus after rejection in the fish, *Oryzias latipes*. Zool. Mag., 92(3): 428-430.
184. Kim, I-C., Y. J. Kim, Y-D. Yoon, Y-S. Lee, J-S. Lee and S. Kawamura. 2004. Cloning of cytochrome P450 1A (CYP1A) genes from the hermaphrodite fish *Rivulus marmoratus* and the Japanese medaka *Oryzias latipes*. Mar. Environ. Res., 58(2-5): 125-129.
185. Kim, I-S and E-H. Lee. 1992. New record of ricefish, *Oryzias latipes sinensis* (Pisces Oryziidae) from Korea. Korean Journal of Systematic Zoology, 8(2): 177-182.
186. Kimura, A., M. Shinohara, R. Ohkura and T. Takahashi. 2001. Expression and localization of transcripts of MT5-MMP and its related MMP in the ovary of the medaka fish *Oryzias latipes*. Biochem. Biophys. Acta., 1518(1-2): 115-123.
187. Kinoshita, M., M. Yamauchi, M. Sasanuma, Y. Ishikawa, T. Osada, Y. Wadamatu, K. Ozato and K. Inoue. 2003. A transgene and its expression profile are stably transmitted to offspring in transgenic medaka generated by the particle gun method. Zool. Sci., 20(7): 869-875.
188. Kinoshita, M., S. Kani, K. Ozato and Y. Wakamatu. 2000. Activity of the medaka translation elongation factor 1 α -A promoter examined using the GFP gene as a reporter. Dev. Growth Differ., 42(5): 469-478.
189. Kiparissis, Y., T. L. Metcalfe, G. C. Balch and C. D. Metcalfe. 2003. Effects of antiandrogens, vi clozolin and cyproterone acetate on gonadal development in Japanese medaka (*Oryzias latipes*). Aquatic Toxicol., 63(4): 391-403.
190. Kitagawa, S., H. Hori and A. Kanamori. 2002. Expression of an oogenesis-specific gene Fig α during sex differentiation in medaka. Zool. Sci., 19(12): 1438.
191. Klaunig, J. E., B. A. Barut and P. J. Goldblatt. 1984. Preliminary studies on the usefulness of medaka, *Oryzias latipes*, embryos in carcinogenicity testing. Natl. Cancer Inst. Monogr., 65: 155-161.
192. Kleaks, P. L., J. A. Nyman and S. Bhattacharyya. 2004. Relationship between hydrocarbon measurements and toxicity to a chironomid, fish larva and daphnid for oils and oil spill chemical treatments in laboratory freshwater marsh microcosms. Environ. Pollut., 129(3): 345-353.
193. Kobayashi, H. and T. Iwamatsu. 2000. Development and fine structure of the yolk nucleus of previtellogenic oocytes in medaka *Oryzias latipes*. Dev. Growth Differ., 42(6): 623-631.
194. Kobza, R. M., J. C. Trexler, W. F. Loftus and S. A. Perry. 2004. Community structure of fishes inhabiting aquatic refuges in a threatened Karst wetland and its implications for ecosystem management. Biol. Conserv., 116(2): 153-165.

195. Koga, A., A. Shimada, M. Sakaizumi, H. Tachida and H. Hori. 2000. Evidence for recent invasion of the medaka fish genome by the Tol2 transposable element. *Genetics*, 155(1): 273-281.
196. Koga, A. and H. Hori. 1999. Homogeneity in the structure of the medaka fish transposable element Tol2. *Genet. Res.*, 73: 7-14.
197. Koga, A. and H. Hori. 2000. Detection of de novo insertion of the medaka fish transposable element tol2. *Genetics*, 156(3): 1243-1247.
198. Koga, A. and H. Hori. 2001. The Tol2 transposable element of the medaka fish: an active DNA-based element naturally occurring in a vertebrate genome. (Review), *Genes. Genet. Syst.*, 76(1): 1-8.
199. Koga, A., H. Hori and M. Sakaizumi. 2002. Transposable elements in medaka fish. *Zool. Sci.*, 19(1): 1-6.
200. Koga, A., H. Hori and M. Sakaizumi. 2002. Gene transfer and cloning of flanking chromosomal regions using the medaka fish Tol2 transposable element. *Mar. Biotechnol.*, 4(1): 6-11.
201. Koga, A., H. Inagaki, Y. Bessho and H. Hori. 1994. Body color mutants and a transposable element affecting the tyrosinase gene of the medaka fish, *Oryzias latipes*. (Review Article), *The Fish Biology Journal MEDAKA*, 6: 7-15.
202. Koga, A., H. Inagaki, Y. Bessho and H. Hori. 1995. Insertion of a novel transposable element in the tyrosinase gene is responsible for an albino mutation in the medaka fish, *Oryzias latipes*. *Mol. Gen. Genet.*, 249: 400-405.
203. Koga, A., M. Suzuki, Y. Maruyama, M. Tsutsumi and H. Hori. 1999. Amino acid sequence of a putative transposase protein of the medaka fish transposable element Tol2 deduced from mRNA nucleotide sequences. *FEBS Lett.*, 461(3): 295-298.
204. Koga, A., Y. Wakamatsu, J. Kurosawa and H. Hori. 1999. Oculocutaneous albinism in the i6 mutant of the medaka fish is associated with a deletion in the tyrosinase gene. *Pigment Cell Res.*, 12: 252-258.
205. Koizumi, N., M. Miyajima and M. Susukida. 1992. Variation of zinc and copper in metallothionein-like protein in killifish (*Oryzias latipes*) exposed to cadmium. *Chemosphere*, 24(12): 1799-1803.
206. Kondo, R., S. Kaneko, H. Sun, S. I. Chigusa and M. Sakaizumi. Diversification of olfactory receptor genes in the Japanese medaka fish, *Oryzias latipes*. *Gene*, 282(1-2): 113-120.
207. Koya, Y. and A. Iwase. 2004. Annual reproductive cycle and rate of the spermatogenic process in male mosquitofish *Gambusia affinis*. *Ichthyol. Res.*, 51(2): 131-136.
208. Koya, Y., M. Inoue, T. Naruse and S. Sawaguchi. 2000. Dynamics of oocyte and embryonic development during ovarian cycle of the viviparous mosquitofish *Gambusia affinis*. *Fish Sci.*, 66(1): 63-70.
209. Kubota, Y. and A. Shima. 1991. Comparative study of embryonic thermoresistance of two inbred strains of the medaka (*Oryzias latipes*). *J. Comp. Physiol. B. Biochem. Stst. Environ. Physiol.*, 160(6): 621-625.
210. Kubota, Y., A. Shimada and A. Shima. 1992. Detection of γ -ray-induced DNA damages in malformed dominant lethal embryos of the Japanese medaka (*Oryzias latipes*) using AP-PCR fingerprinting. *Mutat. Res.*, 283(4): 263-270.
211. Kurita, J., T. Oshiro, F. Takashima and M. Sakaizumi. 1995. Cytogenetic studies on diploid and triploid oogenesis in interspecific hybrid fish between *Oryzias latipes* and *O. ocurvinotus*. *J. Exp. Zool.*, 273(3): 234-241.
212. Kuroiwa, T., A. Ohki, K. Naka and S.

- Maeda. 1994. Biomethylation and biotransformation of arsenic in a freshwater food chain: Green alga (*Chlorella vulgaris*) fwdarw shrimp (*Neocaridina denticulata*) fwdarw killifish (*Oryzias latipes*). Applied Organometallic Chemistry, 8(4): 325-333.
213. Kurosawa, G., K. Yamada, H. Ishiguro and H. Hori. 1999. Hox gene complexity in medaka fish may be similar to that in pufferfish rater than zebrafish. Biochem. Biophys. Res. Comm., 260: 66-77.
214. Kurosawa, G., N. Takamatsu, H. Hori, K. Yamada, H. Ishiguro, M. Matuda, S. Asakawa, N. Shimizu and Y. Kohara. 2002. The genome structure and the genomic comparison of medaka hox cluster. Zool. Sci., 19(12): 1418.
215. Kuwahara, Y., A. Shimada, H. Mitani and A. Shima. 2002. A Critical stage in spermatogenesis for radiation-induced cell death in Medaka Fish, *Oryzias latipes*. Radiat. Res., 157(4): 386-392.
216. Lauren, D. J., S. J. Teh and D. E. Hinton. 1990. Cytotoxicity phase of diethylnitrosamine-induced hepatic neoplasia in medaka. Cancer Res., 50(17): 5504-5514.
217. Lee, C., J. G. NA, K-C. Lee and K. Park. 2002. Choriogenin mRNA induction in male medaka, *Oryzias latipes* as a biomarker of endocrine disruption. Aquatic Toxicol., 61(3-4): 233-241.
218. Lee, C., S. H. Jeon, N. A. J. Gyun and K. Park. 2002. Sequence analysis of choriogenin H gene of medaka (*Oryzias latipes*) and mRNA expression. Environ. Toxicol. Chem., 21(8): 1709-1714.
219. Lelewellyn, G. C., G. A. Stephenson and J. W. Hofman. 1977. Aflatoxin B1 induced toxicity and teratogenicity in Japanese Medaka eggs (*Oryzias latipes*). Toxicon, 15(6): 582-587.
220. Leung, T. S. and R. V. Bulkley. 1979. Effects of petroleum hydrocarbons on length of incubation and hatching success in the Japanese medaka. Bull. Environ. Toxicol., 23(1-2): 236-243.
221. Lindsey, C. C. 1965. The effect of alternating temperature on vertebral count in the medaka (*Oryzias latipes*). Can. J. Zool., 43: 99-104.
222. Magnuson, J. J. 1962. An analysis of the aggressive behavior, growth, and competition for food and space in medaka (*Oryzias latipes* (Pisces, Cyprinodontidae)). Can. J. Zool., 40: 313-363.
223. Martindale, M. Q., S. Meier and A. G. Jacobson. 1987. Mesodermal metamерism in the teleost, *Oryzias latipes* (the medaka). J. Morphol., 193(3): 241-252.
224. Martinez-Morales, J. R., K. Naruse, H. Mitani, A. Shima and J. Wittbrodt. 2004. Rapid chromosomal assignment of medaka mutants by bulked segregant analysis. Gene, 329: 159-165.
225. Marty, G. D., J. M. Munoz, D. J. Lauren and D. E. Hinton. 1990. Age-dependent changes in toxicity of N-nitroso compounds to Japanese medaka (*Oryzias latipes*) embryos. Aquatic Toxicol., 17(1): 45-62.
226. Marty, G. D., S. Wetzlich, J. M. Nunez, A. Craigmill and D. E. Hinton. 1991. Fish-based biomonitoring to determine toxic characteristics of complex chemical mixtures: documentation of bioremediation at a pesticide disposal site. Aquat. toxicol (Amst.), 19(4): 329-340.
227. Maruyama, K., S. Yasumasu and I. Iuchi. 2002. Analysis of globin gene expression in the medaka, *Oryzias latipes*. Zool. Sci., 19(12): 1499.
228. Maruyama, K., S. Yasumasu, I. Iuchi and K. Naruse. 2004. Globin gene organization in the medaka, *Oryzias latipes*. Zool. Sci., 18(Supplement): 27.
229. Maruyama, K., S. Yasumasu, K. Naruse, H. Mitani, A. Shima and I. Iuchi. 2004. Genomic organization and developmental expression of globin genes in the teleost

- Oryzias latipes*. Gene, 335: 89-100.
230. Matsuda, K., I. Iuchi and K. Yamagami. 1991. Analysis of hardening of the egg envelope (chorion) of the fish, *Oryzias latipes*. Dev. Growth Differ., 33(1): 75-83.
231. Matsuda, M. 2002. Identification of sex determining gene in the medaka-DMy is a Y-specific DM-domain gene required for male development. Zool. Sci., 19(12): 1421.
232. Matsuda, M., C. Matsuda, Y. Nagahama, S. Hamaguchi and M. Sakaizumi. 2001. Positional cloning of the sex-determining region of medaka using a Y congenic strain. Zool. Sci., 18(Supplement): 27.
233. Matsuda, M., H. Yonekawa, S. Hamaguchi and M. Sakaizumi. 1997. Geographic variation and diversity in the mitochondrial DNA of the medaka, *Oryzias latipes*, as determined by restriction endonuclease analysis. Zool. Sci., 14: 517-526.
234. Matsuda, M., N. Kawamoto, S. Asakawa, N. Shimizu, Y. Nagahama, S. Hamaguchi, M. Sakaizumi and H. Hori. 2001. Construction of a BCA library derived from the inbred Hd-Rr strain of the teleost fish, *Oryzias latipes*. Genes Genet. Syst., 76(1): 61-63.
235. Matsuda, M., T. Yamagishi, M. Sakaizumi and S-R. Joen. 1997. Mitochondrial DNA variation in the Korean wild population of medaka, *Oryzias latipes*. Korean J. Limnol., 30(2): 119-128.
236. Matsuda, M., Y. Nagahama, A. Shinomiya, S. Hamaguchi, M. Sakaizumi, N. Shibata, S. Asakawa, N. Shimizu and H. Hori. 2002. DMY is a Y-specific DM-domain gene required for male development in the medaka fish. Nature, 417(6888): 559-563.
237. Matsuda, M., Y. Nagahama, T. Sato, Y. Toyozaki, S. Hamaguchi and M. Sakaizumi. 2003. *Oryzias currinotus* has DMY, a gene that is required for male development in the Medaka, *O. latipes*. Zool. Sci., 20(2): 159-161.
238. Matsuzawa, T. and J. Hamilton. 1973. Polymorphism in lactate dehydrogenase of skeletal muscle associated with YY sex chromosomes in medaka (*Oryzias latipes*). Proc. Soc. Exp. Biol. Med., 142(1): 232-236.
239. Michibata, H. 1981. Effect of water hardness on the toxicity of cadmium to the egg of the teleost *Oryzias latipes*. Bull. Environ. Contam. Toxicol., 27(2): 187-192.
240. Michibata, H. and R. Hori. 1979. The accumulation of manganese from the environmental medium by the egg of *Oryzias latipes*. J. Cell Physiol., 98(1): 241-243.
241. Michibata, H., Y. Nojima and M. K. Kojima. 1987. Stage sensitivity of eggs of the teleost *Oryzias latipes* to cadmium exposure. Environ. Res. Sect. A., 42(2): 321-327.
242. Mihoko, Y. and M. Sugimoto. 2002. Is reduction of size required for the apoptosis in fish melanophores? Zool. Sci., 19(12): 1473.
243. Mori, A. 1998. Prey-handling behavior of three species of homalopsine snakes: features associated with piscivory and Duvernoy's glands. Journal of Herpetology, 32(1): 40-50.
244. Mori, S. 1970. The influence of the water volume of containers upon the fighting behavior in *Oryzias latipes*. Ann. Animal Psychol., 20: 109-114.
245. Morishita, F. 1987. Responses of the melanophores of the medaka, *Oryzias latipes*, to adrenergic drugs: evidence for involvement of alpha 2 adrenergic receptors mediating melanin aggregation. Comp. Biochem. Physiol. C., 88(1): 69-74.
246. Marshall, W. S. 2003. Rapid regulation of NaCl secretion by estuarine teleost fish: coping strategies for short-duration

- freshwater exposures. *Biochim. Biophys. Acta.*, 1618(2): 95-105.
247. Nagase, H., T. Hamasaki, T. Sato, H. Kito, Y. Yoshioka and Y. Ose. 1991. Structure-activity relationships for organotin compounds on the red killifish *Oryzias latipes*. *Appl. Organomet. Chem.*, 5(2): 91-97.
248. Nakamoto, M. and N. Shibata. 2003. Cloning and expression of daxi in Medaka gonad. *Zool. Sci.*, 20(12): 1559.
249. Nakamura, H., K. Ozato, Y. Wakamatsu and T. Iwamatsu. 2003. Normal growth of the "See-Through" medaka. *Zool. Sci.*, 20(5): 607-615.
250. Nakatsuru, Y., N. Nemoto, K. Nakagawa, P. Masahito and T. Ishikawa. 1987. O6-methylguanine DNA methyltransferase activity in liver from various fish species. *Carcinogenesis*, 8(8): 1123-1127.
251. Nakayama, K., Y. Oshima, T. Yamaguchi, Y. Tsuruda, I. J. Kang, N. Imada, T. Honjo and M. Kobayashi. 2004. Fertilization success and sexual behavior in male medaka, *Oryzias latipes*, exposed to tributyltin. *Chemosphere*, 55(10): 1331-1337.
252. Namoto, S. 1987. A subtype of adenosine receptors mediating pigment dispersion in leucophores of the medaka: evidence for an A2-receptor. *Comp. Biochem. Physiol. C.*, 88(1): 75-81.
253. Nanda, I., M. Kondo, U. Hornung, C. Winkler, M. Schartl, S. Asakawa, Z. Shan, T. Haaf and A. Shima. 2002. A duplicated copy of DMRT1 in the sex-determining region of the Y chromosome of the medaka, *Oryzias latipes*. *Proc. Natl. Acad. Sci. U.S.A.*, 99(18): 11778-11783.
254. Naruse, K., H. Hori, N. Shimizu, Y. Kohara and H. Takeda. 2004. Medaka genomics: a bridge between mutant phenotype and gene function. *Mech. Dev.*, 121(7-8): 619-628.
255. Naruse, K., H. Mitani, A. Shima, M. Tanaka and K. Mita. 2002. Medaka gene map and the vertebrate genome evolution. *Zool. Sci.*, 19(12): 1496.
256. Naruse, K., H. Mitani, S. Kondo, A. Shima and M. Tanaka. 2001. Medaka EST analysis, gene mapping and the vertebrate genome evolution. *Zool. Sci.*, 18(Supplement): 27.
257. Naruse, K., K. Ijiri, A. Shima and N. Egami. 1985. The production of cloned fish in the medaka (*Oryzias latipes*). *J. Exp. Zool.*, 236(3): 335-341.
258. Newsted, J. L., J. Nakanishi, I. Cousins, K. Werner and J. P. Giesy. 2002. Predicted distribution and ecological risk assessment of a "segregated" hidrofluoroether in the Japanese environment. *Environ. Sci. Technol.*, 36(22): 4761-4769.
259. Nimrod, A. C. and W. H. Benson. 1998. Reproduction and development of Japanese medaka following an early life stage exposure to xenoestrogens. *Aquatic Toxicol.*, 44(1-2): 141-156.
260. Nirmala, K., Y. Oshima, N. Imada, T. Honjo, K. Kobayashi and R. Lee. 1999. Transgenerational toxicity of tributyltin and its combined effects with polychlorinated biphenyls on reproductive processes in Japanese medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.*, 18(4): 717-721.
261. Niwa, Hs. 1965. Effects of castration and administration of methyl testosterone on the nuptial coloration of the Medaka (*Oryzias latipes*). *Embryologia (Nagoya)*, 61: 289-298.
262. Niwa, Hs. 1965. Inhibition by estradiol of methyl testosteroneinduced nuptial-coloration the Medaka (*Oryzias latipes*). *Embryologia (Nagoya)*, 61: 299-307.
263. Nuccitelli, R. 1980. The electrical changes accompanying fertilization and cortical vesicle secretion in the medaka egg. *Dev. Biol.*, 76(2): 483-498.

264. Nuccitelli, R. 1980. The fertilization potential is not necessary for the block to polyspermy or the activation of development in the medaka egg. *Dev. Biol.*, 76(2): 499-504.
265. Oba, Y., M. Yoshikuni, Y. Nagahama and M. Mita. 1997. Inhibitory guanine-nucleotide-binding-regulatory protein α subunits in medaka (*Oryzias latipes*) oocytes. cDNA cloning and decreased expression of proteins during oocytes maturation. *Eur. J. Biochem.*, 249(3): 846-853.
266. Ogawa, K. 1957. Social behavior in the 3-individuals group of *Oryzias latipes*. *Ann. Animal Psychol.*, 7: 72-78.
267. Ogawa, N. 1965. A phenogenetic study of the vertebral fused (f) in the medaka, *Oryzias latipes*. *Embryologia* (Nagoya), 9(1): 13-33.
268. Ogawa, Y. 1958. On the local variation in the number of anal fin-rays of Japanese killifish, *Oryzias latipes* in Tajima district, Hyogo Prefecture. *Collecting and Breeding*, 20: 48-51.
269. Ogiwara, K., M. Shinohara and T. Takahashi. 2002. Identification of the follicles digestion factor using the medaka fish in vitro ovulation system. *Zool. Sci.*, 19(12): 1547.
270. Ogiwara, K., M. Shinohara and T. Takahashi. 2004. Expression of proprotein convertase 2 mRNA in the ovarian follicles of the medaka, *Oryzias latipes*. *Gene*, 337: 79-89.
271. Oh, H. S., S. K. Lee, Y-H. Kim and J. K. Roh. 1991. Mechanism of selective toxicity of diazinon to killifish (*Oryzias latipes*) and loach (*Misgurnus anguillicaudatus*). *ASTM. Spec. Tech. Publ.*, 1124: 343-353.
272. Ohtsuka, M., N. Kikuchi, K. Ozato, H. Inoko and M. Kimura. 2004. Comparative analysis of a 229-kb medaka genomic region, containing the zic1 and zic4 genes, with Fugu, human, and mouse. *Genomics*, 83(6): 1063-1071.
273. Ohya, T. and S. Hayashi. 2003. Subnuclei structure of vasotocinergic (VT) neurons preopticus of the mature Medaka: presence of sexual dimorphism. *Neurosci. Res.*, 46(Supplement): 78.
274. Oikawa, T. 1971. Histochemical and physiological study of chrmaffin cells in the skin of the medaka, *Oryzias latipes*. *Dev. Growth Differ.*, 13(2): 125-130.
275. Okamoto, H. and J. Y. Kuwada. 1991. Outgrowth by fin motor axons in wildtype and a finless mutant of the Japanese medaka fish. *Dev. Biol.*, 146(1): 49-61.
276. Okamoto, H. and J. Y. Kuwada. 1991. Alteration of pectoral fin nerves following ablation of fin buds and by ectopic fin buds in the Japanese medaka fish. *Dev. Biol.*, 146(1): 62-71.
277. Okubo, K., K. Naruse, K. Aida, H. Mitani, M. Kondo and A. Shima. 2002. Structural and functional characterization of gonadotropin-releasing hormones and their receptors in the medaka *Oryzias latipes*. *Zool. Sci.*, 19(12): 1419-1420.
278. Okubo, K., K. Naruse, K. Aida, H. Mitani, M. Kondo, A. Shima, M. Tanaka, S. Asakawa and N. Shimizu. 2002. Structural characterization of GnRH loci in the medaka genome. *Gene*, 293(1-2): 181-189.
279. Onchi, R., Y. Koya and H. Munehara. 2002. Sperm-emission behavior by sneaker and its reproductive success in medaka fish. *Zool. Sci.*, 19(12): 1501.
280. Onitake, K. and T. Iwamatsu. 1986. Immunocytochemical demonstration of steroid hormones in the granulose cells of the medaka, *Oryzias latipes*. *J. Exp. Zool.*, 239(1): 97-103.
281. Onitsuka, S., Y. Kasai and K. Yoshimura. 1989. Quantitative structure-toxic activity relationship of fatty acids and the sodium salts to aquatic organisms. *Chemosphere*,

- 18(7-8): 1621-1631.
282. Ono, Y. and T. Uematsu. 1957. Mating ethogram in *Oryzias latipes*. J. Fac. Sci., Hokkaido Univ., VI, 13: 197-202.
283. Ono, Y. and T. Uematsu. 1958. Seasonal variation of social behavior in *Oryzias latipes*. Ann. Animal Psychol., 8: 63-69.
284. Ortego, L. S., A. C. Nimrod, G. R. Parsons, W. H. Benson and W. T. Brehm. 1994. Early life-stage effects in medaka (*Oryzias latipes*) following in ovo exposure to polyamine biosynthetic inhibitors. Ecotoxicol. Environ. Saf., 28(3): 329-339.
285. Oshima, N., H. Kasukawa, R. Fujii, B. C. Wilkes, V. J. Hruby and M. E. Hadley. 1986. Action of melanin-concentrating hormone (MCH) on teleost chromatophores. Gen. Comp. Endocrinol., 64(3): 381-388.
286. Ostrand, K. G. and G. R. Wilde. 2001. Temperature, dissolved oxygen, and salinity tolerances of five prairie stream fishes and their role in explaining fish assemblage patterns. Trans. Am. Fish Soc., 130(5): 742-749.
287. Owens, K. D. and K. N. Baer. 2000. Modifications of the topical Japanese Medaka (*Oryzias latipes*) embryo larval assay for assessing developmental toxicity of pentachlorophenol and p,p'-dichlorodiphenyltrichloroethane. Ecotoxicol. Environ. Saf., 47(1): 87-95.
288. Packer, A. 2001. Medaka on the move. Nat. Genet., 28(4): 302.
289. Parenti, L. R. and B. Soeroto. 2004. *Adrianichthys roseni* and *Oryzias nebulosus*, two new ricefishes (Atherinomorpha: Beloniformes: Adrianichtyidae) from Lake Poso, Sulawesi, Indonesia. Ichthyol. Res., 51(1): 10-19.
290. Patyna, P. J., K. R. Cooper, R. A. Davi, T. F. Parkerton and R. P. Brown. 1999. A proposed multigeneration protocol for Japanese medaka (*Oryzias latipes*) to evaluate effects of endocrine disruptors. Sci. Total Environ., 233(1-3): 211-220.
291. Philipp, D. and G. Whitt. 1977. Patterns of gene expression during teleost embryogenesis: lactate dehydrogenase isozyme ontogeny in the medaka (*Oryzias latipes*). Dev. Biol., 59(2): 183-197.
292. Rao, S. S., T. A. Neheli, C. D. Metcalfe and B. Schmidt. 1997. Assessing the Toxicity of Environmental Contaminants with Early Life Stages of Japanese Medaka (*Oryzias latipes*). Environ. Toxicol. Water Qual., 12(4): 349-352.
293. Rice, P. J., C. D. Drewes, T. M. Klubertanz, J. R. Coats and S. P. Bradbury. 1997. Acute toxicity and behavioral effects of chlorpyrifos, permethrin, phenol, strychnine, and 2,4-dinitrophenol to 30-day-old Japanese medaka (*Oryzias latipes*). Environ. Toxicol. Chem., 16(4): 696-704.
294. Ridgway, E. B., J. C. Gilkey and L. F. Jaffe. 1977. Free calcium increases explosively in activating medaka eggs. Proc. Natl. Acad. Sci. U.S.A., 74(2): 623-627.
295. Rocchetta, G., M-L. Vanelli and C. Pancaldi. 1983. Size and shape in *Poecilia reticulata*. Experientia, 39(4): 409-410.
296. Rosen, D. and L. Parenti. 1981. Relationship of *Oryzias*, and the groups of atherinomorph fishes. American Museum of Natural History, 2719: 1-25.
297. Rosenthal, G. G. and C. S. Evans. 1998. Female preference for swords in *Xiphophorus helleri* reflects a bias for large apparent size. Proc. Natl. Acad. Sci. U.S.A., 95(8): 4431-4436.
298. Rotchell, J. M., G. K. Ostrander, E. Ulnal and R. J. Van Beneden. 2001. Retinoblastoma gene mutations in chemically induced liver tumor samples of Japanese medaka (*Oryzias latipes*). Mar. Biotechnol., 3(Supplement): S44-S49.
299. Rotchell, J. M., G. K. Ostrander, J. B. Brair, J-K. Shin and W. E. Hawkins.

2001. Cloning of the retinoblastoma cDNA from the Japanese medaka (*Oryzias latipes*) and preliminary evidence of mutational alterations in chemically-induced retinoblastomas. *Gene*, 263(1-2): 231-237.
300. Ruzzante, D. E. and R. W. Doyle. 1991. Rapid behavioral changes in medaka (*Oryzias latipes*) caused by selection for competitive and non competitive growth. *Evolution*, 45: 1936-1946.
301. Ruzzante, D. E. and R. W. Doyle. 1993. Evolution of social behavior in a resource-rich, structured environment: selection experiments with medaka (*Oryzias latipes*). *Evolution*, 47: 456-470.
302. Saiki, A., M. Tamura, M. Matsumoto, J. Katowgi, A. Watanabe and K. Onitake. 1997. Establishment of in vitro spermatogenesis form spermatocytes in the medaka, *Oryzias latipes*. *Dev. Growth Differ.*, 39(3): 337-344.
303. Sakai, N., T. Iwamatsu, K. Ymauchi and Y. Nagahama. 1987. Development of the steroidogenic capacity of medaka (*Oryzias latipes*) ovarian follicles during vitellogenesis and oocyte maturation. *Gen. Comp. Endocrinol.*, 66(3): 333-342.
304. Sakaizumi, M. 1984. Rigid isolation between the northern population and the southern population of the Medaka, *Oryzias latipes*. *Zool. Sci.*, 1: 795-800.
305. Sakaizumi, M., K. Moriwaki and N. Egami. 1983. Allozymic variation and regional differentiation in wild population of the fish *Oryzias latipes*. *Copeia*, 1983: 311-318.
306. Sakaizumi, M., M. Takeda, H. Otake, K. Togashi, A. Shinomiya, S. Hamaguchi, M. Matsuda and Y. Nagahama. 2002. Mutations affecting sex determination or differentiation from wild populations of the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(12): 1497.
307. Sakaizumi, M., Y. Shimizu and S. Hamaguchi. 1992. Electrophoretic studies of meiotic segregation in inter- and intraspecific hybrids among east Asian species of the genus *Oryzias latipes* (Pisces: Oryziatidae). *J. Exp. Zool.*, 246: 85-92.
308. Sato, T., M. Sakaizumi and S. Hamaguchi. 2002. Endocrine mechanisms on onset of vitellogenesis in spring in the viviparous teleost, *Gambusia affinis*. *Zool. Sci.*, 19(12): 1487.
309. Sato, T., M. Sakaizumi and S. Hamaguchi. 2002. Genetic analysis of a secondary sexual character-less mutant, scl, the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(12): 1488.
310. Schartl, M. 2004. A comparative view on sex determination in medaka. *Mech. Dev.*, 121(7-8): 639-645.
311. Schlenk, D. and A. El-Alfy. 1998. Expression of branchial flavin-containing monooxygenase is directly correlated with salinity-induced aldicarb toxicity in the euryhaline fish (*Oryzias latipes*). *Mar. Environ. Res.*, 46(2-3) (Special Issue): 103-106.
312. Scholz, S., C. Kordes, J. Hamann and H. O. Gutzeit. 2004. Induction of vitellogenin in vivo and in vitro in the model teleost medaka (*Oryzias latipes*): comparison of gene expression and protein levels. *Mar. Environ. Res.*, 57(3): 235-244.
313. Schoots, A. F., R. C. Meijer and J. M. Denude. 1983. Dopaminergic regulation of hatching in fish embryos. *Dev. Biol.*, 100(1): 59-63.
314. Seki, M., H. Yokota, H. Matsubara, M. Maeda, H. Tadokoro and K. Kobayashi. 2004. Fish full life-cycle testing for androgen methyltestosterone on Medaka (*Oryzias latipes*). *Environ. Toxicol. Chem.*, 23(3): 774-781.
315. Sguigna, C., R. Fluck and B. Barber. 1988. Calcium dependence of rhythmic contractions of the *Oryzias latipes* blastoderm. *Comp. Biochem. Physiol. C.*,

- 89(2): 369-374.
316. Shibata, N. and S. Hmaguchi. 1988. Evidence for the sexual bipotentiality of spermatogonia in the fish, *Oryzias latipes*. *J. Exp. Zool.*, 245(1): 71-77.
317. Shima, A. and A. Shimada. 1988. Induction of mutations in male of the fish *Oryzias latipes* at a specific locus after gamma-irradiation. *Mutat. Res.*, 198(1): 93-98.
318. Shima, A. and A. Shimada. 1991. Development of a possible nonmammalian test system for radiation-induced germ-cell mutagenesis using a fish, the Japanese medaka (*Oryzias latipes*). *Proc. Natl. Acad. Sci. U.S.A.*, 88(6): 2545-2549.
319. Shima, A. and A. Shimada. 2001. The medaka as a model for studying germ-cell mutagenesis and genomic instability. *Mar. Biotechnol.*, 3(Supplement 1): S162-S167.
320. Shima, A., A. Shimada, J. Komura, K. Isa, K. Naruse, M. Sakaizumi and N. Egami. 1985. The preservation and utilization of wild populations of the medaka *Oryzias latipes*. *Medaka*, (3): 1-4.
321. Shima, A. and N. Egami. 1978. Absence of systematic polyploidization of hepatocyte nuclei during the ageing process of the male medaka, *Oryzias latipes*. *Exp. Gerontol.*, 13(1-2): 51-55.
322. Shimada, A. and A. Shima. 2004. Transgenerational genomic instability as revealed by a somatic mutation assay using the medaka fish. *Mutat. Res.*, 552(1-2): 119-124.
323. Shimada, A., A. Shima, S. Fukumachi, Y. Wakamatsu and K. Ozato. 2002. Induction and characterization of mutations at the b locus of the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(4): 411-417.
324. Shimada, Y. and N. Egami. 1984. The unique responses of the primordial germ cells in the fish *Oryzias latipes* to gamma-rays. *Int. J. Radiat. Biol. Relat. Stud. Phys. Chem. Med.*, 45(3): 227-235.
325. Shimizu, K., S. Sawaguchi, Y. Koya and A. Shimizu. 2002. Endocrine mechanisms on onset of vitellogenesis in spring in the viviparous teleost, *Gambusia affinis*. *Zool. Sci.*, 19(12): 1487.
326. Shimura, R., Y. X. Ma, K. Ijiri, S. Nagaoka and M. Uchiyama. 2004. Nitrate toxicity on visceral organs of medaka fish, *Oryzias latipes*: aiming to raise fish from egg to egg in space. *Biological Sciences in Space*, 18(1): 7-12.
327. Shinomiya, A., H. Otake, K. Togashi, S. Hamaguchi and M. Sakaizumi. 2004. Field survey of sex-reversals in the medaka, *Oryzias latipes*; genotypic sexing of wild populations. *Zool. Sci.*, 21(6): 613-619.
328. Shinomiya, A., S. Hamaguchi, M. Sakaizumi, M. Matsuda and Y. Nagahama. 2002. A strain having a ZZ/ZW sex determination system derived from a naturally occurring recessive mutation in the medaka, *Oryzias latipes*. *Zool. Sci.*, 19(12): 1497.
329. Shugart, L. R., J. F. McCarthy, S. J. D' surney, M. S. JR. Greeley and C. G. Hull. 1990. Molecular and cellular markers of toxicity in the Japanese Medaka *Oryzias latipes*. US. DOE. Rep., pp32.
330. Solomon, F. P. and E. M. Faustman. 1987. Developmental toxicity of four model alkylating agents on Japanese medaka fish (*Oryzias latipes*) embryos. *Environ. Toxicol. Chem.*, 6(10): 747-753.
331. Solomon, H. M. and J. S. Weis. 1979. Abnormal circulatory development in medaka caused by the insecticides carbaryl, malathion and parathion. *Teratology*, 19(1): 51-62.
332. Spitsbergen, J. M. 1995. Cardiovascular toxicity of environmental contaminants to developing fish-molecular mechanisms. AD. Rep., pp27.
333. Srivastava, P. N. 1966. Effect of ionizing radiation on the ovaries of Japanese medaka, *Oryzias latipes* (t. et s.). *Acta.*

- Anat (Basel), 63(3): 434-444.
334. Sugimoto, M. 1993. Morphological color changes in the medaka, *Oryzias latipes*, after prolonged background adaptaion- I. Changes in the population morphology of melanophores. Comparative biochemistry and physiology A Comparative Physiology, 104(3): 513-518.
335. Sun, H., R. Kondo, A. Shima, K. Naruse, H. Hori and S. I. Chigusa. 1999. Evolutionary analysis of putative olfactory receptor genes of medaka fish, *Oryzias latipes*. Gene, 231(1-2): 137-145.
336. Suyama, I., H. Etoh, T. Maruyama, Y. Kato and R. Ichikawa. 1981. Effects of ionizing radiation on the early development of *Oryzias* eggs. J. Radiat. Res., 22(1): 125-133.
337. Suzuki, A. and N. Shibata. 2004. Developmental process of genital ducts in the medaka, *Oryzias latipes*. Zool. Sci., 21(4): 397-406.
338. Suzuki, A., N. Shibata, T. Sato, M. Sakaizumi and S. Hamaguchi. 2002. Development of gonads and genital ducts in secondary sexual character less mutant of medaka. Zool. Sci., 19(12): 1488.
339. Suzuki, A., N. Shibata, T. Sato, M. Sakaizumi and S. Hamaguchi. 2003. Early development of gonads and genital ducts in secondary sexual character less mutant. Scl. of the Medaka. Zool. Sci., 20(12): 1604.
340. Tadokoro, H., M. Maeda, Y. Kawashima, M. Kitano, D. F. Hwang and T. Yoshida. 1991. Aquatic toxicity testing for multicomponent compounds with special reference to preparation of test solution. Ecotoxicol. Environ. Saf., 21(1): 57-67.
341. Takahashi, H. and Y. Iwasaki. 1973. Histochemical demonstration of delta 5-3 beta-hydroxysteroid dehydrogenase activity in the testis of the medaka, *Oryzias latipes*. Endocrinol. Jpn., 20(5): 529-533.
342. Takai, A., N. Kagawa and K. Fujikawa. 2004. Does- and time-dependent responses for micronucleus induction by X-rays and fast neutrons in gill cells of medaka. Environ. Mol. Mutagen., 44(2): 108-112.
343. Takai, A., N. Kagawa and K. Fujikawa. 2004. Susceptibility of male and female medaka (*Oryzias latipes*) fish to spontaneous and X-ray induced micronucleus formation in gill cells. Mutat. Res., 558(1-2): 131-136.
344. Takehana, Y., S. Uchiyama, M. Sakaizumi, M. Matsuda and S-R. Jeon. 2004. Geographic variation and diversity of the cytochrome b gene in wild population of medaka (*Oryzias latipes*) from Korea and China. Zool. Sci., 21(4): 483-491.
345. Takemoto, T., H. Kimura, M. Nonaka, A. Shimada, A. Shima and S. Yanagi. 2001. Functional investigation of the DRP/CRAM family in medaka fish, *Oryzias latipes*. Zool. Sci., 18(Supplement): 70.
346. Takemoto, Y., Y. Ohshima, H. Yamada and J. Miyamoto. 1984. Fate of fenitrothion in several developmental stages of the killifish (*Oryzias latipes*). Arch. Environ. Contam. Toxicol., 13(5): 579-587.
347. Takenaka, Y. and M. Sakaizumi. 2001. Geographic variation in the cytochrome b gene of the Japanese medaka, *Oryzias latipes*. Zool. Sci., 18(Supplement): 29.
348. Takenaka, Y. and M. Sakaizumi. 2002. Phylogeography of medaka (*Oryzias latipes*) based on mitochondrial cytochrome b DNA sequences. Zool. Sci., 19(12): 1497.
349. Takeuchi, K. 1960. A study of the mutant (wavy) in the medaka, *Oryzias latipes*. Annot. Zool. Japon., 33: 124-131.
350. Takeuchi, K. 1966. 'Wavy-fused' mutants in the Medaka, *Oryzias latipes*. Nature, 211(51): 866-867.
351. Takeuchi, K. 1967. Large tooth formation in female medaka, *Oryzias latipes*, given

- methyl testosterone. J. Dent., 46(4): 750.
352. Takeuchi, K. 1967. Normal development of teeth of medaka, *Oryzias latipes*. Aichi Gakuin Daigaku Shigakkaishi, 5(1): 81-83.
353. Takeuchi, K. 1967. Discrimination of genetic sex in embryos of d-rR strain of the medaka (*Oryzias latipes*). Experientia, 23(7): 569-570.
354. Takeuchi, K. 1968. Inhibition of large distal tooth formation in male medaka, *Oryzias latipes*, by estradiol. Experientia, 24(10): 1061-1062.
355. Takeuchi, T. and E. Manabe. 1990. Statistical analysis on the sexual behavior of the medaka, *Oryzias latipes*. Research Bulletin of Shujitsu Women's College, 20: 1-32.
356. Takimoto, Y., M. Ohshima and J. Miyamoto. 1987. Comparative metabolism of fenitrothion in aquatic organisms. I. Metabolism in the euryhaline fish, *Oryzias latipes* and *Mugil cephalus*. Ecotoxicol. Environ. Saf., 13(1): 104-117.
357. Taneda, K., T. Nishimoto and R. Abe. 1996. The sensory cue for making school in Medaka, *Oryzias latipes*. Mem. Fac. Sci. Kochi Univ. Series d Biology, 16-17(0): 69-75.
358. Taneda, K., T. Nishimoto and T. Matsuoka. 1991. The influence of neighbor fish on avoiding direction by sudden startling in medaka *Oryzias latipes*. Mem. Fac. Sci. Kochi Univ. Series d Biology, 12: 73-80.
359. Tatsuzawa, S., Y. Kano and M. Sakaizumi. 2001. The first record of two freshwater fishes, the Medaka *Oryzias latipes* and the loach *Misgurnus anguillicaudatus* from Mage-shima island: ecological, genetical, and historical views for conservation. Biogeography, 3: 89-100.
360. Teather, K., M. Harris, J. Boswell and M. Gray. 2001. Effects of acrobat MZ(R) and tattoo C(R) on Japanese medaka (*Oryzias latipes*) development and adult male behavior. Aquatic Toxicol., 51(4): 419-430.
361. Tiersch, T. R. 2001. Cryopreservation in aquarium fishes. Mar. Biotechnol., 3(Supplement1): S212-S223.
362. Tohya, S., A. Mochizuki and Y. Iwasa. 2000. A cell-sorting model can explain cone mosaic pattern of Medaka. Proc. 1st. Int. Conf. Syst., 2000: 239-244.
363. Toledo, C., J. Hendricks, P. Loveland, J. Wilcox and G. Bailey. 1987. Metabolism and DNA-binding in vivo of aflatoxin B1 in medaka (*Oryzias latipes*). Comp. Biochem. Physiol. C, 87(2): 275-281.
364. Tong, Y., T. Shan, Y. K. Poh, T. Yan, H. Wang, S. H. Lam and Z. Gong. 2004. Molecular cloning of zebrafish and medaka vitellogenin genes and comparison of their expression in response to 17beta-estradiol. Gene, 328: 25-36.
365. Toyozaki, Y., M. Sakaizumi and S. Hamaguchi. 2002. Three species closely related to *Oryzias latipes* have a male-heterogametic (XX-XY) sex-determining system. Zool. Sci., 19(12): 1496.
366. Trexler, J. C. and J. Travis. 1992. Effects of habitat and body size on mortality rates of *Poecilia Latipinna*. Ecology, 73: 2224.
367. Tsuda, T., S. Aoki, T. Inoue and M. Kojima. 1995. Accumulation of IBP, parathion and EPN by killifish: comparison of individual and mixed pesticides. Comparative Biochemistry and Physiology C Pharmacology Toxicology and Endocrinology, 111C(1): 19-22.
368. Tsuda, T., M. Kojima, H. Harada, A. Nakajima and S. Aoki. 1997. Relationships of bioconcentration factors of organophosphate pesticides among species of fish. Comparative Biochemistry and Physiology C Pharmacology Toxicology and Endocrinology, 116C(3): 213-218.
369. Tsuda, T., M. Kojima, H. Harada, A. Nakajima and S. Aoki. 1997. Acute

- toxicity, accumulation and excretion of isoprothiolane and its degradation products in killifish. Water Res., 31(2): 323-327.
370. Tsuda, T., M. Kojima, H. Arada, A. Nakajima and S. Aoki. 1997. Acute toxicity, accumulation and excretion of organophosphorous insecticides and their oxidation products in killifish. Chemosphere, 35(5): 939-949.
371. Tsukuda, H. and T. Katayama. 1957. The temperature adaptation in fishes I. The influence of rearing temperature on temperature tolerance, growth rate and body form. Physiol. and Ecol., 7: 113-122.
372. Tsuneki, K. and T. Sasaki. 1957. Fixation of poison by populations of *Oryzias latipes*. Japanese J. Ecol., 6: 153.
373. Ueda, H., G. Young and Y. Nagahama. 1983. Immunocytochemical identification of thyrotropin (TSH)-producing cells in pituitary glands of several species of teleosts with antiserum to human TSH beta subunit. Cell Tissue Res., 231(1): 199-204.
374. Uematsu, K. 1990. An analysis of sufficient stimuli for the oviposition in the medaka *Oryzias latipes*. J. Fac. Appl. Biol. Sci. Hiroshima Univ., 29(2): 109-116.
375. Urasaki, H. 1976. The role of pineal and eyes in the photoperiodic effect on the gonad of the medaka, *Oryzias latipes*. Chronobiologia, 3(3): 228-234.
376. Utida, S., S. Hatai, T. Hirano and F. I. Kamemoto. 1971. Effect of prolactin on survival and plasma sodium levels in hypophysectomized medaka *Oryzias latipes*. Gen. Comp. Endocrinol., 16(3): 566-573.
377. Uwa, H. 1965. Gynogenetic haploid embryos of the medaka (*Oryzias latipes*). Embryologia (Nagoya), 9(1): 40-48.
378. Uwa, H. 1968. Hormonal inhibitions of ethisterone-induced process formation in adult females of the medaka, *Oryzias latipes*. Embryologia (Nagoya), 10(2): 147-154.
379. Uwa, H. and L. R. Parenti. 1988. Morphometric and meristic variation in ricefishes, genus *Oryzias*: A comparison with cytogenetic data. Ichthyol. Res., 35(2): 159-166.
380. Uwa, H., R. F. Wang and Y. R. Chen. 1988. Karyotype and geographical distribution of ricefishes from Yunnan southwestern China. Japanese J. Ichthyol., 35(3): 332-340.
381. Walker, W. W., C. S. Heard, K. Lotz, T. F. Lytle, W. E. Hawkins, D. H. Barnes and R. M. Overstreet. 1989. Tumorigenic, growth, reproductive, and developmental effects in medaka exposed to bis (tri-n-butyltin) oxide. Oceans, 1989(2): 516-524.
382. Walker, W. W., C. S. Manning, R. M. Overstreet and W. E. Hawkins. 1985. Development of aquarium fish models for environmental carcinogenesis: an intermittent-flow exposure system for volatile, hydrophobic chemicals. J. Appl. Toxicol., 5(4): 255-260.
383. Walter, R., H-Y. Li, S. Kazianis, G. W. Intano and C. A. Walter. 2002. Absence of global genomic cytosine methylation pattern erasure during medaka (*Oryzias latipes*) early embryo development. Comp. Biochem. Physiol. B., 133B(4): 597-607.
384. Walter, R. and J. Hamilton. 1970. Head-up movements as an indicator of sexual unreceptivity in female medaka, *Oryzias latipes*. Anim. Behav., 18: 125-127.
385. Walter, R. and J. Hamilton. 1970. Supermales (YY sex chromosomes) and androgen-treated XY males: competition for mating with female killifish *Oryzias latipes*. Anim. Behav., 18: 128-131.
386. Watanabe, A., E. Kobayashi, T. Ogawa and K. Onitake. 1998. Fibroblast growth factor may regulate the initiation of oocytes growth in the developing ovary of

- the medaka, *Oryzias latipes*. Zool. Sci., 15(4): 531-536.
387. Weber, D. N. and R. E. Spieler. 1987. Effects of the light-dark cycle and scheduled feeding on behavioral and reproductive rhythms of the cyprinodont fish, Medaka, *Oryzias latipes*. Experientia, 43(6): 621-624.
388. Weis, J. S. and P. Weis. 2003. Is the invasion of the common reed, *Phragmites australis*, into tidal marshes of the eastern US an ecological disaster? Mar. Pollut. Bull., 46(7): 816-820.
389. Wester, P. W. 1991. Histopathological effects of environmental pollutants β -HCH and methyl mercury on reproductive organs in freshwater fish. Comp. Biochem. Physiol. C., 100(1-2): 237-239.
390. Wester, P. W. and J. H. Canton. 1986. Histopathological study of *Oryzias latipes* (medaka) after long-term β -hexachlorocyclohexane exposure. Aquatic Toxicol., 9(1): 21-45.
391. Wester, P. W., J. H. Canton, A. A. J. Van Iersel, E. I. Krajnc and H. M. A. G. Vaessen. 1990. The toxicity of bis (tri-n-butyltin) oxide (TBTO) and di-n-butyltindichloride (DBTC) in the small fish species *Oryzias latipes* (medaka) and *Poecilia reticulata* (guppy). Aquatic Toxicol., 16(1): 53-72.
392. Winkler, C., U. Hornung, M. Kondo, C. Neuner, J. Duschl, A. Shima and M. Schartl. 2004. Developmentally regulated and non-sex-specific expression of autosomal dmrt genes in embryos of the Medaka fish (*Oryzias latipes*). Mech. Dev., 121(7-8): 997-1005.
393. Wisk, J. D. and K. R. Cooper. 1990. Comparison of the toxicity of several polychlorinated dibenzo-p-dioxins and 2,3,7,8-tetrachlorodibenzofuran in embryos of the Japanese medaka (*Oryzias latipes*). Chemosphere, 20(3-4): 361-377.
394. Wisk, J. D. and K. R. Cooper. 1990. The stage specific toxicity of 2,3,7,8-tetrachlorodibeno-p-dioxin in embryos of the Japanese medaka (*Oryzias latipes*). Environ. Toxicol. Chem., 9(9): 1159-1169.
395. Wisk, J. D. and K. R. Cooper. 1992. Effect of 2,3,7,8-tetrachlorodibeno-p-dioxin on benzo(a)pyrene hydroxylase activity in embryos of the Japanese medaka (*Oryzias latipes*). Arch. Toxicol., 66(4): 245-249.
396. Wyban, J. A. 1982. Soluble peptidase isozymes of the Japanese Medaka (*Oryzias latipes*): tissue distributions and substrate specificities. Biochem. Genet., 20(9-10): 849-858.
397. Yamada, G., Y. Ogino, S. Ohta, Y. Sato, H. Ogi, H. Kato, M. Kamikawa, R. Haraguchi and K. Suzuki. 2003. Molecular analysis of external genitalia formation: morphogenesis for a copulatory organ. Biochemical J., 75(8): 695.
398. Yamagami, K. 1975. Relationship between two kinds of hatching enzymes in the hatching liquid of the medaka, *Oryzias latipes*. J. Exp. Zool., 192(1): 127-132.
399. Yamamoto, K. 1962. Origin of the yearly crop of eggs in the medaka, *Oryzias latipes*. Annot. Zool. Japon., 35: 156-161.
400. Yamamoto, T. 1956. The physiology of fertilization in the medaka (*Oryzias latipes*). Exp. Cell Res., 10(2): 387-393.
401. Yamamoto, T. 1958. Artificial induction of functional sex-reversal in genotypic females of the medaka (*Oryzias latipes*). J. Exp. Zool., 137(2): 227-263.
402. Yamamoto, T. 1959. The effects of estrone dosage level upon the percentage of sex-reversals in genetic male (XY) of the medaka (*Oryzias latipes*). J. Exp. Zool., 141: 133-153.
403. Yamamoto, T. 1961. Progenies of sex-reversal females mated with sex-reversed males in the medaka, *Oryzias latipes*. J. Exp. Zool., 146: 163-179.
404. Yamamoto, T. 1964. The problem of viability of YY zygotes in the medaka,

- Oryzias latipes*. Genetics, 50: 45-58.
405. Yamamoto, T. 1964. Linkage map of sex chromosomes in the medaka, *Oryzias latipes*. Genetics, 50: 59-64.
406. Yamamoto, T. 1965. Estriol-induced XY females of the medaka (*Oryzias latipes*) and their progenies. Gen. Comp. Endocrinol., 5(5): 527-533.
407. Yamamoto, T. 1967. Estrone-induced white YY females and mass production of white YY males in the medaka, *Oryzias latipes*. Genetics, 55(2): 329-336.
408. Yamamoto, T. 1968. Effects of 17-alpha-hydroxyprogesterone and androstenedione upon sex differentiation in the medaka, *Oryzias latipes*. Gen. Comp. Endocrinol., 10(1): 8-13.
409. Yamamoto, T., K. Takeuchi and M. Takai. 1968. Male-inducing action of androsterone and testosterone propionate upon XX zygotes in the medaka, *Oryzias latipes*. Embryologia (Nagoya), 10(2): 116-125.
410. Yamamoto, T. and N. Matsuda. 1963. Effects of estradiol and some alkylcarbonyl androstanes upon sex differentiation in the medaka. *Oryzias latipes*. Gen. Comp. Endocrinol., 3: 101-110.
411. Yano, S. and H. Kaizu. 1954. Malformation in the medaka. Collecting and Breeding, 16: 213.
412. Yasumasu, S. 2001. Active hatching enzyme, HCE, of medaka prepared by an *E. coli* expression system. Zool. Sci., 18(Supplement): 67.
413. Yasutake, J., K. Inohaya and A. Kudo. 2003. Twist functions in the Medaka vertebral column formation. Zool. Sci., 20(12): 1559.
414. Yokota, T. and T. Oishi. 1992. Seasonal change in the locomotor activity rhythm of the medaka, *Oryzias latipes*. Int. J. Biometeorol., 36: 39-44.
415. Yoshioka, H. 1962. On the effects of environmental factors upon the reproduction of fishes. I. The effects of day-length on the reproduction of the Japanese killifish, *Oryzias latipes*. Bull. Fac. Fish., Hokkaido Univ., 13: 123-136.
416. Yoshioka, H. 1966. On the effects of environmental factors upon the reproduction of fishes. III. The occurrence and regulation of refractory period in the photoperiodic response of medaka, *Oryzias latipes*. J. Hokkaido Univ. Educ., 17: 23-33.
417. Yoshizaki, K. 1981. A morphological study on the melanophores of swordtail *Xiphophorus helleri*. Nippon Suisan Gakkaishi, 47(6): 747-751.
418. Zhu, J. and L. Li. 1997. Acute toxicity of DMAH to *Oryzias latipes* as modified by some influential factors. J. Environ. Sci. (China), 9(1): 20-26.
419. 相沢直樹・氏平政伸・島本敦介・谷下一夫. 1995. 生体組織凍結過程の熱的評価. 日本機械学会通常総会講演会講演論文集, 72(Pt1): 324-325.
420. 安芸昌彦. 2004. 栗林公園の淡水魚類. 香川生物, 31: 13-22.
421. 秋月岩魚. 1999. ブラックバスがメダカを食う－日本の生態系が危ない－. 宝島社新書, 222pp.
422. 安藤正典・西村哲治・神野透人・鈴木玲央・青木康展・有薗幸司・石橋弘志・今川正良・中室克彦. 2004. 水域環境における内分泌かく乱化学物質の次世代への影響評価法確立に関する分子遺伝学的研究. 環境保全研究成果集, 2002 (1): 11.1-11.20
423. 青井 透. 2001. 研究現場からの最新報告 (6) 水の循環利用と豊かな生態系復元を目的とするビオトープ創り. 環境施設, 83: 81-87.
424. 青木 清・大木浩子. 1985. メダカの視覚と定位. 生物の科学 遺伝, 39(8): 37-38.
425. 青木 宙・隆島史夫・平野哲也. 1997. 魚類のDNA-分子遺伝学的アプローチ. 恒星社厚生閣, 467pp.

426. 新家淳治・中川喜明・荒木恵一・早川修二・金丸 豪. 1989. 化学物質の環境汚染に対する研究 水中微生物分解性試験及び魚類(ヒメダカ)急性毒性試験. 三重県環境科学センター研究報告, (9): 53-60.
427. 有薗幸司. 2003. 環境ホルモンの生態系への影響. NIRS-M, 169: 133-140.
428. 伴 幸成. 1993. 田んぼの生き物たち. 大日本図書, 35pp.
429. 大日本水産会. 2000. 藤沢メダカの学校をつくる会. SAKANA, Vol.10.
430. 大日本水産会. 2001. 藤沢メダカの学校をつくる会. SAKANA, Vol.11.
431. 土井麻記子・深津和彦・篠原亮太・花嶋正孝・樋口壮太郎・友田啓二郎. 2004. 廃棄物埋立地浸出水の短期曝露によるヒメダカ(*Oryzias latipes*)への影響. 廃棄物学会論文誌, 15(4): 294-301.
432. 土井麻記子・高村嘉幸・深津和彦・鴻野 雅一. 2004. アクアリウムを用いた生態系に対する排水の安全性評価に関する研究. 水環境学会誌, 27(5): 343-348.
433. 土井敏男. 2001. 封鎖的ビオトープで見られたメダカの体型変異. 兵庫陸水生物, (53): 73-79.
434. 江上信雄. 1953. メダカの尻鰭軟条数の変異に関する研究 I. 日本各地産野生メダカの軟条数の変異. 魚類学雑誌, 3(33-3): 87-89.
435. 江上信雄. 1953. メダカの尻鰭軟条数の変異に関する研究 II. 軟条数についての交配実験. 魚類学雑誌, 3: 171-178.
436. 江上信雄. 1989. メダカに学ぶ生物学. 中央公論社, 238pp.
437. 江上信雄・山上健次郎・嶋 昭紘. 1990. メダカの生物学. 東京大学出版会, 315pp.
438. 江上信雄・吉野道仁. 1954. メダカの尻鰭軟条数の変異に関する研究III. 野生メダカ軟条数の地理的変異(資料の追加). 魚類学雑誌, 7: 83-88.
439. 江口星雄. 1995. IML-2 “メダカの宇宙における交尾・産卵行動”実験の概要. 東京大学原子力研究総合センター技術発表会技術報告, 4: 26-33.
440. 遠藤 徹. 2004. メダカの里「家根台地区」水田生態系保全の取り組みと地域づくり. ほ場と土壤, 36(2): 29-34.
441. 藤田肖子・石嶋直之・西内康浩. 1987. 水田除草剤に対する淡水ヒメダカと海水順化ヒメダカの感受性の相違. 農薬検査所報告, (27): 32-36.
442. 藤原茂樹・山田一裕・西村 修・須藤隆一. 1999. ヒメダカに対する界面活性剤とアンモニアの複合影響. 用水と排水, 41(7): 598-602.
443. 福田博道. 2001. メダカを川に戻したい. FRONT, 7月号.
444. 福本紘一・宮下敏夫・南部和人. 2003. 水生植物の移植とビオトープ. 水草研究会誌, 79: 29-32.
445. 古川昭雄・椿 宣高・高村健二・永田尚志・可知直毅・高橋慎司・安岡善文・田村正行・渡辺 信. 1995. 地球環境研究総合推進費による 野生生物種の減少に関する研究. 国立環境研究所地球環境研究年報, 1994: 79-88.
446. 古川瑠美・野呂晋一・長尾和右・井尻憲一・江口星雄. 1999. メダカ稚魚の成長に伴う視覚の発達および微小重力下での挙動. 宇宙生物科学, 13(3): 164-165.
447. 古田寿昭. 2002. 生体系の非破壊分析(2)光で生理活性をオンーオフできるスイッチBhcケージド化合物の設計, 合成そして応用. 現代化学, 378: 24-31.
448. 古屋康則・石川早苗・沢口小有美. 2004. カダヤシ*Gambusia affinis*の卵巣周期に及ぼす水温および光周期の影響. 魚類学雑誌, 51(1): 43-50.
449. 古屋康則・恩地理惠. 2002. メダカの交接・産卵行動の観察法. 岐阜大学教育学部研究報告 自然科学, 26(2): 19-22.
450. 後出明子・坂 正臣・大西正健. 2003. 京都府下におけるメダカ生息調査と評価分析. 生物高分子, 3(1): 31-34.
451. 端 憲二. 1999. 近自然工法4 小さな魚道による休耕田への魚類遡上試験. 農業土木学会誌, 67(5): 497-502.
452. 端 憲二. 2000. 魚道の評価 田圃につける

- 小さな魚道. 応用生態工学, 3(2): 231-234.
453. 端 憲二・本間新哉・中 達男. 2001. 流速増大時のメダカの避難行動. 農業工学関係研究成果情報, 2000: 13-14.
454. 端 憲二・竹村武士・本間新哉・佐藤政良. 2001. 流れにおけるメダカの遊泳行動に関する実験的考察. 農業土木学会誌, 69(9): 987-992.
455. 畠山 仁・狩野賢司. 2001. メダカ *Oryzias latipes* の雌の配偶者選択 雄の体サイズ・体重に対する好み. 日本獣医畜産大学研究報告, 50: 25-33.
456. 日高秀雄・八反田 誠・立川 涼. 1984. メダカによる農薬の忌避. 日本農芸化学会誌, 58(2): 145-151.
457. 日野隆信・藤曲正登・福田芳生. 1981. 六価クロムの生体影響に関する研究 ヒメダカを対象とした慢性毒性試験. 千葉県衛星研究所研究報告, 5: 58-60.
458. 平松和昭・四ヶ所四男美・福田信二. 2003. ファジィ推論によるメダカの環境応答モデルの開発. 農業土木学会論文集, 228: 789-796.
459. 平松和也・山本雅照. 1998. 生態系維持回復のための水際帯調査. 大阪府立淡水魚試験場業務報告, 30: 17-19.
460. 平岡幸夫. 1980. 水質汚濁の生物指標としてのメダカのせきつい骨異常 I 某農業かんがい用水路における経時的観察所見. 広島大学医学雑誌, 28(5): 611-618.
461. 平山 真. 2003. 透明メダカ現る!. 化学と生物, 41(2): 110-111.
462. 堀 寛. 1999. 生物多様性の分子機構へのアプローチ 脊索動物ゲノムの倍数化と多様性の分子機構 トランスポゾン関与の可能性. 蛋白質 核酸 酵素, 44(3): 234-244.
463. 堀尾 剛. 2003. 農業生態系の持つ自然循環機能に基づいた食料と環境の安全性の確保 メダカを使って化学物質の内分泌かく乱作用を簡易に検出する. 農業環境研究成果情報, 19: 28-29.
464. 堀尾 剛・伊藤文成. 2001. 成育初期段階のメダカに対する内分泌かく乱物質作用に関する研究. 環境化学討論会予稿集, 10: 358-359.
465. 堀 由紀子. 1998. 水族館のはなし. 岩波書店, 200pp.
466. 細谷和海. 2000. メダカの生息状況と保護. 水環境学会誌, 23(3): 7-11.
467. 市川憲平. 1993. 雄が子守をする虫たち 5, メダカの生態(2). 海洋と生物, 15(6): 421-426.
468. 市川憲平. 2004. 放棄田ビオトープによる里の自然再生とタガメやその他の水生動物の定着. ホシザキグリーン財団研究報告, (7): 137-150.
469. 猪早敬二・工藤 明. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指して I. パターン形成と器官形成骨形成モデルとしてのメダカ. 蛋白質 核酸 酵素, 45(17): 2745-2751.
470. 石川祐二・荒木和男. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指して III. 突然変異体作製・マッピング・クローニング メダカにおけるENUおよびX線による誘発突然変異体スクリーニング. 蛋白質 核酸 酵素, 45(17): 2820-2828.
471. 岩切 稔・神野展光・長尾泰之・小寺智曉. 1994. 福岡県宗像市釣川水系産魚類Ⅳ. 宗像地域におけるメダカの分布. 福岡教育大学紀要 第三分冊 数学・理科・技術科編, (43): 65-74.
472. 岩切 稔・高尾立実・渡辺 清・中畑公宏. 1986. メダカうろこ色素胞の神経支配Ⅱ 光学顕微鏡と走査電子顕微鏡の対比による銀染色標本の観察. 福岡教育大学紀要 第三分冊 数学・理科・技術科編, 35(1985): 97-104.
473. 岩切 稔・坂中一恵・竹重宏子. 1987. メダカの形態学的体色変化. 福岡教育大学紀要 第3分冊 数学・理科・技術科編, 36(1986): 63-72.
474. 岩松鷹司. 1981. 異種精子によるメダカ (*Oryzias latipes*) 卵の発生. 愛知教育大学研究報告 自然科学, 30: 141-151.
475. 岩松鷹司. 1985. メダカ卵の実験的操作. 生物の科学 遺伝, 39(8): 42-46.
476. 岩松鷹司. 1993. メダカ学. サイエンティ

- スト社, 324pp.
477. 岩松鷹司. 1997. メダカ学全書. 大学教育出版, 370pp.
478. 岩松鷹司. 1993. メダカ卵における第一卵割面とはい軸について. 愛知教育大学研究報告自然科学, 42: 79-86.
479. 岩松鷹司. 2002. メダカと日本人. 青弓社, 216pp.
480. 和泉克雄. 1971. 热帯メダカ族百科. 東京書店, 357pp.
481. JICA. 2000. タイメダカ公開セレモニー. JICAフロンティア, 7月.
482. 城条義興. 2001. 藤沢メダカの学校をつくる会アドバイザー. 小学館, 小四教育技術, 7-8号.
483. 神奈川県公害センター. 1984. 神奈川県の水生生物. 神奈川県公害センター.
484. 神奈川県立生命の星・地球博物館. 1995. 神奈川県レッドデータ生物調査報告書, 神奈川県立博物館調査研究報告, 第7号, 神奈川県レッドデータ生物調査団編集.
485. 金森 章. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解をめざしてIV. ゲノム・進化・種分化 メダカを用いて脊椎動物の生殖巣の性分化を探る 蛋白質・核酸・酵素, 45(17): 2949-2953.
486. 金子美智雄. 2000. 田んぼに行こう里山探検をしよう, テーマ発見!総合学習体験ブック. ほるぶ出版, 40pp.
487. 環境庁. 1999. 汽水・淡水魚類のレッドリストの見直しについて. 環境庁自然保護局野生生物課, pp.1-6.
488. 環境庁自然保護局. 1997. 第4回自然環境保全基礎調査, 動植物分布調査報告書, 淡水魚類, pp.146, 342.
489. 環境教育部会. 2001. 水のしらべー藤沢の水と奏でる環境教育. 藤沢市教育文化センター.
490. 環境省. 2003. 改訂・日本の絶滅のおそれのある野生生物 - レッドデータブック (汽水・淡水魚類編). 自然環境研究センター, pp.162-163.
491. 環境省 自然環境局. 2004. 農林水産省と環境省の連携による「田んぼの生きもの調査2003」の結果について. かんきょう(東京), 29(6): 38-39.
492. 春日清一. 1980. メダカにおけるカドミウム急性毒性の雌雄差. 日本水産学会誌, 46(9): 1073-1076.
493. 川地啓文. 2001. 藤沢メダカと環境教育. ふじさわ教育, 第126号. 藤沢市教育文化センター.
494. 建設省. 1998. 平成2~7年度河川水辺の国勢調査(河川版), 生物調査結果総括. 建設省河川局河川環境課, pp.25-26.
495. 建設省. 1998. 平成2~7年度河川水辺の国勢調査(河川版), 生物調査結果総括(参考資料). 建設省河川局河川環境課.
496. 菊池久登. 1996. 再発見された藤沢メダカ. ふじさわ教育, 第108号. 藤沢市教育文化センター.
497. 北野 聰・山形哲也・市川 寛・小林 尚. 2002. 長野県北部の水田用水路における魚類群集及びメダカの推定個体数. 滋賀自然教育研究施設研究業績, 39: 17-20.
498. 北野 聰・山形哲也・市川 寛・小林 尚. 2003. 長野県北部の水田用水路に生息する淡水魚類: 耕作放棄1年後の状況. 滋賀自然教育研究施設研究業績, 40: 29-32.
499. 小林重光. 2002. メダカはいにおける脈管系の発生について. 医学検査, 51(6): 915-919.
500. 小林哲邦・菱田富雄. 1989. 卵膜除去したメダカ胚に対するメチルテストステロンの影響 特に生殖細胞数の変化に関連して. 朝日大学教養部研究報告, 15: 113-121.
501. 古賀章彦. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してIV. ゲノム・進化・種分化 メダカ属でのトランスポゾンの進化. 蛋白質・核酸・酵素, 45(17): 2900-2908.
502. 小島 昭. 2001. 炭素繊維による藻場形成榛名湖での挑戦. 水, 43(12): 27-31.
503. 小島 覚. 1999. 環境を考える・シリーズ10森林の減少と生物多様性(その3). 菌じん, 45(12): 32-38.
504. 上月康則・村上仁士・森 裕行・佐藤陽一. 1998. 都市近郊河川におけるメダカを中心とした魚類相調査. 土木学会年次学術講演会公演概要集, 53: 492-493.

505. 工藤 明. 2002. 重力研究に適した実験動物, メダカの骨形成とヒレ再生. 宇宙ステーション利用計画ワークショップ予稿集, 24(2): 133-134.
506. 草野慎司. 1996. メダカのくらし. あかね書房, 54pp.
507. Lin B-L・東海明宏・吉田喜久雄・米沢義堯・富永 衛・荻野 哲・籠島通夫・芦田昭二・岩松鷹司. 2003. メダカ (*Oryzias latipes*) 精巣卵の定量的検出のための新手法(小片化法). 水環境学会誌, 26(11): 725-730.
508. Lin B-L・東海明宏・吉田喜久雄・米沢義堯・富永 衛・荻野 哲・籠島通夫・芦田昭二・岩松鷹司. 2004. メダカ(*Oryzias latipes*) 第二次性徴による量的内分泌搅乱影響評価のための有効な手法(分節計数法). 水環境学会誌, 27(1): 47-51.
509. 松原喜代松. 1995. 魚類の形態と検索 I. 石崎書店, pp.367-370.
510. 松田 勝・長浜嘉孝. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカBACライブラリーの構築. 蛋白質 核酸 酵素, 45(17): 2880-2885.
511. 松本清志. 1995. メダカのダイアローグ. ツバサ企画.
512. 三谷啓志. 2004. メダカから脊椎動物ゲノムの痕跡をたどる フグやゼブラフィッシュとの比較から何が見えてくるか?. 化学と生物, 42(4): 221-223.
513. 宮本敦史. 2001. 水生生物の分布, 生態調査 淡水魚類の分布と希少魚類の繁殖基礎試験. 三重県科学技術振興センター水産技術センター事業報告, 2000: 93-94.
514. 宮本敦史・藤吉利彦. 2003. 水生生物の分布, 生体調査-I 淡水魚類の分布と希少魚類の繁殖基礎試験. 三重県科学技術振興センター水産研究部事業報告, 2002: 98-99.
515. 水野利恵・井尻憲一. 1999. メダカ突然変異体haの発生・成長過程における耳石の観察. 宇宙生物科学, 13(3): 166-167.
516. 森下郁子. 2000. 淡水生態系の保護と展望. 水環境学会誌, 23(3): 20-23.
517. 森下郁子. 2000. メダカを考える 淡水生態系の保護の展望. 水環境学会誌, 23(3): 148-151.
518. 森下郁子. 2001. 生態系の保全と回復に関する下水道の役割について(1). 月刊下水道, 24(7): 52-55.
519. 鍋島靖信・平松和也・上原一彦. 2003. 魚類の生息・繁殖に対する親水水路整備の効果. 大阪食とみどりの新技術, 2002: 21-22.
520. 長尾和右・井尻憲一・西脇優子. 1999. 3Dクリノスタットによる疑似微小重力下におけるメダカはい発生 網膜および軟骨形成. 宇宙生物科学, 13(3): 162-163.
521. 長屋雅人・工藤 明. 2001. 小型硬骨魚類(ゼブラフィッシュ/メダカ)を用いた再生研究. 遺伝子医学, 5(4): 653-658.
522. 中川久機・石尾真弥. 1991. 重金属イオンに対するメダカの卵, 仔魚および成魚の感受性の比較. 水産増殖, 39(4): 435-440.
523. 中川雄三. 1998. 富士のすそ野のメダカの学校(科学で環境探検). 大日本図書, 55pp.
524. 中村滝男. 1999. 空とぶメダカ 絶滅危惧種 メダカの不思議. ポプラ社, 79pp.
525. 中村滝男. 2000. メダカ保護活動の新たな展開. 水環境学会誌, 23(3): 12-15.
526. 中村滝男. 2000. メダカを考える メダカ保護活動の新たな展開. 水環境学会誌, 23(3): 140-143.
527. 成瀬 清. 1985. 人為雌性発生によるクローンメダカの作成. 生物の科学 遺伝, 39(8): 22-24.
528. 成瀬 清・近藤 周・三谷啓志・嶋昭絃・深町昌司・近藤真理子. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカ連鎖地図からみえるもの ポジショナルクローニング・比較マッピングからゲノム進化まで. 蛋白質 核酸 酵素, 45(17): 2844-2852.
529. 西原重行. 1998. メダカと共に生きて. 電算印刷株式会社, 140pp.
530. 西岡健太郎・倉田健悟・上月康則・佐良

- 家康・村上仁士・佐藤陽一・福田 守. 2000. 都市近郊用水路網内のメダカの生息環境に関する統計解析. 土木学会年次学術講演会公演概要集, 55: 18-19.
531. 西 隆行・神崎たかし・谷口三紀生・立石 康彦・岩崎英二. 2001. ビオトープの構築に関する基礎研究. 日本環境衛生センター所報, 27: 43-50.
532. 西 隆行・谷口三紀生・立石康彦・宮原 哲也. 2001. ビオトープ構築の試み. 日本環境衛生センター所報, 28: 98-103.
533. 沖津由季. 2000. 希少魚が暮らせる川づくり奮戦記6 小田原メダカがいる水田環境. 養殖, 37(8): 60-61.
534. 沖津由季・勝呂尚之. 2001. メダカを中心とした小田原市桑原・鬼柳農業用水路の魚類. 神奈川自然誌資料, 22: 51-59.
535. 小村潤一郎. 1985. メダカの細胞培養. 生物の科学 遺伝, 39(8): 28-30.
536. 大沢一爽. 1982. メダカの実験 -33章-. 共立出版. 160pp.
537. 太田克明・吉田元一・小野珠乙・山岸 哲. 1994. 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成5年度研究成果報告集2, pp.270-273.
538. 太田滋規・西森克浩・山中 治. 2003. 8. 琵琶湖沿岸帶魚類等調査費1)琵琶湖および河川下流の魚類等生息状況調査. 滋賀県水産試験場事業報告, 2002: 112-113.
539. 太田忠之・岩松鷹司. 1981. 精子べん毛微細管による魚卵の卵割誘導. 生物の科学 遺伝, 35(6): 43-45.
540. 大塚正人・木村 橙. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅢ. 突然変異体作製・マッピング・クローニング メダカ背腹構造突然変異体Daの原因遺伝子のポジショナルクローニング. 蛋白質 核酸 酵素, 45(17): 2864-2871.
541. 尾里健二郎・若松佑子・依田欣哉. 1996. メダカと遺伝子. 平成8年度社会人公開実験講座96 “バイオサイエンスを体験する” 報告書, pp.12-32.
542. 小澤祥司. 1999. 生きもの豊かな水田を再び. 水情報, 19(4): 8-11.
543. 小澤祥司. 2000. メダカが消える日-自然の再生をめざして-. 岩波書店, 224pp.
544. ピー・エス・ティー. 2004. メダカセンサー. クリモト技報, (50): 118-120.
545. 酒泉 満. 1987. メダカの分子生物地理学. 水野信彦・後藤 晃(編) 日本の淡水魚類-その分布, 変異, 種分化をめぐって-. 東海大学出版会, pp.81-90.
546. 酒泉 満. 1997. 淡水魚地方個体群の遺伝的特性と系統保存. 長田芳和・細谷和海(編) 日本の希少淡水魚の現状と系統保存, よみがえれ日本産淡水魚. 緑書房, pp.218-227.
547. 酒泉 満. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してⅣ. ゲノム・進化・種分化 メダカの系統と種内構造. 蛋白質 核酸 酵素, 45(17): 2909-2917.
548. 酒泉 満. 2000. 分子情報に基づくメダカの種内構造の解析. 日本植物分類学会会報, 15(2): 27-36.
549. 酒泉 満. 2000. メダカを考える メダカとはどういう生きものか. 水環境学会誌, 23(3): 130-134.
550. 酒泉 満. 2002. メダカの遺伝的地域差と地域個体群の保全. 日本水環境学会セミナー, 43: 90-100.
551. 酒泉 満・久保孝一. 1981. カラー自然シリーズ35. メダカ. 偕成社, 38pp.
552. 笹井隆邦. 2001. 藍那のため池の生物. 神戸常盤短期大学紀要, 22: 23-30.
553. 佐々木 昆. 1993. 小さないのちーメダカの誕生. フレーベル館, 32pp.
554. 佐藤英毅. 1988. カダヤシ, メダカ, グッピーおよびフナにおける台所用洗剤に対する感受性. 川崎市衛生研究所年報, 23(1987): 81-82.
555. 瀬能 宏. 1993. カダヤシ科・メダカ科. 中坊徹次(編) 日本産魚類検索 -全種の同定. 東海大学出版会, pp.414-415.
556. 茂岡忠義・山県民次・蓑田妙子・山内文雄. 1988. クロロフェノール類のヒメダカへの

- 急性毒性と胚のふ化阻害性及び構造との相
関. 衛生化学, 34(4): 343-349.
557. 嶋 昭絢. 2004. メダカの生物学研究最前
線. 生物の化学 遺伝, 58(5): 89-92.
558. 嶋 昭絢・桑原義和・江口星雄. 2003.
メダカ雄性殖細胞における γ 線誘発ゲノム
損傷排除機構の解析. 共用設備管理部門年
報, 30: 21-24.
559. 嶋 昭絢・島田敦子. 1995. メダカ生殖細
胞の突然変異. 共用設備管理部門総合研究
部門年報, 22: 31-34.
560. 嶋 昭絢・島田敦子. 1996. メダカ生殖細
胞の突然変異. 共用設備管理部門総合研究
部門年報, 23: 27-30.
561. 嶋 昭絢・島田敦子. 1997. メダカ生殖細
胞の突然変異. 共用設備管理部門総合研究
部門年報, 24: 21-24.
562. 島田敦子. 1985. メダカを用いた環境汚染
の検定. 生物の科学 遺伝, 39(8): 34-36.
563. 島田敦子・嶋 昭絢. 2000. メダカを用い
た生殖細胞突然変異成立機構の解析 生殖系
列における遺伝的不安定性. 放射線生物研
究, 35(4): 370-382.
564. 島田敦子・嶋 昭絢. 2000. 小型魚類研究
の新展開 脊椎動物の発生・遺伝・進化の理
解を目指してⅢ. 突然変異体作製・マッピ
ング・クローニング メダカにおける自然お
よび誘発生殖細胞突然変異率とその系統差.
蛋白質 核酸 酵素, 45(17): 2886-2892.
565. 篠原亮太. 2000. メダカ保護の意味すると
ころ. 水環境学会誌, 23(3): 1.
566. 白井康子・木村正英. 2004. 香川県のため
池の現状－希少動植物保護に向けた取り組
み－. 国立環境研究所研究報告, 183: 16-
25.
567. 杉本雅純. 1998. 魚の色と模様 色素胞の数
や形態の変化による体色変化 メダカの背地
適応を例として. 海洋と生物, 20(6): 488-
494, I.
568. 勝呂尚之. 2002. 日本産希少淡水魚の保護
水産試験場による希少淡水魚の生息地の復
元. 生物の科学 遺伝, 56(6): 76-79,
12(2)-12(3).
569. 水産庁. 1998. 日本の希少な野生水生生物
に関するデータブック. 日本水産資源保護
協会.
570. 鈴木正貴・水谷正一・後藤 章. 2000.
環境に配慮した新技術4 水田生態系保全の
ための小規模水田魚道の開発. 農業土木學
会誌, 68(12): 1263-1266, 1241(1).
571. 田部和博・安部伸治. 2003. 「自然再生推
進法」の制定と「バイオマス・ニッポン総
合戦略」の策定～2 生物多様性国家戦略と
環境省の自然再生事業について. 農業土木
学会誌, 71(8): 705-708.
572. 田口泰子. 1985. メダカの近交系の作出と
その応用. 生物の科学 遺伝, 39(8): 18-21.
573. 田口泰子. 1990. 海洋生物のジーンバンク
系統保存・凍結保存 メダカの系統保存につ
いて. 月刊海洋, 22(3): 142-148.
574. 高木勇夫. 1992. 地図に刻まれた歴史と景
観[藤沢市]. 新人物往来社, 158pp.
575. 竹田純一. 2002. 生物多様性の保全 新・生
物多様性国家戦略の実践 里地里山と生物多
様性. かんきょう (東京), 27(5): 15-16.
576. 竹花佑介・酒泉 満. 2002. 日本産希少淡
水魚の保護 メダカの遺伝的多様性の危機.
生物の科学 遺伝, 56(6): 66-71.
577. 竹村武士・奥島修二・山本勝利・小出水
規行・端 憲二. 2003. 小水路においてメ
ダカが高い選好性を示す底質条件. 農業工
学研究所研究成果情報, 2002: 17-18.
578. 竹内邦輔. 1981. 誰にでもできるメダカの
実験. 新光印刷株式会社, 110pp.
579. 竹内哲郎. 1988. めだかのお話. めだかの
学校株式会社, 18pp.
580. 多紀保彦. 2000. メダカの学校 (タイメダ
カと藤沢メダカ). 食器と容器, 6月.
- 581.. 田中宏明・高橋明宏・東谷 忠・矢古宇
靖子・斎藤正義. 2001. 下水道による水生
生態系への影響に関する調査. 国土技術政
策総合研究所資料, (10): 225-230.
582. 田中正治・平松和也. 2004. 淀川の生息魚
類の分布およびその生態. 大阪府立食とみ
どりの総合技術センター水生生物センター
研究報告, 12: 69.
583. 田中正彦. 2000. 都市の生物 生物からみた
都市の姿 メダカはどこへいった. 生物の科
学 遺伝, 54(8): 27-31.
584. 田中 実・木下政人. 2003. メダカの細胞

- ひとつひとつを光らせて生殖腺形成機構を解明する 生殖細胞系列特異的にRNA翻訳制御を指示するエレメントがわかった. 化学と生物, 41(5): 278-280.
585. 田中 実・友永健士・木下政人・若松祐子. 2003. 遺伝子機能解析のためのポストゲノム実験動物メダカ(生きたまま内臓(細胞)を見られる「透明・遺伝子導入」メダカ). 生物物理, 43(2): 93-96.
586. たたら迫典久. 2003. メダカを用いた内分泌搅乱化学物質の評価法. ファルマシア, 39(9): 881-883.
587. 寺尾 修. 1985. 野生メダカの生態. 生物の科学 遺伝, 39(8): 47-50.
588. 富森聰子・古川宏幸・長屋祐一・谷山鉄郎. 1994. ヒメダカにおよぼすゴルフ場使用農薬の複合致死毒性. 人間と環境, 20(3): 162-167, 169-171.
589. 豊泉龍児・小笠原 強. 2001. 尾(尾鰭)から肢(他の鰭)への分化転換個体の作出を目的とした、有尾両生類の肢ならびに魚類の鰭の再生現象に関する研究. 神奈川大学総合理学研究所年報, 2000: 179-181.
590. 椿 宜高. 2000. 現代社会と遺伝学 生物多様性の危機と遺伝学. 生物の科学 遺伝, 54(10): 58-64.
591. 椿 宜高・高村健二・永田尚志・富山清升・太田克明・吉田元一・山岸 哲・河田雅圭・土田浩治. 1996. 野生生物の種の減少 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成7年度終了研究成果報告書4, pp.271-287.
592. 椿 宜高・高村健二・富山清升・太田克明・吉田元一・山岸 哲・柏谷栄一・河田雅圭・伊沢雅子. 1995. 野生生物の種の減少 希少野生動物の遺伝的多様性とその保存に関する研究 希少野生動物の遺伝的多様性に関する研究 遺伝的多様性減少の影響の解明. 地球環境研究総合推進費 平成6年度研究成果報告書2, pp.280-284.
593. 辻井要介・上田哲行. 2003. コンクリート化された水路におけるメダカの分布とそれに影響を及ぼす環境要因について. 環動昆, 14(3): 179-192.
594. 上田成一・宮崎憲明・白井玄爾・山口道雄. 1995. 化学物質の魚類急性毒性および藻類生長阻害. 長崎県衛生公害研究所報, (40): 134, 135, 135(1).
595. 上山尊士. 1998. 生態影響試験業務. TRC News, 62: 13-16.
596. 氏平政伸・島本敦介・青木勝敏・相沢直樹・谷下一夫. 1995. メダカ受精卵の細胞外凍結における有効熱伝導率の測定. 日本機械学会熱工学部門講演会講演論文集, 1995: 117-119.
597. 氏平政伸・島本敦介・岡浩太郎・谷下一夫. 1996. 生体凍結過程における有効熱伝導率測定. バイオエンジニアリング学術講演会講演論文集, 7: 162-163.
598. 氏平政伸・山口 亮・相沢直樹・谷下一夫. 1995. mmオーダの生体組織の細胞外凍結による損傷. 日本機械学会論文集, 61(588): 3066-3074.
599. 氏平政伸・山口 亮・谷下一夫. 1995. mmオーダの生体組織の解凍過程に関する研究: 解凍後の顕微鏡観察によるメダカ受精卵の形態保存. 日本伝熱シンポジウム講演論文集, 32(1): 99-100.
600. 梅津 剛. 2002. 微生物からの生態系を考えた海水魚養殖装置の開発. 養殖, 39(7): 84-87.
601. 宇根 豊. 1999. 田んぼのエコロジー 田んぼの虫たちは, 自然の生きもの. 生物の科学 遺伝, 53(4): 26-30.
602. 宇根 豊. 2000. いのちが集まる・命が育む「田んぼの学校」入学編. 農村環境整備センター, pp.38-41.
603. 牛野 正. 2004. 農業土木とランドスケープエコロジー3 スイスの景観コンセプト. 農業土木学会誌, 72(1): 27-31, 1(1).
604. 宇和ひろし. 1985. メダカ属の種と系統. 生物の科学 遺伝, 39(8): 6-11.
605. 和田克彦・小林敬典・細谷和海・名古屋博之・前川光司・木曾克裕・小野里 勉・高田啓介・柴田直樹. 1995. 希少魚の個体群維持機構の解明と保存技術開発に関する予備的研究. 地球環境研究総合推進費 平成6年度研究成果報告書2, pp.344-348.

606. 若松佑子・丹羽勝利・可児修一・尾里健二郎. 2000. 小型魚類研究の新展開 脊椎動物の発生・遺伝・進化の理解を目指してV. 新技術の展開 メダカの核移植. 蛋白質核酸酵素, 45(17): 2962-2966.
607. 若松佑子・尾里健二郎. 1998. メダカの地域集団 近交系 突然変異系統. 蛋白質核酸 酵素, 43(11): 1487-1491.
608. 渡辺かほり. 2000. “藤沢メダカ”復活作戦. 養殖, 6月.
609. 渡辺かほり. 2000. メダカと環境教育「藤沢メダカの学校をつくる会」の活動から. 水環境学会誌, 23(3): 16-19.
610. 渡辺かほり. 2001. 藤沢の自然No.26, 鶴沼の自然. 藤沢市環境部みどり課.
611. 渡辺克己. 1994. 神奈川県産野生メダカの系統維持と教材化. 神奈川県立教育センター.
612. 渡辺 志. 1992. モノアルキルフォスフェイト(MAP)のヒメダカ魚毒性に及ぼすアルミニウムイオンの影響. 合成洗剤研究会誌, 16(1): 13-14.
613. 篠本美孝・上野輝彌. 1984. メダカ *Oryzias latipes* の骨学的研究. 北九州市立自然史博物館誌, 5: 143-161.
614. 山口 亮・氏平政伸・相沢直樹・谷下一夫. 1995. 生体組織の細胞外凍結による損傷の評価. 日本伝熱シンポジウム講演論文集, 32(1): 97-98.
615. 山本時男 (編). 1975. メダカの生物学と品種. 佑学社.
616. 山本義和・井上雅代・小林けい子・永井 京子. 1982. ヒメダカに対するカドミウムの急性毒性に及ぼす水温の影響. 生態化学, 5(1): 31-35.
617. 山本義和・井上雅代・小林けい子・永井 京子. 1982. ヒメダカに対するカドミウムの急性毒性に及ぼす水温の影響 特に死個体について. 生態化学, 5(2): 15-20.
618. 横山 正. 1987. ポプラ社の実験観察シリーズ3, メダカのそだちかた. ポプラ社. 39pp.
619. 吉家豊洋・森上義考. 1987. そだててみよう4, メダカ. あかね書房.
620. 吉岡敏行・萩野泰夫・森 忠繁. 1995. 水生生物に対する界面活性剤および衣料洗剤の毒性評価. 全国公害研会誌, 20(3): 138-143.
621. 吉岡義正・小瀬洋喜・後藤幹保・ヒキノヒロシ. 1987. 主として水棲の動物及び植物を用いた生態影響試験法間の関係. 衛生化学, 33(1): 11-19.
622. 吉崎正巳・太田一男. 1983. どきどき・自然, めだか. 福音館書店, 27pp.